

NON-HOMOGENEOUS BIQUADRATIC EQUATION WITH FOUR UNKNOWNNS

$$xy(x + y) + 4zw^3 = 0$$

G. DHANALAKSHMI\*<sup>1</sup>, M. A. GOPALAN<sup>2</sup>

<sup>1</sup>Assistant professor, Department of Mathematics,  
Chidambaram pillai college for women, Mannachanallur, Trichy. India.

<sup>2</sup>Professor, Department of Mathematics,  
Shrimati Indira Gandhi College, Trichy-620002, India.

(Received On: 29-10-21; Revised & Accepted On: 10-11-21)

---

ABSTRACT

The non-homogeneous bi-quadratic equation with four unknowns given by  $xy(x + y) + 4zw^3 = 0$  is analysed for finding its non-zero distinct integer solutions through different methods. A few interesting relations among the solutions are presented.

**Keywords:** Non- homogeneous bi-quadratic, Bi-quadratic with four unknowns, Integer solutions.

---

INTRODUCTION

The theory of Diophantine equations offers a rich variety of fascinating problems. In particular, biquadratic diophantine equations, homogeneous and non-homogeneous have aroused the interest of numerous mathematicians since antiquity [1-5]. In this context, one may refer [6-17] for various problems on the biquadratic diophantine equations with four variables. However, often we come across non-homogeneous biquadratic equations and as such one may require its integral solution in its most general form.

It is towards this end, this paper concerns with the problem of determining non-trivial integral solutions of the non-homogeneous equation with four unknowns given by  $xy(x + y) + 4zw^3 = 0$

NOTATIONS

$t_{m,n}$  : polygonal number of rank n with size m.

$P_m^n$  : pyramidal number of rank n with size m.

METHOD OF ANALYSIS

The non- homogeneous bi-quadratic equation with four unknowns to be solved is

$$xy(x + y) + 4zw^3 = 0 \tag{1}$$

Substitution of the linear transformations

$$x = u + v, y = u - v, z = u, (u \neq v \neq 0) \tag{2}$$

in (1) leads to

$$v^2 - u^2 = 2w^3 \tag{3}$$

---

**Corresponding Author: G. Dhanalakshmi\*<sup>1</sup>,**  
**<sup>1</sup>Assistant professor, Department of Mathematics,**  
**Chidambaram pillai college for women, Mannachanallur, Trichy. India.**

Solving (3) through different ways for  $u, v, w$  and using (2), one obtains different sets of solutions to (1) that are illustrated below:

**WAY 1:**

Write (3) as the system of double equations as shown in Table 1 below:

**Table-1: System of Double Equations**

SYSTEM	1	2	3	4	5	6
v+u	$w^3$	$2w^2$	$w^2$	$2w$	$w$	2
v-u	2	$w$	$2w$	$w^2$	$2w^2$	$w^3$

Solving each of the system of equations in Table 1, the corresponding values of  $u, v$  and  $w$  are obtained. Substituting the values of  $u$  and  $v$  in (2), the respective values of  $x$  and  $y$  are determined. The integer solutions to (1) obtained through solving each of the above system of equations are exhibited.

Solutions through SYSTEM 1:  $x = 8k^3, y = -2, z = 4k^3 - 1, w = 2k$

**OBSERVATIONS:**

- $x + w^3 = 16P_k^6$
- $x^3 + y^3 + 6xyz = 8z^3$
- $z + w = 6P_k^4 + y + 1$
- $x - w = 6P_k^8$
- $2x - 5w = 6P_k^{16}$

Solutions through SYSTEM 2:  $x = 8k^2, y = -2k, z = 4k^2 - k, w = 2k$

**OBSERVATIONS:**

- $y^3 + 12P_k^4 = 2w$
- $xy + 5w + 6P_k^{16} = 0$
- $x + w^3 + 2k = 6P_k^4 + 2z$
- $3w + 2t_{10,k} = x$
- $2x + 7y = 2t_{18,k}$

Solutions through SYSTEM 3:  $x = 4k^2, y = -4k, z = 2k^2 - 2k, w = 2k$

**OBSERVATIONS:**

- $3(z-w-y)$  is a nasty number.
- $z = 2k^2 + w + y$
- $x - w = 2t_{6,k}$
- $5y^2 = 32t_{7,k} + 24w$
- $2z + w^2 = 8t_{3,k-1} + x$
- $7w^3 + 8y = 24P_k^{14}$

Solutions through SYSTEM 4:  $x = 4k, y = -4k^2, z = 2k - 2k^2, w = 2k$

**OBSERVATIONS:**

- $13w^3 - 14x = 48P_k^{13}$
- $3y + 2x + 4t_{8,k} = 0$
- $2y + 3w + 2t_{10,k} = 0$
- $2z + w^3 = 12P_k^4 + y$
- $x + y + 8t_{3,k-1} = 0$

Solutions through SYSTEM 5:  $x = 2k, y = -8k^2, z = k - 4k^2, w = 2k$

**OBSERVATIONS:**

- $z^2 + w + 6P_k^8 = x^4 + k^2$
- $6(z^2 + w - x^4 + 6P_k^8)$  is a nasty number.
- $yw + 5x + 6P_k^{16} = 0$
- $5y + 12w + 16t_{7,k} = 0$

Solutions through SYSTEM 6:  $x = 2, y = -8k^3, z = 1 - 4k^3, w = 2k$

**OBSERVATIONS:**

- $y + 4w + 6P_k^8 = 6k$
- $w + 1 = z + 6P_k^4$
- $y + 8P_k^6 = 0$
- $x + y + 6P_k^5 + 2P_k^9 = 2$
- $y + z + 6P_k^{12} + 6k = 1$

**WAY 2:**

Taking the linear transformations

$$x = u + w, y = u - w, z = u, (u \neq w \neq 0) \tag{4}$$

in (1), it is written as

$$u^2 = w^2(1 - 2w) \tag{5}$$

After some algebra, it is seen that (5) is satisfied by the following two choices of u and w:

Choice 1:  $w = -2(k^2 + k), u = 2(k^2 + k)(2k + 1)$

Choice 2:  $w = -2(k^2 + k), u = -2(k^2 + k)(2k + 1)$

In view of (4), the corresponding sets of integer solutions to (1) are represented below:

Solutions to choice 1:

$$\begin{aligned} x &= 4k(k^2 + k) \\ y &= 4(k^2 + k)(k + 1) \\ z &= 2(k^2 + k)(2k + 1) \\ w &= -2(k^2 + k) \end{aligned}$$

Solutions to choice 2:

$$x = -4(k^2 + k)(k + 1)$$

$$y = -4k(k^2 + k)$$

$$z = -2(k^2 + k)(2k + 1)$$

$$w = -2(k^2 + k)$$

## CONCLUSION

In this paper, an attempt has been made to determine non-zero distinct integer solutions to the non-homogeneous bi-quadratic equation with four unknowns given by  $xy(x + y) + 4zw^3 = 0$ . The researchers in this field may search for other sets of integer solutions to the equation under consideration and other forms of bi-quadratic equations with four or more variables.

## REFERENCES

1. L.E. Dickson, History of Theory of Numbers, Vol.11, Chelsea publishing company, New York (1952).
2. L.J. Mordell, Diophantine equations, Academic press, London (1969).
3. Carmichael, R.D., The theory of numbers and Diophantine Analysis, Dover publications, New York (1959).
4. Telang, S.G., Number theory, Tata Mc Graw Hill publishing company, New Delhi (1996)
5. Nigel, D. Smart, The Algorithmic Resolutions of Diophantine Equations, Cambridge University press, London (1999).
6. M.A. Gopalan, V. Pandichelvi On the Solutions of the Biquadratic equation  $(x^2 - y^2)^2 = (z^2 - 1)^2 + w^4$  paper presented in the international conference on Mathematical Methods and Computation, Jamal Mohammed College, Tiruchirappalli, July 24-25, 2009.
7. M.A. Gopalan, P. Shanmuganandham, On the biquadratic equation  $x^4 + y^4 + z^4 = 2w^4$ , Impact J.Sci tech; Vol.4, No.4, 111-115, 2010.
8. M.A. Gopalan, G. Sangeetha, Integral solutions of Non-homogeneous Quartic equation  $x^4 - y^4 = (2\alpha^2 + 2\alpha + 1)(z^2 - w^2)$ , Impact J.Sci Tech; Vol.4 No.3, 15-21, 2010.
9. M.A. Gopalan, R. Padma, Integral solution of Non-homogeneous Quartic equation  $x^4 - y^4 = z^2 - w^2$ , Antarctica J. Math., 7(4), 371-377, 2010.
10. M.A. Gopalan, P. Shanmuganandham, On the Biquadratic Equation  $x^4 + y^4 + (x + y)z^3 = 2(k^2 + 3)^n w^4$ , Bessel J. Math., 2(2) 87-91, 2012.
11. M.A. Gopalan, S. Vidhyalakshmi, K. Lakshmi, On the bi-quadratic equation with four unknowns  $x^2 + xy + y^2 = (z^2 + zw + w^2)^2$ , IJPAMS, 5 (1), 73-77, 2012.
12. M.A. Gopalan, B. Sivakami, Integral solutions of Quartic equation with four unknowns  $x^3 + y^3 + z^3 = 3xyz + 2(x + y)w^3$ , Antartica J. Math., 10 (2), 151-159, 2013.
13. M.A. Gopalan, S. Vidhyalakshmi, A. Kavitha, Integral solutions to the bi-quadratic equation with four unknowns  $(x + y + z + w)^2 = xyzw + 1$ , IOSR, Vol.7(4), 11-13, 2013.
14. K. Meena, S. Vidhyalakshmi, M.A. Gopalan, S. Aarthi Thangam, On the bi-quadratic equation with four unknowns  $x^3 + y^3 = 39zw^3$ , International Journal of Engineering Research Online, Vol. 2(1), 57-60, 2014.
15. M.A. Gopalan, V. Sangeetha, Manju Somanath, Integer solutions of non-homogeneous biquadratic equation with four unknowns  $4(x^3 - y^3) = 31(k^2 + 3s^2)zw^2$ , Jamal Academic Research Journal, Special Issue, 296-299, 2015.
16. R.Anbuselvi and K.S.Araththi, "On the Biquadratic Equations with four unknowns  $x^4 - y^4 = 10(z+w) p^2$ ", International journal of Emerging Technologies in computational and Applied Sciences, 25(1), Pp :13-16, June – August 2018.
17. A.Vijayasankar, Sharadha Kumar, M.A.Gopalan, "On the Non-Homogeneous Bi-Quadratic Equation with Four Unknowns  $8xy + 5z^2 = 5w^4$ ", Jouranal of Xi'an University of architecture & Technology, 12(2), 2020, Pp:1108-1115.

**Source of support: Nil, Conflict of interest: None Declared.**

**[Copy right © 2021. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**