

**CERTAIN SUBCLASS OF MULTIVALENT  $\alpha$  - SPIRALLIKE FUNCTIONS**

**O. KARTHIYAYINI<sup>1</sup> AND V. SIVASANKARI\*<sup>2</sup>**

<sup>1,2</sup>Department of Science and Humanities,  
 PES University, Electronic City Campus, Bangalore - 560100, Karnataka, India.

(Received On: 27-12-21; Revised & Accepted On: 21-01-22)

**ABSTRACT**

The aim of this paper is to study certain subclasses of multivalent  $\alpha$ - spiral starlike functions and  $\alpha$ - spiral convex functions defined by subordination. We obtain an upper bound estimate for the second Hankel determinant of functions belonging to these classes using Toeplitz determinants. Also, the bounds rendered in this paper generalize some previous results.

**Keywords and Phrases:** Analytic function, convex  $\alpha$  – spiral function, multivalent function,  $\alpha$ - spiral starlike function, Hankel determinant, Toeplitz determinants.

**2000 Mathematics Subject Classification:** Primary 30C45.

**1. INTRODUCTION**

For a fixed integer  $p \geq 1$ , let  $A_p$  denote the class of functions  $f$  of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n, \tag{1.1}$$

which are analytic in the open unit disc  $\mathbb{U} = \{z: |z| < 1\}$  with  $p \in N = \{1,2,3, \dots\}$ .

Let  $S$  be the subclass of  $A_1 =: A$ , consisting of univalent functions.

Let  $\Omega$  be the class of Schwarzian functions

$$w(z) = \sum_{n=p+1}^{\infty} d_n z^n,$$

which are analytic in the open unit disc  $\mathbb{U} = \{z: |z| < 1\}$  and satisfies the conditions  $w(0) = 0$  and  $|w(z)| < 1$ .

Let  $f$  and  $g$  be analytic functions in  $\mathbb{U}$ , we say that  $f$  is subordinate to  $g$ , written as  $f < g$  if there exist a Schwarz function  $w \in \Omega$ , such that  $f(z) = g(w(z))$ , ( $z \in \mathbb{U}$ ) [2].

In 1976, Noonan and Thomas [18] defined the  $q^{th}$  Hankel determinant of  $f$  given by (1.1) for integers  $n \geq 1$  and  $q \geq 1$  by

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \dots & a_{n+q} \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q} & \dots & a_{n+2q-2} \end{vmatrix}$$

This determinant has been investigated by several authors in the literature [1, 18]. It is interesting to note that the Hankel determinants  $H_2(1) = |a_3 - a_2^2|$  and  $H_2(2) = |a_2 a_4 - a_3^2|$  are well known as Fekete-Szegő functional and second Hankel determinant respectively.

**Corresponding Author: V. Sivasankari\*<sup>2</sup>**

<sup>1,2</sup>Department of Science and Humanities,  
 PES University, Electronic City Campus, Bangalore - 560100, Karnataka, India.

The Fekete-Szegő problem for the well known classes

$$ST := \left\{ f \in A : \operatorname{Re} \left( \frac{zf'(z)}{f(z)} \right) > 0, z \in \mathbb{U} \right\}$$

and

$$CV := \left\{ f \in A : \operatorname{Re} \left( \frac{(zf'(z))'}{f'(z)} \right) > 0, z \in \mathbb{U} \right\}$$

was investigated by Keogh and Merkes [12]. Recently, many authors have discussed upper bounds for the Hankel determinant of functions belonging to various subclasses of univalent functions [3, 4, 22] and references therein.

Janteng et al. discussed the Hankel determinant problem for the classes of starlike functions with respect to symmetric points and convex functions with respect to symmetric points in [9] and for the functions whose derivative has a positive real part in [10]. In their work, they have shown that if  $f \in RT$  then  $|a_2a_4 - a_3^2| \leq \frac{4}{9}$ . In [11], the authors also obtained the second Hankel determinant and sharp bounds for the familiar subclasses of  $S$ , namely starlike and convex functions denoted by  $ST$  and  $CV$  and showed that  $|a_2a_4 - a_3^2| \leq 1$  and  $|a_2a_4 - a_3^2| \leq \frac{1}{8}$  respectively. Mishra and Gochhayat [16] have obtained the sharp bound to the non-linear functional  $|a_2a_4 - a_3^2|$  for the class of analytic functions denoted by  $R_\lambda(\alpha, \rho)$ ,  $(0 \leq \rho \leq p, |\alpha| < \frac{\pi}{2p})$ .

Analogous to the Hankel determinant of univalent functions, we consider the Hankel determinant in the case  $q = 2$  and  $n = p$ , known as second Hankel determinant for multivalent functions given by

$$H_2(p) = \begin{vmatrix} a_{p+1} & a_{p+2} \\ a_{p+2} & a_{p+3} \end{vmatrix}.$$

Estimate on the functional  $|a_{p+1}a_{p+3} - a_{p+2}^2|$  for the classes of  $p$ -valent starlike and  $p$ -valent convex functions were obtained by Krishna and Ramreddy [21]. However, for any real number  $\mu$ , the sharp estimate on the functional  $|a_{p+2} - \mu a_{p+1}^2|$  for the classes of  $p$ -valent starlike and convex functions of order  $\alpha$  were obtained by Hayami and Owa [7].

Inspired by the earlier works obtained by different researchers in this direction, we in the present paper, obtain an upper bound to the functional  $|a_{p+1}a_{p+3} - a_{p+2}^2|$  for the functions  $f$  belonging to multivalent  $\alpha$ -spiral starlike and  $\alpha$ -spiral convex functions which are defined as follows.

**Definition 1.1:** For  $-1 \leq B < A \leq 1$ , a function  $f \in A_p$ , given by (1.1), is said to be  $p$ -valently  $\alpha$ -spiral starlike if it satisfies the inequality

$$\frac{e^{-i\alpha} zf'(z)}{pf(z)} < \cos \alpha \left( \frac{1+Az}{1+Bz} \right) - i \sin \alpha, \text{ for all } z \in \mathbb{U}, |\alpha| \leq \frac{\pi}{2p}. \quad (1.2)$$

We denote this class of functions by  $SP_{p,\alpha}(A, B)$ .

By specializing on the values of  $A, B, \alpha$  and  $p$ , we obtain subclasses of analytic functions that were studied earlier in literature.

1.  $SP_{p,\alpha}(1, -1) = SP_p(\alpha)$ , the class of  $p$ -valently  $\alpha$ -spiral functions.
2.  $SP_{p,0}(1 - 2\alpha, -1) = ST_p(\alpha)$ , the class of  $p$ -valent starlike functions of order  $\alpha$  was studied by Hayami and Owa[7] and Vamshee Krishna *et al.* [21]
3.  $SP_{1,0}(A, B) = ST^*(A, B)$ , the subclass of starlike functions was studied by Goel and Mehrotra [5] and G.Singh *et al.* [4].
4.  $SP_{1,\alpha}(1, -1) = SP(\alpha)$ , the class of  $\alpha$ -spiral functions introduced by Spacsek [20].
5.  $SP_{1,0}(1, -1) = ST$ , the class of starlike functions, studied by Janteng *et al.* [11].

**Definition 1.2:** For  $-1 \leq B < A \leq 1$ , a function  $f \in A_p$ , given by (1.1), is said to be  $p$ -valently convex  $\alpha$ -spiral if it satisfies the inequality

$$\frac{1}{p} \left[ e^{-i\alpha} \left( 1 + \frac{zf''(z)}{f'(z)} \right) \right] < \cos \alpha \left( \frac{1+Az}{1+Bz} \right) - i \sin \alpha, \text{ for all } z \in \mathbb{U}, |\alpha| \leq \frac{\pi}{2p}. \quad (1.3)$$

We denote this class of functions by  $CVSP_{p,\alpha}(A, B)$ .

By specializing on the values of  $A, B, \alpha$  and  $p$ , we obtain subclasses of analytic functions that were studied earlier in literature.

1.  $CVSP_{p,\alpha}(1, -1) = CVSP_p(\alpha)$ , the class of  $p$ -valently convex  $\alpha$  - spiral functions.
2.  $CVSP_{p,0}(1 - 2\alpha, -1) = CV_p(\alpha)$ , the class of  $p$ -valent convex functions of order  $\alpha$  was studied by Hayami and Owa [7] and Vamshee Krishna *et al.* [21].
3.  $CVSP_{1,0}(A, B) = K(A, B)$ , the subclass of convex functions was studied by Goel and Mehrook [5] and G.Singh *et al.* [4].
4.  $CVSP_{1,\alpha}(1, -1) = CVSP(\alpha)$ , the class of convex  $\alpha$  - spiral functions studied by Vamshee Krishna *et al.* [22].
5.  $CVSP_{1,0}(1, -1) = CV$ , the class of convex functions, studied by Janteng *et al.* [11].

In order to prove our main results, we shall need the following preliminary lemma.

Let  $P$  denote the class of functions  $p(z)$  of the form

$$p(z) = 1 + c_1z + c_2z^2 + c_3z^3 + \dots, \tag{1.4}$$

which are analytic in the open unit disk  $\mathbb{U}$  for which  $Re\{p(z)\} > 0$ .

**Lemma 1.3:** [19] *If the function  $p \in P$  is given by the series (1.4), then the following sharp estimate holds :*

$$|c_k| \leq 2, \quad k = 1, 2, \dots \tag{1.5}$$

**Lemma 1.4:** [13, 14] *If the function  $p \in P$  is given by the series (1.4), then*

$$\begin{aligned} 2c_2 &= c_1^2 + x(4 - c_1^2), \\ 4c_3 &= c_1^3 + 2c_1(4 - c_1^2)x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)y, \end{aligned}$$

for some  $x, y$  with  $|x| \leq 1, |y| \leq 1$  and  $c_1 \in [0, 2]$ .

## 2. MAIN RESULTS ON $\alpha$ - SPIRAL STARLIKE FUNCTION

**Theorem 2.1:** *Let the function  $f$  given by (1.1) be in the class  $SP_{p,\alpha}(A, B)$ . Then*

$$|a_{p+1}a_{p+3} - a_{p+2}^2| \leq \frac{(A-B)^2 p^2 \cos^2(\alpha)}{4} \left( -\frac{\pi}{2p} \leq \alpha \leq \frac{\pi}{2p} \right). \tag{2.1}$$

**Proof:** If  $f(z) \in SP_{p,\alpha}(A, B)$ , then there exist a Schwarz function  $w(z) \in \Omega$  such that

$$\frac{e^{-i\alpha} z f'(z)}{p f(z)} = \cos \alpha \phi(w(z)) - i \sin \alpha, \quad z \in U \tag{2.2}$$

where

$$\begin{aligned} \phi(z) &= \frac{1 + Az}{1 + Bz} = 1 + (A - B)z - B(A - B)z^2 + B^2(A - B)z^3 + \dots \\ &= 1 + B_1z + B_2z^2 + B_3z^3 + \dots \end{aligned} \tag{2.3}$$

Define the function  $p_1(z)$  by

$$p_1(z) = \frac{1+w(z)}{1-w(z)} = 1 + c_1z + c_2z^2 + c_3z^3 + \dots, \quad z \in \mathbb{U} \tag{2.4}$$

Since  $w(z)$  is a schwarz function, we see that  $Re(p_1(z)) > 0$  and  $p_1(0) = 1$ .

Define the function  $h(z)$  by

$$h(z) = \frac{e^{-i\alpha} z f'(z) + i \sin \alpha f(z)}{p f(z) \cos \alpha}, \quad z \in \mathbb{U} \tag{2.5}$$

In view of the equations (2.2), (2.4) and (2.5), we have

$$\begin{aligned} h(z) &= \phi \left( \frac{p_1(z)-1}{p_1(z)+1} \right) = \phi \left( \frac{c_1z + c_2z^2 + c_3z^3 + \dots}{2 + c_1z + c_2z^2 + c_3z^3 + \dots} \right) \\ &= \phi \left( \frac{1}{2} c_1z + \frac{1}{2} \left( c_2 - \frac{c_1^2}{2} \right) z^2 + \frac{1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) z^3 + \dots \right) \\ &= 1 + \frac{B_1c_1}{2} z + \left[ \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2c_1^2}{4} \right] z^2 \\ &+ \left[ \frac{B_1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) + \frac{B_2c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3c_1^3}{8} \right] z^3 + \dots \end{aligned} \tag{2.6}$$

From (2.5), we have

$$e^{-i\alpha} z f'(z) + i p \sin \alpha f(z) = p \cos \alpha \{f(z) \times h(z)\}. \tag{2.7}$$

Replacing  $f(z), f'(z)$  by their equivalent  $p$ -valent expressions and also the equivalent expression for  $h(z)$  in (2.6), we have

$$\begin{aligned} & e^{-i\alpha} z(pz^{p-1} + \sum_{n=p+1}^{\infty} na_n z^{n-1}) + ip \sin\alpha(z^p + \sum_{n=p+1}^{\infty} a_n z^n) \\ &= p \cos \alpha \left[ (z^p + \sum_{n=p+1}^{\infty} a_n z^n) \times \left( 1 + \frac{B_1 c_1}{2} z + \left[ \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2 c_1^2}{4} \right] z^2 \right. \right. \\ & \left. \left. + \left[ \frac{B_1}{2} \left( c_3 - c_1 c_2 + \frac{c_1^3}{4} \right) + \frac{B_2 c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3 c_1^3}{8} \right] z^3 + \dots \right) \right]. \end{aligned}$$

Upon simplification, we obtain

$$\begin{aligned} & e^{-i\alpha} (a_{p+1} z^{p+1} + 2a_{p+2} z^{p+2} + 3a_{p+3} z^{p+3} + \dots) \\ &= p \cos \alpha (z^p + a_{p+1} z^{p+1} + a_{p+2} z^{p+2} + a_{p+3} z^{p+3} + \dots) \\ & \left( \frac{B_1 c_1}{2} z + \left[ \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2 c_1^2}{4} \right] z^2 + \dots \right) \end{aligned} \tag{2.8}$$

Equating the coefficients of like powers of  $z^{p+1}, z^{p+2}$  and  $z^{p+3}$  respectively in (2.8), we have

$$\begin{aligned} a_{p+1} e^{-i\alpha} &= \frac{B_1 c_1}{2} p \cos \alpha, \\ 2a_{p+2} e^{-i\alpha} &= a_{p+1} \frac{B_1 c_1}{2} p \cos \alpha + \left( \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2 c_1^2}{4} \right) p \cos \alpha, \\ 3a_{p+3} e^{-i\alpha} &= a_{p+2} \frac{B_1 c_1}{2} p \cos \alpha + a_{p+1} \left( \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2 c_1^2}{4} \right) p \cos \alpha \\ & \quad + \left[ \frac{B_1}{2} \left( c_3 - c_1 c_2 + \frac{c_1^3}{4} \right) + \frac{B_2 c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3 c_1^3}{8} \right] p \cos \alpha. \end{aligned}$$

After simplifying using (2.3), we obtain

$$\begin{aligned} a_{p+1} &= e^{i\alpha} \frac{(A-B)c_1}{2} p \cos \alpha, \\ a_{p+2} &= \frac{e^{i\alpha}(A-B)}{8} [e^{i\alpha}(A-B)c_1^2 p \cos \alpha + 2c_2 - c_1^2 - Bc_1^2] p \cos \alpha, \\ a_{p+3} &= \frac{e^{i\alpha}}{48} (A-B) [8c_3 + e^{2i\alpha}(A-B)^2 c_1^3 p^2 \cos^2(\alpha) - 3(A-B)e^{i\alpha} p \cos \alpha c_1^3 \\ & \quad - 3B(A-B)e^{i\alpha} c_1^3 + (2+4B+2B^2)c_1^3 + (6e^{i\alpha} p \cos \alpha(A-B) - 8-8B)c_1 c_2] p \cos \alpha. \end{aligned}$$

Substituting the values of  $a_{p+1}, a_{p+2}$  and  $a_{p+3}$  in the second Hankel functional, we have

$$\begin{aligned} |a_{p+1} a_{p+3} - a_{p+2}^2| &= \left| \frac{e^{2i\alpha}(A-B)^2 p^2 \cos^2 \alpha}{192} [16c_3 c_1 - e^{2i\alpha}(A-B)^2 c_1^4 p^2 \cos^2 \alpha \right. \\ & \quad \left. - 12c_2^2 + (1+B)^2 c_1^4 - 4(1+B)c_2 c_1^2] \right|. \end{aligned}$$

By using the facts  $|xp + yq| \leq x|p| + y|q|$ , where  $x, y, p$  and  $q$  are real numbers and  $|e^{ni\alpha}| = 1$ , upon simplification, we obtain

$$\begin{aligned} |a_{p+1} a_{p+3} - a_{p+2}^2| &\leq \frac{(A-B)^2 p^2 \cos^2 \alpha}{192} |[16c_3 c_1 - (A-B)^2 c_1^4 p^2 \cos^2 \alpha \\ & \quad - 12c_2^2 + (1+B)^2 c_1^4 - 4(1+B)c_2 c_1^2]|. \end{aligned} \tag{2.9}$$

Substituting the values of  $c_2$  and  $c_3$  from Lemma 1.4, we have

$$\begin{aligned} |a_{p+1} a_{p+3} - a_{p+2}^2| &\leq \frac{(A-B)^2 p^2 \cos^2 \alpha}{192} |4c_1 [c_1^3 + 2c_1(4-c_1^2)x - c_1(4-c_1^2)x^2 \\ & \quad + 2(4-c_1^2)(1-|x|^2)y] - (A-B)^2 c_1^4 p^2 \cos^2 \alpha + (1+B)^2 c_1^4 \\ & \quad - 3[c_1^4 + (4-c_1^2)^2 x + 2c_1^2(4-c_1^2)x] - 2(1+B)c_1^2 [c_1^2 + (4-c_1^2)x]|. \end{aligned}$$

Assume that  $c_1 = c$  and  $c \in [0,2]$ , using triangular inequality and  $|y| \leq 1$ , we have

$$\begin{aligned} |a_{p+1} a_{p+3} - a_{p+2}^2| &\leq \frac{(A-B)^2 p^2 \cos^2 \alpha}{192} (|10 + 4B + B^2|c^4 + (A-B)^2 p^2 \cos^2 \alpha c^4 \\ & \quad + |(16 + 2B)c^2(4 - c^2)|\delta + |(12 - 7c^2 - 8c)(4 - c^2)|\delta^2) \\ &= \frac{(A-B)^2 p^2 \cos^2 \alpha}{192} F(c, \delta), \text{ where } \delta = |x| \leq 1 \text{ and} \end{aligned}$$

$$F(c, \delta) = |10 + 4B + B^2|c^4 + (A-B)^2 p^2 \cos^2 \alpha c^4 + |(16 + 2B)c^2(4 - c^2)|\delta + |(12 - 7c^2 - 8c)(4 - c^2)|\delta^2. \tag{2.10}$$

Now the function  $F(c, \delta)$  is maximized on the closed square  $[0,2] \times [0,1]$ . Differentiating  $F(c, \delta)$  in (2.10), partially with respect to  $\delta$ , we get

$$\frac{\partial F}{\partial \delta} = [| (16 + 2B)c^2(4 - c^2) | + |(12 - 7c^2 - 8c)(4 - c^2)|](2\delta). \tag{2.11}$$

For  $0 < \delta < 1$ , and for fixed  $c$  with  $0 < c < 2$ , from (2.11) we observe that  $\frac{\partial F}{\partial \delta} > 0$ .

Consequently,  $F(c, \delta)$  is an increasing function of  $\delta$  and hence cannot have maximum value at any point in the interior of the closed square  $[0,2] \times [0,1]$ . Moreover, for fixed  $c \in [0,2]$ , we have

$$\max_{0 \leq \delta \leq 1} F(c, \delta) = F(c, 1) = G(c). \quad (2.12)$$

Upon simplifying the relation (2.10) and (2.12), we obtain

$$G(c) = (A - B)^2 p^2 \cos^2 \alpha c^4 + (1 + B)^2 c^4 + 8c^3 + |(24 + 8B)|c^2 - 32c + 48. \quad (2.13)$$

Differentiation yields :

$$G'(c) = (A - B)^2 p^2 \cos^2 \alpha (4c^3) + (1 + B)^2 (4c^3) + 24c^2 + 2|(24 + 8B)|c - 32. \quad (2.14)$$

From the expression (2.14), we observe that  $G'(c) \leq 0$  from all values of  $c$  in the interval  $0 \leq c \leq 2$  and for a fixed valued of  $\alpha$  with  $\left(\frac{-\pi}{2p} \leq \alpha \leq \frac{\pi}{2p}\right)$ .

Therefore,  $G(c)$  is a monotonically decreasing function of  $c$  in the interval  $[0,2]$ . So, that its maximum value occurs at  $c = 0$ . From (2.13), we get

$$\max_{0 \leq c \leq 2} G(0) = 48. \quad (2.15)$$

After simplifying the expressions (2.9) and (2.15) we obtain

$$|16c_3 c_1 - (A - B)^2 c_1^4 p^2 \cos^2 \alpha - 12c_2^2 + (1 + B)^2 c_1^4 - 4(1 + B)c_2 c_1^2| \leq 48. \quad (2.16)$$

Upon simplifying the expressions (2.9) and (2.16), we get

$$|a_{p+1} a_{p+3} - a_{p+2}^2| \leq \frac{(A-B)^2 p^2 \cos^2(\alpha)}{4}. \quad (2.17)$$

Choosing  $c_1 = c = 0$  and selecting  $x = -1$  in Lemma 1.3, we find that  $c_2 = -2$  and  $c_3 = 0$ . Substituting these values in (2.16), it is observed that equality is attained which shows that our result is sharp. This completes the proof.

Choosing  $A = 1, B = -1$ , Theorem 2.1 gives the following result.

**Corollary 2.2:** If  $f(z) \in SP_p(\alpha)$   $\left(\frac{-\pi}{2p} \leq \alpha \leq \frac{\pi}{2p}\right)$ , then  $|a_{p+1} a_{p+3} - a_{p+2}^2| \leq p^2 \cos^2(\alpha)$ .

Choosing  $A = 1 - 2\beta$  ( $0 \leq \beta < 1$ ),  $B = -1, \alpha = 0$  Theorem 2.1 gives the following result.

**Corollary 2.3:** If  $f(z) \in ST_p(\beta)$   $\left(0 \leq \beta \leq \left(p - \frac{1}{2}\right)\right)$ , then  $|a_{p+1} a_{p+3} - a_{p+2}^2| \leq (p - \beta)^2$ .

Choosing  $p = 1, \alpha = 0$  Theorem 2.1 gives the following result.

**Corollary 2.4:** If  $f(z) \in ST^*(A, B)$  ( $-1 \leq B < A \leq 1$ ), then  $|a_2 a_4 - a_3^2| \leq \frac{(A-B)^2}{4}$ .

Choosing  $A = 1, B = -1$ , and  $p = 1$  Theorem 2.1 gives the following result.

**Corollary 2.5** If  $f(z) \in SP(\alpha)$  and for  $\left(\frac{-\pi}{2} \leq \alpha \leq \frac{\pi}{2}\right)$  then  $|a_2 a_4 - a_3^2| \leq \cos^2(\alpha)$ .

Choosing  $A = 1, B = -1, p = 1$  and  $\alpha = 0$  Theorem 2.1 gives the following result.

**Corollary 2.6:** If  $f(z) \in ST$ , then  $|a_2 a_4 - a_3^2| \leq 1$ .

This inequality is sharp and coincides with that of Janteng, Halim and Darus [11].

### 3. MAIN RESULTS ON $\alpha$ - SPIRAL CONVEX FUNCTION

**Theorem 3.1:** Let the function  $f$  given by (1.1) be in the class  $CVSP_{p,\alpha}(A, B)$ . Then

$$\frac{|a_{p+1} a_{p+3} - a_{p+2}^2| \leq p^4 (A-B)^2 \cos^2(\alpha) [6 - (1+B)(p^2 + 4p + 7) + 3(A-B)p \cos \alpha]^2 + 48(p+1)(p+3)\Delta(A,B)}{12(p+1)(p+2)^2(p+3)\Delta(A,B)} \quad (3.1)$$

where

$$\Delta(A, B) = 2(p^2 + 4p + 7) + (A - B)^2(p^2 + 4p + 1)p^2 \cos^2(\alpha) + 6(A - B)p \cos(\alpha)(1 + B) - 4(1 + B)(p^2 + 4p + 1) - (p^2 + 4p + 7)(B^2 + 4B + 3).$$

**Proof:** If  $f(z) \in CVSP_{p,\alpha}(A, B)$ , then there exist a Schwarz function  $w(z) \in \Omega$  such that

$$\frac{1}{p} \left[ e^{-i\alpha} \left( 1 + \frac{zf''(z)}{f'(z)} \right) \right] = \cos \alpha \phi(w(z)) - i \sin \alpha, \quad z \in \mathbb{U} \tag{3.2}$$

where

$$\begin{aligned} \phi(z) &= \frac{1 + Az}{1 + Bz} = 1 + (A - B)z - B(A - B)z^2 + B^2(A - B)z^3 + \dots \\ &= 1 + B_1z + B_2z^2 + B_3z^3 + B_4z^4 + \dots. \end{aligned} \tag{3.3}$$

Define the function  $p_1(z)$  by

$$p_1(z) = \frac{1+w(z)}{1-w(z)} = 1 + c_1z + c_2z^2 + c_3z^3 + \dots, \quad z \in \mathbb{U} \tag{3.4}$$

Since  $w(z)$  is a schwarz function, we see that  $Re(p_1(z)) > 0$  and  $p_1(0) = 1$ . Define the function  $h(z)$  by

$$h(z) = \frac{e^{-i\alpha}[f'(z) + zf''(z)] + i \sin \alpha f'(z)}{p \cos \alpha f'(z)}, \quad z \in \mathbb{U} \tag{3.5}$$

In view of the equations (3.2), (3.4) and (3.5), we have

$$\begin{aligned} h(z) &= \phi \left( \frac{p_1(z)-1}{p_1(z)+1} \right) = \phi \left( \frac{c_1z + c_2z^2 + c_3z^3 + \dots}{2 + c_1z + c_2z^2 + c_3z^3 + \dots} \right) \\ &= \phi \left( \frac{1}{2}c_1z + \frac{1}{2} \left( c_2 - \frac{c_1^2}{2} \right) z^2 + \frac{1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) z^3 + \dots \right) \\ &= 1 + \frac{B_1c_1}{2}z + \left[ \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2c_1^2}{4} \right] z^2 \\ &\quad + \left[ \frac{B_1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) + \frac{B_2c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3c_1^3}{8} \right] z^3 + \dots. \end{aligned} \tag{3.6}$$

From (3.5), we have

$$e^{-i\alpha} [f'(z) + zf''(z)] + i \sin \alpha f'(z) = p \cos \alpha \{f'(z) \times h(z)\}.$$

Replacing  $f'(z)$  and  $f''(z)$  by their equivalent  $p$ -valent expressions and also the equivalent expression for  $h(z)$  in (3.6), we have

$$\begin{aligned} &e^{-i\alpha} (pz^{p-1} + \sum_{n=p+1}^{\infty} na_n z^{n-1}) + z(p(p-1)z^{p-2} + \sum_{n=p+1}^{\infty} n(n-1)a_n z^{n-2}) \\ &\quad + i \sin \alpha (pz^{p-1} + \sum_{n=p+1}^{\infty} na_n z^{n-1}) \\ &= p \cos \alpha \left[ (pz^{p-1} + \sum_{n=p+1}^{\infty} na_n z^{n-1}) \times \left\{ 1 + \frac{B_1c_1}{2}z + \left[ \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2c_1^2}{4} \right] z^2 \right. \right. \\ &\quad \left. \left. + \left[ \frac{B_1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) + \frac{B_2c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3c_1^3}{8} \right] z^3 + \dots \right\} \right]. \end{aligned} \tag{3.7}$$

Equating the coefficients of like powers of  $z^p, z^{p+1}$  and  $z^{p+2}$  respectively in (3.7), we have

$$\begin{aligned} (p+1)a_{p+1}e^{-i\alpha} &= p \frac{B_1c_1}{2} p \cos \alpha, \\ 2(p+2)a_{p+2}e^{-i\alpha} &= (p+1)a_{p+1} \frac{B_1c_1}{2} p \cos \alpha + p \left( \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2c_1^2}{4} \right) p \cos \alpha, \\ 3(p+3)a_{p+3}e^{-i\alpha} &= (p+2)a_{p+2} \frac{B_1c_1}{2} p \cos \alpha + (p+1)a_{p+1} \left( \frac{B_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_2c_1^2}{4} \right) p \cos \alpha \\ &\quad + p \left[ \frac{B_1}{2} \left( c_3 - c_1c_2 + \frac{c_1^3}{4} \right) + \frac{B_2c_1}{2} \left( c_2 - \frac{c_1^2}{2} \right) + \frac{B_3c_1^3}{8} \right] p \cos \alpha. \end{aligned}$$

After simplifying using (3.3), we obtain

$$\begin{aligned} a_{p+1} &= e^{i\alpha} \frac{(A-B)c_1}{2(p+1)} p^2 \cos \alpha, \\ a_{p+2} &= \frac{e^{i\alpha}(A-B)}{8(p+2)} [e^{i\alpha}(A-B)c_1^2 p \cos \alpha + 2c_2 - c_1^2(1+B)] p^2 \cos \alpha, \\ a_{p+3} &= \frac{e^{i\alpha}}{48(p+3)} (A-B) [8c_3 + e^{2i\alpha}(A-B)^2 c_1^3 p^2 \cos^2(\alpha) \\ &\quad - 3(A-B)e^{i\alpha} p \cos \alpha c_1^3(1+B) + 2(1+B)^2 c_1^3 + 6e^{i\alpha} p \cos \alpha (A-B)c_1c_2 - 8c_1c_2(1+B)] p^2 \cos \alpha. \end{aligned}$$

Substituting the values of  $a_{p+1}, a_{p+2}$  and  $a_{p+3}$  in the second Hankel functional and applying the same procedure as described in Theorem 2.1, upon simplification, we have

$$\begin{aligned}
 |a_{p+1}a_{p+3} - a_{p+2}^2| \leq & \frac{p^4 \cos^2(\alpha)(A-B)^2}{12(p+1)(p+2)^2(p+3)} \left| [(p+2)^2 c_1 c_3 - c_1^2 c_2 (1+B)(p+2)^2 \right. \\
 & + \frac{c_1^4}{4} (p+2)^2 (1+B)^2 + \frac{3}{4} (A-B) c_1^2 c_2 (p+2)^2 p \cos \alpha \\
 & - \frac{3}{8} (A-B) c_1^4 p \cos \alpha (1+B)(p+2)^2 \\
 & + \frac{(A-B)^2}{8} c_1^4 p^2 \cos^2(\alpha) (p+2)^2 - 3(p+1)(p+3) \frac{c_2^2}{4} \\
 & - \frac{3}{16} (p+1)(p+3) c_1^4 (1+B)^2 - \frac{3}{16} (A-B)^2 (p+1)(p+3) c_1^4 p^2 \cos^2(\alpha) \\
 & + \frac{3}{4} c_1^2 c_2 (1+B)(p+1)(p+3) + \frac{3}{8} (A-B) c_1^4 (1+B) p \cos \alpha (p+1)(p+3) \\
 & \left. - \frac{3}{4} (p+1)(p+3)(A-B) c_1^2 c_2 p \cos \alpha \right|. \tag{3.8}
 \end{aligned}$$

The above expression is equivalent to

$$|a_{p+1}a_{p+3} - a_{p+2}^2| \leq \frac{p^4 \cos^2(\alpha)(A-B)^2}{12(p+1)(p+2)^2(p+3)} \times |d_1 c_1 c_3 + d_2 c_1^2 c_2 + d_3 c_2^2 + d_4 c_1^4| \tag{3.9}$$

where

$$\begin{aligned}
 d_1 &= (p+2)^2 \\
 d_2 &= -(1+B)(p+2)^2 + \frac{3}{4}(1+B)(p^2+4p+3) + \frac{3}{4}(A-B)p \cos \alpha \\
 d_3 &= -\frac{3}{4}(p^2+4p+3) \\
 d_4 &= \frac{1}{16}(1+B)^2(p^2+4p+7) - \frac{3}{8}(A-B)(1+B)p \cos \alpha - \frac{1}{16}(A-B)^2 p^2 \cos^2 \alpha (p^2+4p+1)
 \end{aligned}$$

Substituting the values of  $c_2$  and  $c_3$  from Lemma (1.4) in the right hand side of (3.9), we have

$$\begin{aligned}
 & |d_1 c_1 c_3 + d_2 c_1^3 c_2 + d_3 c_2^2 + d_4 c_1^4| = \\
 & |d_1 c_1 \times \frac{1}{4} \{c_1^3 + 2c_1(4-c_1^2)x - c_1(4-c_1^2)x^2 + 2(4-c_1^2)(1-|x|^2)y\} + d_2 c_1^2 \\
 & \times \frac{1}{2} \{c_1^2 + x(4-c_1^2)\} + d_3 \times \frac{1}{4} \{c_1^2 + x(4-c_1^2)\}^2 + d_4 c_1^4|. \tag{3.10}
 \end{aligned}$$

After simplifying, we get

$$\begin{aligned}
 & 4|d_1 c_1 c_3 + d_2 c_1^2 c_2 + d_3 c_2^2 + d_4 c_1^4| = \\
 & |(d_1 + 2d_2 + d_3 + 4d_4)c_1^4 + 2d_1 c_1(4-c_1^2)y \\
 & + 2(d_1 + d_2 + d_3)c_1^2(4-c_1^2)|x| - \{(d_1 + d_3)c_1^2 + 2d_1 c_1 - 4d_3\}(4-c_1^2)|x|^2 y|. \tag{3.11}
 \end{aligned}$$

Substituting the values of  $d_1, d_2, d_3$  and  $d_4$ , we obtain

$$\begin{aligned}
 d_1 + 2d_2 + d_3 + 4d_4 &= \frac{1}{4}(p^2+4p+7) + (1+B)(p^2+4p+1) + \frac{1}{4}(1+B)^2(p^2+4p+7) \\
 - \frac{3}{2}B(A-B)p \cos \alpha - \frac{1}{4}(A-B)^2(p^2+4p+1)p^2 \cos^2 \alpha, \tag{3.12}
 \end{aligned}$$

$$d_1 + d_2 + d_3 = \frac{1}{4}(p^2+4p+7) - \frac{1}{4}(1+B)(p^2+4p+7) + \frac{3}{4}(A-B)p \cos \alpha, \tag{3.13}$$

$$(d_1 + d_3)c_1^2 + 2d_1 c_1 - 4d_3 = \frac{1}{4}(p^2+4p+7)c_1^2 + 2(p^2+4p+4)c_1 + 3(p^2+4p+3). \tag{3.14}$$

Consider

$$\begin{aligned}
 & (p^2+4p+7)c_1^2 + 8(p^2+4p+4)c_1 + 12(p^2+4p+3) \\
 & = (p^2+4p+7) \times \left[ c_1^2 + \frac{8(p^2+4p+4)}{p^2+4p+7} c_1 + \frac{12(p^2+4p+3)}{p^2+4p+7} \right] \\
 & = (p^2+4p+7) \times \left[ \left\{ c_1 + \frac{4(p^2+4p+4)}{(p^2+4p+7)} \right\}^2 - \frac{16(p^2+4p+4)}{(p^2+4p+7)^2} + \frac{12(p^2+4p+3)}{(p^2+4p+7)} \right].
 \end{aligned}$$

Upon simplification, the above expression can also be expressed as

$$\begin{aligned} & (p^2 + 4p + 7)c_1^2 + 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3) \\ &= (p^2 + 4p + 7) \times \left[ \left\{ c_1 + \frac{4(p^2 + 4p + 4)}{(p^2 + 4p + 7)} \right\}^2 - \left\{ \frac{2\sqrt{p^4 + 8p^3 + 18p^2 + 8p + 1}}{(p^2 + 4p + 7)} \right\}^2 \right] \\ &= (p^2 + 4p + 7) \times \left[ c_1 + \left\{ \frac{4(p^2 + 4p + 4)}{(p^2 + 4p + 7)} + \frac{2\sqrt{p^4 + 8p^3 + 18p^2 + 8p + 1}}{(p^2 + 4p + 7)} \right\} \right] \\ & \quad \times \left[ c_1 + \left\{ \frac{4(p^2 + 4p + 4)}{(p^2 + 4p + 7)} - \frac{2\sqrt{p^4 + 8p^3 + 18p^2 + 8p + 1}}{(p^2 + 4p + 7)} \right\} \right]. \end{aligned} \quad (3.15)$$

Since  $c_1 \in [0,2]$ , using the result  $(c_1 + a)(c_1 + b) \geq (c_1 - a)(c_1 - b)$ , where  $a, b > 0$  in the right-hand side of (3.15), upon simplification, we obtain

$$\begin{aligned} & (p^2 + 4p + 7)c_1^2 + 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3) \geq \\ & (p^2 + 4p + 7)c_1^2 - 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3). \end{aligned} \quad (3.16)$$

From the relations (3.14) and (3.16), we obtain

$$-4(d_1 + d_3)c_1^2 + 2d_1c_1 - 4d_3 \leq -\{(p^2 + 4p + 7)c_1^2 - 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3)\}. \quad (3.17)$$

Substituting the calculated values from (3.13) and (3.17) in the right-hand side of the relation (3.12), we get

$$\begin{aligned} & 16|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \\ & \leq \{[(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + (1 + B)^2(p^2 + 4p + 7) \\ & \quad - 6B(A - B)p\cos \alpha - (A - B)^2(p^2 + 4p + 1)p^2\cos^2 \alpha]c_1^4 + 8(p^2 + 4p + 4)c_1(4 - c_1^2)y \\ & \quad + 2[(p^2 + 4p + 7) - (1 + B)(p^2 + 4p + 7) + 3(A - B)p\cos \alpha]c_1^2(4 - c_1^2)|x| \\ & \quad + [(p^2 + 4p + 7)c_1^2 - 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3)](4 - c_1^2)|x|^2y\}. \end{aligned} \quad (3.18)$$

Choosing  $c_1 = c \in [0,2]$ , applying triangle inequality and using  $|y| \leq 1$ , and also replacing  $|x|$  by  $\delta$  in the right hand side of (3.18), it reduces to

$$\begin{aligned} & 16|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \\ & \leq \{[(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + (1 + B)^2(p^2 + 4p + 7) \\ & \quad - 6B(A - B)p\cos \alpha - (A - B)^2(p^2 + 4p + 1)p^2\cos^2 \alpha]c^4 + 8(p^2 + 4p + 4)c_1(4 - c^2) \\ & \quad + 2[(p^2 + 4p + 7) - (1 + B)(p^2 + 4p + 7) + 3(A - B)p\cos \alpha]c^2(4 - c^2)\delta \\ & \quad + [(p^2 + 4p + 7)c^2 - 8(p^2 + 4p + 4)c_1 + 12(p^2 + 4p + 3)](4 - c^2)\delta^2\} \\ & = F(c, \delta), \quad \text{for } 0 \leq \delta = |x| \leq 1. \end{aligned} \quad (3.19)$$

We assume that the upper bound for (3.19) occurs at an interior point of the set  $\{\delta, c\}$ :  $\delta \in [0,1]$  and  $c \in [0,2]$ .

Differentiating  $F(c, \delta)$  in (3.19) partially with respect to  $\delta$ , we get

$$\begin{aligned} \frac{\partial F}{\partial \delta} &= \{2(p^2 + 4p + 7) - (1 + B)(p^2 + 4p + 7) + 3(A - B)p\cos \alpha\}c^2(4 - c^2) \\ & \quad + \{2(p^2 + 4p + 7)c^2 - 8(p^2 + 4p + 4)c + 12(p^2 + 4p + 3)\}(4 - c^2)\delta. \end{aligned} \quad (3.20)$$

For  $0 \leq \delta \leq 1$ , and for fixed  $c$  with  $0 \leq c \leq 2$  and  $\left(\frac{-\pi}{2p} \leq \alpha \leq \frac{\pi}{2p}\right)$ , from (3.20) we observe that  $\frac{\partial F}{\partial \delta} > 0$ .

Consequently,  $F(c, \delta)$  is an increasing function of  $\delta$  and hence cannot have maximum value at any point in the interior of the closed square  $[0,2] \times [0,1]$ . Moreover, for fixed  $c \in [0,2]$ , we have

$$\max_{0 \leq \delta \leq 1} F(c, \delta) = F(c, 1) = G(c). \quad (3.21)$$

Upon simplifying the relation (3.19) and (3.21), we obtain

$$\begin{aligned} G(c) &= \{[-2(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + (p^2 + 4p + 7)(B^2 + 4B + 3) \\ & \quad - 6(A - B)p\cos \alpha(1 + B) - (A - B)^2(p^2 + 4p + 1)p^2\cos^2 \alpha]c^4 \\ & \quad + [48 - 8(1 + B)(p^2 + 4p + 7) + 24(A - B)p\cos \alpha]c^2 + 48(p + 1)(p + 3)\} \end{aligned} \quad (3.22)$$



Differentiation yields:

$$G'(c) = \{[-2(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + (p^2 + 4p + 7)(B^2 + 4B + 3) - 6(A - B)pcos \alpha(1 + B) - (A - B)^2(p^2 + 4p + 1)p^2cos^2\alpha]\}(4c^3) + [48 - 8(1 + B)(p^2 + 4p + 7) + 24(A - B)pcos \alpha](2c). \quad (3.23)$$

$$G''(c) = \{[-2(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + 12(p^2 + 4p + 7)(B^2 + 4B + 3) - (A - B)pcos \alpha(1 + B) - (A - B)^2(p^2 + 4p + 1)p^2cos^2\alpha]\}(12c^2) + [48 - 8(1 + B)(p^2 + 4p + 7) + 24(A - B)pcos \alpha](2). \quad (3.24)$$

The maximum of minimum value of  $G(c)$  is obtained for the values of  $G'(c) = 0$ . From the expression (3.23), we get

$$G'(c) = \{[-2(p^2 + 4p + 7) + 4(1 + B)(p^2 + 4p + 1) + (p^2 + 4p + 7)(B^2 + 4B + 3) - 6(A - B)pcos \alpha(1 + B) - (A - B)^2(p^2 + 4p + 1)p^2cos^2\alpha]\}(4c^3) + [48 - 8(1 + B)(p^2 + 4p + 7) + 24(A - B)pcos \alpha](2c) = 0. \quad (3.25)$$

We now discuss the following cases.

**Case-1:** If  $c = 0$ , then from (3.24), we obtain

$$G''(c) = 96 - 16(1 + B)(p^2 + 4p + 7) + 48(A - B)pcos \alpha > 0 \quad \text{because } |\alpha| \leq \frac{\pi}{2p}.$$

Therefore, by the second derivative test,  $G(c)$  has a minimum value at  $c = 0$ , which is ruled out.

**Case-2:** If  $c \neq 0$ , then from (3.25), we obtain

$$c^2 = \frac{4(6 - (1 + B)(p^2 + 4p + 7) + 3(A - B)pcos \alpha)}{\Delta(A, B)} \quad (3.26)$$

where

$$\Delta(A, B) = 2(p^2 + 4p + 7) + (A - B)^2(p^2 + 4p + 1)p^2cos^2(\alpha) + 6(A - B)pcos(\alpha)(1 + B) - 4(1 + B)(p^2 + 4p + 1) - (p^2 + 4p + 7)(B^2 + 4B + 3)$$

Using the value of  $c^2$  in (3.24), after simplifying, we get

$$G''(c) = -(192 - 32(1 + B)(p^2 + 4p + 7) + 96(A - B)pcos \alpha) < 0 \quad \text{because } |\alpha| \leq \frac{\pi}{2p}.$$

From the second derivative test,  $G(c)$  has a maximum value at  $c$ , where  $c^2$  is given by (3.26).

From the expression (3.22), we have  $G$ -maximum value at  $c^2$ , after simplifying it is given by

$$\max_{0 \leq c \leq 2} G(c) = \frac{16[6 - (1 + B)(p^2 + 4p + 7) + 3(A - B)pcos \alpha]^2 + 48(p + 1)(p + 3)\Delta(A, B)}{12(p + 1)(p + 2)^2(p + 3)\Delta(A, B)} \quad (3.27)$$

where

$$\Delta(A, B) = 2(p^2 + 4p + 7) + (A - B)^2(p^2 + 4p + 1)p^2cos^2(\alpha) + 6(A - B)pcos(\alpha)(1 + B) - 4(1 + B)(p^2 + 4p + 1) - (p^2 + 4p + 7)(B^2 + 4B + 3)$$

Considering only the maximum value of  $G(c)$  at  $c$ , where  $c^2$  is given by (3.26). From the expression (3.19) and (3.27), upon simplification, we obtain

$$|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4| \leq \frac{[6 - (1 + B)(p^2 + 4p + 7) + 3(A - B)pcos \alpha]^2 + 48(p + 1)(p + 3)\Delta(A, B)}{12(p + 1)(p + 2)^2(p + 3)\Delta(A, B)} \quad (3.28)$$

From the expression (3.9) and (3.28), after simplifying, we get

$$|a_{p+1}a_{p+3} - a_{p+2}^2| \leq \frac{p^4(A - B)^2cos^2(\alpha)[6 - (1 + B)(p^2 + 4p + 7) + 3(A - B)pcos \alpha]^2 + 48(p + 1)(p + 3)\Delta(A, B)}{12(p + 1)(p + 2)^2(p + 3)\Delta(A, B)} \quad (3.29)$$

where

$$\Delta(A, B) = 2(p^2 + 4p + 7) + (A - B)^2(p^2 + 4p + 1)p^2cos^2(\alpha) + 6(A - B)pcos(\alpha)(1 + B) - 4(1 + B)(p^2 + 4p + 1) - (p^2 + 4p + 7)(B^2 + 4B + 3)$$

Choosing  $x = -1$  in Lemma 1.3, we find that  $c_2 = c_1^2 - 2$  and  $c_3 = c_1^3 - 3c_1$ . The result is sharp for  $c_1 = c$ ,  $c_2 = c^2 - 2$  and  $c_3 = c^3 - 3c$  where  $c^2$  is given by (3.26).

This completes the proof.

Choosing  $A = 1, B = -1$ , Theorem 3.1 gives the following result.

**Corollary 3.2:** If  $f(z) \in CVSP_p(\alpha)$  ( $-\frac{\pi}{2p} \leq \alpha \leq \frac{\pi}{2p}$ ), then

$$\frac{|a_{p+1}a_{p+3} - a_{p+2}^2| \leq p^4(6(1 + 2p\cos(\alpha)) + p^2\cos^2(\alpha) + (p + 1)(p + 3)(p^2 + 4p + 7 + 2(p^2 + 4p + 1)p^2\cos^2(\alpha))}{(p + 1)(p + 2)^2(p + 3)\{2(p^2 + 4p + 1) + (p^2 + 4p + 7)p^2\sec^2(\alpha)\}}.$$

Choosing  $A = 1 - 2\beta$  ( $0 \leq \beta < 1$ ),  $B = -1$ ,  $\alpha = 0$  Theorem 3.1 gives the following result.

**Corollary 3.3:** If  $f(z) \in CV_p(\beta)$  ( $0 \leq \beta \leq (p - \frac{1}{2})$ ), then

$$\frac{|a_{p+1}a_{p+3} - a_{p+2}^2| \leq p^2(p - \beta)^2[6(p + 1 - \beta)^2 + (p + 1)(p + 3)\{2\beta(\beta - 2p)(p^2 + 4p + 1) + (2p^4 + 8p^3 + 3p^2 + 4p + 7)\}]}{(p + 1)(p + 2)^2(p + 3)[2\beta(\beta - 2p)(p^2 + 4p + 1) + (2p^4 + 8p^3 + 3p^2 + 4p + 7)]}.$$

Choosing  $p = 1$ ,  $\alpha = 0$  Theorem 3.1 gives the following result.

**Corollary 3.4:** If  $f(z) \in K(A, B)$  ( $-1 \leq B < A \leq 1$ ), then

$$|a_2a_4 - a_3^2| \leq \frac{(A - B)^2}{576} \left[ \frac{16| - A^2 + 2B^2 + AB| - |A - 5B|^2 - 12|A - 5B| - 36}{| - A^2 + 2B^2 + AB| - |A - 5B| - 2} \right].$$

Choosing  $A = 1$ ,  $B = -1$ , and  $p = 1$  Theorem 3.1 gives the following result.

**Corollary 3.5:** If  $f(z) \in CVSP(\alpha)$  and for ( $-\frac{\pi}{2} \leq \alpha \leq \frac{\pi}{2}$ ) then  $|a_2a_4 - a_3^2| \leq \frac{17(1+\cos^2\alpha)+2\cos \alpha}{144(1+\sec^2\alpha)}$ .

Choosing  $A = 1$ ,  $B = -1$ ,  $p = 1$  and  $\alpha = 0$  Theorem 3.1 gives the following result.

**Corollary 3.6:** If  $f(z) \in CV$ , then  $|a_2a_4 - a_3^2| \leq \frac{1}{8}$ .

This inequality is sharp and coincides with that of Janteng, Halim and Darus [11].

## REFERENCES

1. P. Dienes, *The Taylor series: An introduction to the theory of functions of a complex variable*, Dover Publications, Inc., New York, 1957.
2. P.L.Duren, *Univalent Functions, Grundlehren der Mathematischen Wissenschaften, Springer, Newyork*, (259)(1983).
3. M. Elumalai, C. Selvaraj, *Second Hankel Determinant for Multivalent spirallike and convex functions of order  $\alpha$* , Inter.J.Sci.Res,5(4), 2029-2034 (2016).
4. Gagandeep Singh, Gurcharanjit Singh, *Second Hankel determinant for Subclasses of Starlike and Convex Functions*, Open Science Journal of Mathematics and Application, 2(6), 48-51 (2014).
5. R.M. Goel and B.S. Mehrok, *On the coefficeints of a subclass of starlike functions*, Ind.J. Pure and Appl.MATH., 12(5), 634-647 (1981).
6. U. Grenander and G. Szegö, *Toeplitz forms and their applications*, California Monographs in Mathematical Sciences, University of California Press, Berkeley, 1958.
7. T. Hayami and S. Owa, *Hankel determinant for  $p$ -valently starlike and convex functions of order  $\alpha$* , Gen. Math. 17, no. 4, 29-44 (2009).
8. T. Hayami and S. Owa, *Applications of Hankel Determinant for  $p$ -valent starlike and convex functions of order  $\alpha$* , Far East J. Appl. Math., 46(1),1-23(2010).
9. A. Janteng, S. A. Halim and M. Darus, *Hankel determinant for functions starlike and convex with respect to symmetric points*, Journal of Quality Mea surement and Analysis, 2(1), 37-43 (2006).
10. A. Janteng, S. A. Halim and M. Darus, *Coefficient inequality for a function whose derivative has a positive real part*, JIPAM. J. Inequal. Pure Appl. Math. 7(2), Article 50, 5 pp (2006).
11. A. Janteng, S. A. Halim and M. Darus, *Hankel Determinant for starlike and convex functions*, Int. J. Math.Anal. 1(13), 619-625 (2007).
12. F. R. Keogh and E. P. Merkes, *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. 20, 8-12 (1969).
13. R. J. Libera and E. J. Zł otkiewicz, *Early coefficients of the inverse of a regular convex function*, Proc. Amer. Math. Soc. 85, no. 2, 225-230,(1982).
14. R. J. Libera and E. J. Zł otkiewicz, *Coefficient bounds for the inverse of a function with derivative in  $P$* , Proc. Amer. Math. Soc. 87, no. 2, 251-257, (1983).

15. T.H. Mac Gregor, *Functions whose derivative have a positive real part*, Trans. Amer. Math. Soc.104(3), 532-537 (1962).
16. A. K. Mishra and P. Gochchayat, *second Hankel determinant for a class of Analytic functions Defined by Fractional Derivative*, Int. J. Math. Sci., vol 2008, Article ID 153280,1-10 (2008).
17. G. Murugusundaramoorthy and N. Magesh, *Coefficient inequalities for certain classes of analytic functions associated with Hankel determinant*, Bull. Math. Anal. Appl. 1, no. 3, 85–89 (2009).
18. J. W. Noonan and D. K. Thomas, *On the second Hankel determinant of areally mean  $p$ -valent functions*, Trans. Amer. Math. Soc. 223 , 337–346 (1976).
19. C. Pommerenke, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
20. L. Spaček, *Contribution à la theorie des fonctions univalentes*, Časopis Pěst. Mat. 62, 12–19 (1932).
21. D. Vamshee Krishna and T. Ramreddy, *Hankel determinant for  $p$ -valent starlike and convex functions of order  $\alpha$* , Novi Sad J. Math, 42(2), 89-102 (2012).
22. D. Vamshee Krishna and T. Ramreddy, *Coefficient inequality for certain subclasses of analytic functions associated with Hankel Determinant*, Indian J. Pure Appl. Math., 46(1), 91-106 (2015).

**Source of support: Nil, Conflict of interest: None Declared.**

**[Copy right © 2022. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**