

HYPER E-BANHATTI INDICES OF CERTAIN NETWORKS

V. R. KULLI*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

(Received On: 15-11-22; Revised & Accepted On: 05-12-22)

ABSTRACT

We introduce the first and second hyper E-Banhatti indices and their corresponding polynomials of a graph. In this paper, we compute these newly defined hyper E-Banhatti indices of some standard classes of graphs. We also determine the first and second hyper E-Banhatti indices and their corresponding polynomials for wheel graphs, friendship graphs, silicate and honeycomb networks.

Mathematics Subject Classification: 05C05, 05C12, 05C35.

Keywords: E-Banhatti index, hyper E-Banhatti index, hyper E-Banhatti polynomial, network.

1. INTRODUCTION

Throughout this paper, we consider simple graphs which are finite, connected, undirected graphs without loops and multiple edges. Let G be such a graph with vertex set $V(G)$ and edge set $E(G)$. The degree $d_G(u)$ of a vertex u is the number of vertices adjacent to u . The edge e connecting the vertices u and v is denoted by uv . If $e=uv$ is an edge of G , then the vertex u and edge e are incident as are v and e . Let $d_G(e)$ denote the degree of an edge e in G , which is defined by $d_G(e) = d_G(u) + d_G(v) - 2$ with $e=uv$. For term and concept not given here, we refer [1].

A molecular graph is a simple graph, representing the carbon atom skeleton of an organic molecule of the hydrocarbon. Therefore the vertices of a molecular graph represent the carbon atoms and its edges the carbon-carbon bonds. Chemical Graph Theory is a branch of Mathematical Chemistry which has an important effect on the development of Chemical Sciences. Several graph indices [2] have found some applications in Chemistry, especially in QSPR/QSAR research [3, 4, 5].

In [6], Kulli defined the Bhanatti degree of a vertex u of a graph G as

$$B(u) = \frac{d_G(e)}{n - d_G(u)},$$

where n is the number of vertices of G and the vertex u and edge e are incident in G .

In [6], Kulli proposed the first and second E-Banhatti indices of a graph G and they are defined as

$$EB_1(G) = \sum_{uv \in E(G)} [B(u) + B(v)],$$

$$EB_2(G) = \sum_{uv \in E(G)} B(u)B(v).$$

We now introduce the first and second hyper E-Banhatti indices of a graph G and they are defined as

$$HEB_1(G) = \sum_{uv \in E(G)} [B(u) + B(v)]^2,$$

$$HEB_2(G) = \sum_{uv \in E(G)} [B(u)B(v)]^2.$$

Corresponding Author: V. R. Kulli*

Department of Mathematics, Gulbarga University, Gulbarga - 585106, India.

Considering the first and second hyper E-Banhatti indices, we define the first and second hyper E-Banhatti polynomials of a graph G as

$$HEB_1(G, x) = \sum_{uv \in E(G)} x^{[B(u)+B(v)]^2},$$

$$HEB_2(G, x) = \sum_{uv \in E(G)} x^{[B(u)B(v)]^2}.$$

In Graph Index Theory, several graph indices were introduced and studied such as the Wiener index [7, 8, 9, 10], the Zagreb indices [11, 12, 13, 14], the Revan indices [15, 16, 17, 18], the reverse indices [19, 20, 21, 22], the Banhatti indices [23, 24, 25, 26], and the Gourava indices [27, 28, 29, 30, 31].

In this paper, we compute the first and second hyper E-Banhatti indices and their corresponding polynomials for wheel graphs, friendship graphs, silicate networks and honeycomb networks.

2. RESULTS FOR SOME STANDARD GRAPHS

2.1. First Hyper E-Banhatti Index

Proposition 1: If G is an r -regular graph with n vertices and $r \geq 2$, then

$$HEB_1(G) = \frac{8nr(r-1)^2}{(n-r)^2}.$$

Proof: Let G be an r -regular graph with n vertices and $r \geq 2$. Then G has $\frac{nr}{2}$ edges. For any edge $uv = e$ in G ,

$$d_G(e) = d_G(u) + d_G(v) - 2 = 2r - 2.$$

From definition we have

$$HEB_1(G) = \sum_{uv \in E(G)} [B(u) + B(v)]^2 = \frac{nr}{2} \left[\frac{2r-2}{n-r} + \frac{2r-2}{n-r} \right]^2 = \frac{8nr(r-1)^2}{(n-r)^2}.$$

Corollary 1.1: Let C_n be a cycle with $n \geq 3$ vertices. Then

$$HEB_1(C_n) = \frac{16n}{(n-2)^2}.$$

Corollary 1.2: Let K_n be a complete graph with $n \geq 3$ vertices. Then

$$HEB_1(K_n) = 8n(n-1)(n-2)^2.$$

Proposition 2: Let P_n be a path with $n \geq 3$ vertices. Then

$$HEB_1(P_n) = 2 \left[\frac{1}{n-1} + \frac{2}{n-2} \right]^2 + (n-3) \left[\frac{2}{n-2} + \frac{2}{n-2} \right]^2$$

$$= \frac{2(3n-4)^2}{(n-1)^2(n-2)^2} + \frac{16(n-3)}{(n-2)^2}.$$

Proposition 3: Let $K_{m,n}$ be a complete bipartite graph with $1 \leq m \leq n$ and $n \geq 2$. Then

$$HEB_1(K_{m,n}) = \frac{1}{mn} [(m+n)(m+n-2)]^2.$$

Proof: Let $K_{m,n}$ be a complete bipartite m n graph with $m+n$ vertices and mn edges such that $|V_1| = m$, $|V_2| = n$, $V(K_{r,s}) = V_1 \cup V_2$ for $1 \leq m \leq n$, and $n \geq 2$. Every vertex of V_1 is incident with n edges and every vertex of V_2 is incident with m edges. Then $d_G(e) = d_G(u) + d_G(v) - 2 = m + n - 2$.

$$HEB_1(K_{m,n}) = \sum_{uv \in E(G)} [B(u) + B(v)]^2 = mn \left[\frac{m+n-2}{m+n-n} + \frac{m+n-2}{m+n-m} \right]^2$$

$$= \frac{1}{mn} [(m+n)(m+n-2)]^2.$$

Corollary 3.1: Let $K_{n,n}$ be a complete bipartite graph with $n \geq 2$. Then

$$HEB_1(K_{n,n}) = 16(n-1)^2.$$

Corollary 3.2: Let $K_{1,n}$ be a star with $n \geq 2$. Then

$$HEB_1(K_{1,n}) = \frac{1}{n}(n^2 - 1)^2.$$

2.2. Second Hyper E-Banhatti Index

Proposition 4: If G is an r -regular graph with n vertices and $r \geq 2$, then

$$HEB_2(G) = \frac{8nr(r-1)^4}{(n-r)^4}.$$

Proof: Let G be an r -regular graph with n vertices and $r \geq 2$. Then G has $\frac{nr}{2}$ edges. For any edge $uv = e$ in G ,
 $d_G(e) = d_G(u) + d_G(v) - 2 = 2r - 2$.

From definition we have

$$HEB_2(G) = \sum_{uv \in E(G)} [B(u) \times B(v)]^2 = \frac{nr}{2} \left[\frac{2r-2}{n-r} \times \frac{2r-2}{n-r} \right]^2 = \frac{8nr(r-1)^4}{(n-r)^4}.$$

Corollary 4.1: Let C_n be a cycle with $n \geq 3$ vertices. Then

$$HEB_2(C_n) = \frac{16n}{(n-2)^4}.$$

Corollary 4.2: Let K_n be a complete graph with $n \geq 3$ vertices. Then

$$HEB_2(K_n) = 8n(n-1)(n-2)^4.$$

Proposition 5: Let P_n be a path with $n \geq 3$ vertices. Then

$$\begin{aligned} HEB_2(P_n) &= 2 \left[\frac{1}{n-1} \times \frac{2}{n-2} \right]^2 + (n-3) \left[\frac{2}{n-2} \times \frac{2}{n-2} \right]^2 \\ &= \frac{2(3n-4)^2}{(n-1)^2(n-2)^2} + \frac{16(n-3)}{(n-2)^2}. \end{aligned}$$

Proposition 6: Let $K_{m,n}$ be a complete bipartite graph with $1 \leq m \leq n$ and $n \geq 2$. Then

$$HEB_2(K_{m,n}) = \frac{(m+n-2)^4}{mn}.$$

Proof: Let $K_{m,n}$ be a complete bipartite graph with $m+n$ vertices and mn edges such that $|V_1| = m$, $|V_2| = n$,
 $V(K_{r,s}) = V_1 \cup V_2$ for $1 \leq m \leq n$, and $n \geq 2$. Every vertex of V_1 is incident with n edges and every vertex of V_2 is incident with m edges. Then $d_G(e) = d_G(u) + d_G(v) - 2 = m + n - 2$.

$$\begin{aligned} HEB_2(K_{m,n}) &= \sum_{uv \in E(G)} [B(u) \times B(v)]^2 \\ &= mn \left[\frac{m+n-2}{m+n-n} \times \frac{m+n-2}{m+n-m} \right]^2 = \frac{(m+n-2)^4}{mn}. \end{aligned}$$

Corollary 6.1: Let $K_{n,n}$ be a complete bipartite graph with $n \geq 2$. Then

$$HEB_2(K_{n,n}) = \frac{16(n-1)^4}{n^2}.$$

Corollary 6.2: Let $K_{1,n}$ be a star with $n \geq 2$. Then

$$HEB_2(K_{1,n}) = \frac{(n-1)^4}{n}.$$

3. RESULTS FOR FRIENDSHIP GRAPHS

A friendship graph F_n , $n \geq 2$, is a graph that can be constructed by joining n copies of C_3 with a common vertex. A graph F_4 is shown in Figure 1.




Figure-1: Friendship graph F_4

Let F_n be a friendship graph with $2n+1$ vertices and $3n$ edges. By calculation, we obtain that there are two types of edges as follows:

$$E_1 = \{uv \in E(F_n) \mid d_{F_n}(u) = d_{F_n}(v) = 2\}, \quad |E_1| = n.$$

$$E_2 = \{uv \in E(F_n) \mid d_{F_n}(u) = 2, d_{F_n}(v) = 2n\}, \quad |E_2| = 2n.$$

Therefore, in F_n , there are two types of Banhatti edges based on Banhatti degrees of end vertices of each edge follow:

$$BE_1 = \{uv \in E(F_n) \mid B(u) = B(v) = \frac{2}{2n-1}\}, \quad |BE_1| = n.$$

$$BE_2 = \{uv \in E(F_n) \mid B(u) = \frac{2n}{2n-1}, B(v) = 2n\}, \quad |BE_2| = 2n.$$

We now compute the first hyper E-Banhatti index of a friendship graph F_n .

Theorem 1: Let F_n be a friendship graph with $2n + 1$ vertices and $3n$ edges. Then

$$HEB_1(F_n) = \frac{32n^5 + 16n}{(2n-1)^2}.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of F_n , we obtain

$$\begin{aligned} HEB_1(F_n) &= \sum_{uv \in E(F_n)} [B(u) + B(v)]^2 = n \left(\frac{2}{2n-1} + \frac{2}{2n-1} \right)^2 + 2n \left(\frac{2n}{2n-1} + 2n \right)^2 \\ &= \frac{32n^5 + 16n}{(2n-1)^2}. \end{aligned}$$

In the following theorem, we obtain the second hyper E-Banhatti index of a friendship graph F_n .

Theorem 2: Let F_n be a friendship graph with $2n + 1$ vertices and $3n$ edges. Then

$$HEB_2(F_n) = \frac{16n}{(2n-1)^4} + \frac{32n^5}{(2n-1)^2}.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of F_n , we obtain

$$\begin{aligned} HEB_2(F_n) &= \sum_{uv \in E(F_n)} [B(u)B(v)]^2 = n \left(\frac{2}{2n-1} \times \frac{2}{2n-1} \right)^2 + 2n \left(\frac{2n}{2n-1} \times 2n \right)^2 \\ &= \frac{16n}{(2n-1)^4} + \frac{32n^5}{(2n-1)^2}. \end{aligned}$$

By using definitions and by cardinalities of the Banhatti edge partition of F_n , we obtain the first and second hyper E-Banhatti polynomials of F_n .

Theorem 3: The first hyper E-Banhatti polynomial of F_n is given by

$$HEB_1(F_n, x) = nx^{\left(\frac{4}{2n-1}\right)^2} + 2nx^{\left(\frac{4n^2}{2n-1}\right)^2}.$$

Theorem 4: The second hyper E-Banhatti polynomial of F_n is given by

$$HEB_2(F_n, x) = nx^{\left(\frac{2}{2n-1}\right)^4} + 2nx^{\left(\frac{4n^2}{2n-1}\right)^2}.$$

4. RESULTS FOR WHEEL GRAPHS

A wheel graph W_n is the join of C_n and K_1 . Then W_n has $n+1$ vertices and $2n$ edges. A graph W_n is presented in Figure 2.




Figure-2: Wheel graph W_n

In W_n , there are two types of edges as follows:

$$E_1 = \{uv \in E(W_n) \mid d(u) = d(v) = 3\}, \quad |E_1| = n.$$

$$E_2 = \{uv \in E(W_n) \mid d(u) = 3, d(v) = n\}, \quad |E_2| = n.$$

Therefore, in W_n , there are two types of Banhatti edges based on Banhatti degrees of end vertices of each edge follow:

$$BE_1 = \left\{ uv \in E(W_n) \mid B(u) = B(v) = \frac{4}{(n-2)} \right\}, \quad |BE_1| = n.$$

$$BE_2 = \left\{ uv \in E(W_n) \mid B(u) = \frac{n+1}{n-2}, B(v) = n+1 \right\}, \quad |BE_2| = n.$$

We determine the first hyper E-Banhatti index of a wheel graph W_n .

Theorem 5: Let W_n be a wheel graph with $n+1$ vertices and $2n$ edges. Then

$$HEB_1(W_n) = \frac{64n}{(n-2)^2} + \frac{n(n^2-1)^2}{(n-2)^2}.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of W_n , we obtain

$$\begin{aligned} HEB_1(W_n) &= \sum_{uv \in E(W_n)} [B(u) + B(v)]^2 = n \left(\frac{4}{n-2} + \frac{4}{n-2} \right)^2 + n \left(\frac{n+1}{n-2} + n+1 \right)^2 \\ &= \frac{64n}{(n-2)^2} + \frac{n(n^2-1)^2}{(n-2)^2}. \end{aligned}$$

In the next theorem, we compute the second hyper E-Banhatti index of a wheel graph W_n .

Theorem 6: Let W_n be a wheel graph with $n+1$ vertices and $2n$ edges. Then

$$HEB_2(W_n) = \frac{256n}{(n-2)^4} + \frac{n(n+1)^2}{(n-2)^2}.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of W_n , we obtain

$$\begin{aligned} HEB_2(W_n) &= \sum_{uv \in E(W_n)} [B(u)B(v)]^2 = n \left(\frac{4}{n-2} \times \frac{4}{n-2} \right)^2 + n \left(\frac{n+1}{n-2} \times (n+1) \right)^2 \\ &= \frac{256n}{(n-2)^4} + \frac{n(n+1)^2}{(n-2)^2}. \end{aligned}$$

By using definitions and by cardinalities of the Banhatti edge partition of W_n , we obtain the first and second hyper E-Banhatti polynomials of W_n .

Theorem 7: The first hyper E-Banhatti polynomial of W_n is given by

$$HEB_1(W_n, x) = nx^{\left(\frac{8}{n-2}\right)^2} + nx^{\left(\frac{n^2-1}{n-2}\right)^2}.$$

Theorem 8: The second hyper E-Banhatti polynomial of W_n is given by

$$HEB_2(W_n, x) = nx^{\frac{256}{(n-2)^4}} + nx^{\frac{(n+1)^2}{(n-2)^2}}.$$

5. RESULTS FOR SILICATE NETWORKS

Silicates are obtained by fusing metal oxide or metal carbonates with sand. A silicate network is symbolized by SL_n , where n is the number of hexagons between the center and boundary of SL_n . A 2-dimensional silicate network is depicted in Figure 3.




Figure-3: A 2-dimensional silicate network

Let G be the graph of a silicate network SL_n . By calculation, we obtain that G has $15n^2 + 3n$ vertices and $36n^2$ edges. In G , by calculation, there are three types of edges based on the degree of end vertices of each edge as follows:

$$\begin{aligned} E_1 &= \{uv \in E(SL_n) \mid d_G(u) = d_G(v) = 3\}, & |E_1| &= 6n. \\ E_2 &= \{uv \in E(SL_n) \mid d_G(u) = 3, d_G(v) = 6\}, & |E_2| &= 18n^2 + 6n. \\ E_3 &= \{uv \in E(SL_n) \mid d_G(u) = d_G(v) = 6\}, & |E_3| &= 18n^2 - 12n. \end{aligned}$$

Therefore, in SL_n , there are three types of Banhatti edges based on Banhatti degrees of end vertices of each edge as follow:

$$\begin{aligned} BE_1 &= \left\{ uv \in E(SL_n) \mid B(u) = B(v) = \frac{4}{(15n^2 + 3n - 3)} \right\}, & |BE_1| &= 6n. \\ BE_2 &= \left\{ uv \in E(SL_n) \mid B(u) = \frac{7}{15n^2 + 3n - 3}, B(v) = \frac{7}{15n^2 + 3n - 6} \right\}, & |BE_2| &= 18n^2 + 6n. \\ BE_3 &= \left\{ uv \in E(SL_n) \mid B(u) = B(v) = \frac{10}{15n^2 + 3n - 6} \right\}, & |BE_3| &= 18n^2 - 12n. \end{aligned}$$

In Theorem 9, we establish the first hyper E-Banhatti index of a silicate network SL_n .

Theorem 9: Let SL_n be a silicate network. Then

$$HEB_1(SL_n) = 6n \left(\frac{8}{15n^2 + 3n - 3} \right)^2 + (18n^2 + 6n) \left(\frac{7}{15n^2 + 3n - 3} + \frac{7}{15n^2 + 3n - 6} \right)^2 + (18n^2 - 12n) \left(\frac{20}{15n^2 + 3n - 6} \right)^2.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of SL_n , we obtain

$$HEB_1(SL_n) = \sum_{uv \in E(SL_n)} [B(u) + B(v)]^2 = 6n \left(\frac{4}{15n^2 + 3n - 3} + \frac{4}{15n^2 + 3n - 3} \right)^2 + (18n^2 + 6n) \left(\frac{7}{15n^2 + 3n - 3} + \frac{7}{15n^2 + 3n - 6} \right)^2 + (18n^2 - 12n) \left(\frac{10}{15n^2 + 3n - 6} + \frac{10}{15n^2 + 3n - 6} \right)^2.$$

After simplification, we get the desired result.

In the following theorem, we obtain the second hyper E-Banhatti index of a silicate network SL_n .

Theorem 10: Let SL_n be a silicate network. Then

$$HEB_2(SL_n) = 6n \left(\frac{4}{15n^2 + 3n - 3} \right)^4 + (18n^2 + 6n) \left(\frac{49}{(15n^2 + 3n - 3)(15n^2 + 3n - 6)} \right)^2 + (18n^2 - 12n) \left(\frac{10}{15n^2 + 3n - 6} \right)^4.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of SL_n , we obtain

$$HEB_2(SL_n) = \sum_{uv \in E(SL_n)} [B(u)B(v)]^2 = 6n \left(\frac{4}{(15n^2 + 3n - 3)} \times \frac{4}{(15n^2 + 3n - 3)} \right)^2 + (18n^2 + 6n) \left(\frac{7}{15n^2 + 3n - 3} \times \frac{7}{15n^2 + 3n - 6} \right)^2 + (18n^2 - 12n) \left(\frac{10}{15n^2 + 3n - 6} \times \frac{10}{15n^2 + 3n - 6} \right)^2$$

gives the desired result after simplification.

By using definitions and by cardinalities of the Banhatti edge partition of SL_n , we obtain the first and second hyper E-Banhatti polynomials of SL_n .

Theorem 11: The first hyper E-Banhatti polynomial of SL_n is given by

$$HEB_1(SL_n, x) = 6nx \left(\frac{8}{15n^2 + 3n - 3} \right)^2 + (18n^2 + 6n)x \left(\frac{7}{15n^2 + 3n - 3} + \frac{7}{15n^2 + 3n - 6} \right)^2 + (18n^2 - 12n)x \left(\frac{20}{15n^2 + 3n - 6} \right)^2.$$

Theorem 12: The second hyper E-Banhatti polynomial of SL_n is given by

$$HEB_2(SL_n, x) = 6nx \left(\frac{4}{15n^2 + 3n - 3} \right)^4 + (18n^2 + 6n)x \left(\frac{49}{(15n^2 + 3n - 3)(15n^2 + 3n - 6)} \right)^2 + (18n^2 - 12n)x \left(\frac{10}{15n^2 + 3n - 6} \right)^4.$$

6. RESULTS FOR HONEYCOMB NETWORKS

Honeycomb networks are useful in Computer Graphics and Chemistry. A honeycomb network of dimension n is denoted by HC_n , where n is the number of hexagons between central and boundary hexagon. A 4-dimensional honeycomb network is shown in Figure 4.




Figure-4: A 4-dimensional honeycomb network

Let G be the graph of a honeycomb network HC_n . By calculation, we obtain that G has $6n^2$ vertices and $9n^2 - 3n$ edges. In G , by algebraic method, there are three types of edges based on the degree of end vertices of each edge as follows:

$$\begin{aligned} E_1 &= \{uv \in E(HC_n) \mid d_G(u) = d_G(v) = 2\}, & |E_1| &= 6. \\ E_2 &= \{uv \in E(HC_n) \mid d_G(u) = 2, d_G(v) = 3\}, & |E_2| &= 12n - 12. \\ E_3 &= \{uv \in E(HC_n) \mid d_G(u) = d_G(v) = 3\}, & |E_3| &= 9n^2 - 15n + 6. \end{aligned}$$

Therefore, in HC_n , there are three types of Banhatti edges based on Banhatti degrees of end vertices of each edge as follow:

$$\begin{aligned} BE_1 &= \left\{ uv \in E(HC_n) \mid B(u) = B(v) = \frac{2}{6n^2 - 2} \right\}, & |BE_1| &= 6. \\ BE_2 &= \left\{ uv \in E(HC_n) \mid B(u) = \frac{2}{6n^2 - 2}, B(v) = \frac{3}{6n^2 - 3} \right\}, & |BE_2| &= 12n - 12. \\ BE_3 &= \left\{ uv \in E(HC_n) \mid B(u) = B(v) = \frac{4}{6n^2 - 3} \right\}, & |BE_3| &= 9n^2 - 15n + 6. \end{aligned}$$

We now compute the first hyper E-Banhatti index of a honeycomb network HC_n .

Theorem 13: Let HC_n be a honeycomb network. Then

$$HEB_1(HC_n) = 6 \left(\frac{4}{6n^2 - 2} \right)^2 + (12n - 12) \left(\frac{2}{6n^2 - 2} + \frac{3}{6n^2 - 3} \right)^2 + (9n^2 - 15n + 6) \left(\frac{8}{6n^2 - 3} \right)^2.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of HC_n , we obtain

$$\begin{aligned} HEB_1(HC_n) &= \sum_{uv \in E(HC_n)} [B(u) + B(v)]^2 \\ &= 6 \left(\frac{2}{6n^2 - 2} + \frac{2}{6n^2 - 2} \right)^2 + (12n - 12) \left(\frac{2}{6n^2 - 2} + \frac{3}{6n^2 - 3} \right)^2 \\ &\quad + (9n^2 - 15n + 6) \left(\frac{4}{6n^2 - 3} + \frac{4}{6n^2 - 3} \right)^2. \end{aligned}$$

After simplification, we obtain the desired result.

We determine the second hyper E-Banhatti index of a honeycomb network HC_n .

Theorem 14: Let HC_n be a honeycomb network. Then

$$HEB_2(HC_n) = 6 \left(\frac{2}{6n^2 - 2} \right)^4 + (12n - 12) \left(\frac{6}{(6n^2 - 2)(6n^2 - 3)} \right)^2 + (9n^2 - 15n + 6) \left(\frac{4}{6n^2 - 3} \right)^4.$$

Proof: From definition and by cardinalities of the Banhatti edge partition of HC_n , we obtain

$$\begin{aligned} HEB_2(HC_n) &= \sum_{uv \in E(HC_n)} [B(u) B(v)]^2 \\ &= 6 \left(\frac{2}{6n^2 - 2} \times \frac{2}{6n^2 - 2} \right)^2 + (12n - 12) \left(\frac{2}{6n^2 - 2} \times \frac{3}{6n^2 - 3} \right)^2 \\ &\quad + (9n^2 - 15n + 6) \left(\frac{4}{6n^2 - 3} \times \frac{4}{6n^2 - 3} \right)^2 \end{aligned}$$

gives the desired result after simplification.

By using definitions and by cardinalities of the Banhatti edge partition of HC_n , we obtain the first and second hyper E-Banhatti polynomials of HC_n .

Theorem 15: The first hyper E-Banhatti polynomial of HC_n is given by

$$HEB_1(HC_n, x) = 6x^{\left(\frac{4}{6n^2-2}\right)^2} + (12n-12)x^{\left(\frac{2}{6n^2-2} + \frac{3}{6n^2-3}\right)^2} + (9n^2-15n+6)x^{\left(\frac{8}{6n^2-3}\right)^2}.$$

Theorem 16: The second hyper E-Banhatti polynomial of HC_n is given by

$$HEB_2(HC_n, x) = 6x^{\left(\frac{2}{6n^2-2}\right)^4} + (12n-12)x^{\left(\frac{6}{(6n^2-2)(6n^2-3)}\right)^2} + (9n^2-15n+6)x^{\left(\frac{4}{6n^2-3}\right)^4}.$$

7. CONCLUSION

In this study, we have introduced the first and second hyper E-Banhatti indices of a graph. Furthermore, we have determined these newly defined indices for some standard graphs, wheel graphs, friendship graphs and certain networks. This study is a new direction in The Theory of Graph Index in Graphs.

Many questions are suggested by this research, among them are the following:

1. Obtain properties of the first and second hyper E-Banhatti indices.
2. Compute these two indices for other chemical nanostructures.

REFERENCES

1. V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. V.R.Kulli, Graph indices, in *Hand Book of Research on Advanced Applications of Application Graph Theory in Modern Society*, M. Pal. S. Samanta and A. Pal, (eds.) IGI Global, USA (2020) 66-91.
3. R.Todeschini and V. Consonni, *Molecular Descriptors for Chemoinformatics*, Wiley-VCH, Weinheim, (2009).
4. I.Gutman and O.E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer, Berlin (1986).
5. V.R.Kulli, *Multiplicative connectivity indices of nanostructures*, LAP LEMBERT Academic Publishing (2018).
6. V.R.Kulli, New direction in the theory of graph index in graphs, submitted
7. H.Wiener, Structural determination of paraffin boiling points, *Journal of the American Chemical Society*, 69(1) (1947) 17-20.
8. A.A.Dobrynin, R.Entringer and I.Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.* 66 (2001) 211-249.
9. H.S.Ramane, K.P.Narayakar, S.S.Shirkol and A.B.Ganagi, Terminal Wiener index of line graphs, *MATCH Commun. Math. Comput. Chem.* 69 (2013) 775-782.
10. F.Yang, Z-D Wang and Y-P Huang, Modification of the Wiener index 4, *Journal of Computational Chemistry*, 25(6) (2004) 881-887.
11. I. Gutman and N. Trinajstić, Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* vol. 17, pp 535-538, 1972.
12. G.Caporossi, P.Hansen and D.Vukicevic, Comparing Zagreb indices of cyclic graphs, *MATCH Commun. Math. Comput. Chem.*, 63 (2010) 441-451.
13. K.C.Das, I.Gutman and B.Horoldagva, Comparing Zagreb indices and coindices of trees, *MATCH Commun. Math. Comput. Chem.*, 67 (2012) 189-198.
14. T.Doslic, B.Furtula, A.Graovac, I.Gutman, S.Moradi and Z.Yarahmadi, On vertex degree based molecular structure descriptors, *MATCH Commun. Math. Comput. Chem.*, 66 (2011) 613-626.
15. V.R. Kulli, Revan indices of oxide and honeycomb networks, *International Journal of Mathematics and its Applications*, 5(4-E) (2017) 663-667.
16. A.Q.Baig, M.Nadeem and W.Gao, Revan and hyper Revan indices of Octahedral and icosahedral networks, *Applied Mathematics and Nonlinear Sciences*, 3(1) (2018) 33-40.
17. P.Kandan, E.Chandrasekaran and M.Priyadharshini, The Revan weighted Szeged index of graphs, *Journal of Emerging Technologies and Innovative Research*, 5(9) (2018) 358-366.
18. R.Aguilar-Sanchez, I.F.Herrera-Gonzalez, J.A.Mendez-Bermudez and J.M.Sigarreta, Revan degree indices on random graphs, arXiv:2210.04749v1 [math.CO] 10 Oct 2022.
19. S.Ediz, Maximal graphs of the first reverse Zagreb beta index, *TWMS J. Appl. Eng. Math.* 8 (2018) 306-310.
20. S.Ediz and M.Canacan, Reverse Zagreb indices of cartesian product of graphs, *International Journal of Mathematics and Computer Science*, 11(1) (2016) 51-58.
21. W.Gao, M.Younas, A.Farooq, A ur R Virk and W.Nazeer, Some reverse degree based topological indices and polynomials of dendrimers, *Mathematics* 2018, 6, 214; doi:10.3390/math6100214.

22. V.R.Kulli, On the sum connectivity reverse index of oxide and honeycomb networks, *Journal of Computer and Mathematical Sciences*, 8(9) (2017) 408-413.
23. V.R.Kulli, On K Banhatti indices of graphs, *Journal of Computer and Mathematical Sciences*, 7 (2016) 213-218.
24. I.Gutman, V.R.Kulli, B.Chaluvaraju and H.S.Boregowda, On Banhatti and Zagreb indices, *Journal of the international Mathematical Virtual Institute*, 7 (2017) 53-67.
25. W.Wang, M.Naeem, A.Rauf, A.Riasat, A.Aslam and K.A.Yannick, On analysis of Banhatti indices for hyaluronic acid curcumin and hydroxychloroquine, *Journal of Chemistry*, 2021 Article ID 7468857 10 pages.
26. D.Zhao, M.A.Zahid, R.Irfan, M.Arshad, A.Fahad, Z.Ahmad and L.Li, Banhatti, Revan and hyper indices of silicon carbide $Si_2C_3-III[n, m]$, *Open Chemistry*, 19 (2021) 646-652.
27. V.R.Kulli, The Gourava indices and coindices of graphs, *Annals of Pure and Applied Mathematics*, 14(1) (2017).
28. M.Aruvi, M.Joseph and E.Ramganes, The second Gourava index of some graph products, *Advances in Mathematics: Scientific Journal* 9(12) (2020) 10241-10249.
29. B.Basavanagoud and S.Policepatil, Chemical applicability of Gourava and hyper Gourava indices, *Nanosystems: Physics, Chemistry, Mathematics* 12(2) (2021) 142-150.
30. V.R.Kulli, V.Lokesha, S.Jain and M.Manjunath, The Gourava index of four operations on graphs, *International J. Math. Combin.* 4 (2018) 65-76.
31. V.R.Kulli, G.N.Adithya and N.D.Soner, Gourava indices of certain windmill graphs, *International Journal of Mathematics Trends and Technology*, 68(9) (2022) 51-59.

Source of support: Nil, Conflict of interest: None Declared.

[Copy right © 2022. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]