



## ANNIHILATOR INJECTIVE MODULES

Himanshu Ranjan\*, B. M. Pandeya and A. J. Gupta

Department of Applied Mathematics, Institute of Technology,  
Banaras Hindu University, Varanasi-221005 (India)

E-mail: [hranjan.rs.apm@itbhu.ac.in](mailto:hranjan.rs.apm@itbhu.ac.in)

(Received on: 18-10-11; Accepted on: 03-11-11)

### ABSTRACT

Annihilator Injective Module is a generalization of Injective Modules, in this paper we give several properties of Annihilator Injective Modules and discuss the question when Annihilator Injective is Injective Module.

**Key words:** Simple Module, Semi simple, Injective Module, Annihilator injective Modules,  $l_M(R)$  and  $r_R(X)$ .

(2000)AMS Mathematics subject classification: 16D50, 16D60, 16D80

### INTRODUCTION:

Throughout this paper  $R$  represent an associative ring with identity  $1 \neq 0$  and all modules are unitary  $R$ -module we write  $M_R$  (res,  ${}_R M$ ) to indicate that  $M$  is a right (res, left)  $R$ -module, we also write  $J, Z_r, S_r$  for the Jacobson radical, the right singular ideal and the right socal respectively of  $R$  and  $E(M_R)$  the injective hull of  $M_R$ , If  $X$  is a subset of  $M$  then the right annihilator of  $X$  in  $R$  denoted by  $r_R(X)$  and if  $A$  is subset of  $R$  then the set  $l_M(A)$  is called left annihilator of  $A$ , If  $N$  is a submodule of  $M$  (res, proper submodule) we denoted by  $N \leq M$  (res,  $N \prec M$ ) moreover we write  $N \leq^e M, N \ll M, N \leq^{\oplus} M$ , and  $N \leq^{\max} M$  to indicate that  $N$  is an essential submodule, a small submodule, a direct summand and a maximal submodule of  $M$  respectively, A module  $M$  is called uniform if  $M \neq 0$  and every non zero submodule of  $M$  is essential in  $M$ , Let  $M$  and  $N$  be  $R$ -Modules  $N$  is called  $M$ -Injective Module if for every submodule  $A$  of  $M$  and the inclusion map  $i: A \rightarrow M$  any homomorphism  $\alpha: A \rightarrow N$  can be extended to a homomorphism  $\beta: M \rightarrow N$  such that the diagram is commutative.

### 1. PRELIMINARIES:

**Definition: 1.1** Let  $M$  be an  $R$ -Module and  $X \subseteq M$  then set  $r_R(X) = \{r \in R : xr = 0 \forall x \in X\}$  is called the right annihilator of  $X$  in  $R$ .

**Definition: 1.2** For each  $A \subseteq R$  the set  $l_M(A) = \{m \in M : ma = 0 \forall a \in A\}$  called left annihilator of  $A$  in  $M$ , for a singleton set  $\{x\}$  and  $\{a\}$  we denote the annihilator as  $r_R(x)$  and  $l_M(a)$  respectively.

**Proposition: 1.3** Let  $M$  be an  $R$ -Module and  $X \subseteq M$  and  $A \subseteq R$  then,

1.  $r_R(X)$  is a right ideal of  $R$
2.  $l_M(A)$  is a submodule of  $M$

**Definition: 1.4**  $R$  modules  $N$  are called injective if it is  $M$  injective for every  $R$ -Module.

**Definition: 1.5**  $R$  modules  $N$  is called quasi injective if  $N$  is  $N$  injective.

\*Corresponding author: Himanshu Ranjan\*, \*E-mail: [hranjan.rs.apm@itbhu.ac.in](mailto:hranjan.rs.apm@itbhu.ac.in)

**Definition: 1.6**  $M$  and  $N$  be  $R$ -modules, then  $N$  is called  $M$  **Annihilator Injective Module** if the inclusion map  $i: l_M(R) \rightarrow M$  and any homomorphism  $\alpha: l_M(R) \rightarrow N$  can be extended to a homomorphism  $\beta: M \rightarrow N$  such that the diagram is commutative i.e.  $\alpha = \beta i$ .

**Definition: 1.7**  $N$  is called **Annihilator Injective Module** if for all  $R$ -Module  $M$  the inclusion map  $i: l_M(R) \rightarrow M$  and any homomorphism  $\alpha: l_M(R) \rightarrow N$  can be extended to a homomorphism  $\beta: M \rightarrow N$  such that the diagram is commutative i.e.  $\alpha = \beta i$ .

### 1. ANNIHILATOR INJECTIVE MODULS:

**Proposition: 2.1** Let  $N$  and  $N'$  be  $R$ -Modules if  $N \cong N'$  and  $N$  is Annihilator Injective then  $N'$  is also Annihilator Injective.

**Proof:** Since  $N$  is annihilator Injective then for any  $R$ -module  $M$   $i: l_M(R) \rightarrow M$  be an injective map and  $\alpha: l_M(R) \rightarrow N$  be a homomorphism then there exist a homomorphism  $\beta: M \rightarrow N$  the map is commute i.e.  $\alpha = \beta i$ , let  $f: N \rightarrow N'$  be an isomorphism now define a homomorphism  $\alpha': l_M(R) \rightarrow N'$  and  $\beta': M \rightarrow N'$  where  $\alpha' = f\alpha$  and  $\beta' = f\beta$  then  $\alpha' = f\alpha = f\beta i = \beta' i$  therefore  $N'$  is annihilator injective module.

**Proposition: 2.2** If  $N_1 \oplus N_2$  is Annihilator Injective module then  $N_1$  and  $N_2$  is relatively Annihilator Injective Modules.

**Proof:** Let  $N_1 \oplus N_2$  be Annihilator Injective Module then we have to show that  $N_1$  is  $N_2$  Annihilator Injective Modue,

Let  $i: l_{N_2}(R) \rightarrow N_2$  be the inclusion map and  $\alpha: l_{N_2}(R) \rightarrow N_1 \oplus N_2$  be a homomorphism where  $\alpha(n) = (\alpha'(n), n)$  so  $\alpha$  is monomorphism since  $N_1 \oplus N_2$  is Annihilator Injective Module then there exist a homomorphism  $\beta: N_2 \rightarrow N_1 \oplus N_2$  such that  $\alpha = \beta i$ , Let  $\pi_1$  be natural projection from  $\pi_1: N_1 \oplus N_2 \rightarrow N_1$  and let  $\beta': N_2 \rightarrow N_1$  be a homomorphism where  $\beta' = \pi_1 \beta$  now define a homomorphism  $\alpha': l_{N_2}(R) \rightarrow N_1$  where  $\alpha' = \pi_1 \alpha = \pi_1 \beta i = \beta' i$  therefore  $N_1$  is  $N_2$  Annihilator Injective Modules similarly we can show that  $N_2$  is  $N_1$  Annihilator Injective Module.

**Proposition: 2.3** Let  $\{N_\lambda\}_{\lambda \in \Lambda}$  be a family of Annihilator Injective modules, then  $\prod_{\lambda \in \Lambda} N_\lambda$  is Annihilator Injective Module if and only if each  $N_\lambda$  is Annihilator Injective Modules.

**Proof:** Let  $M$  be an  $R$ -Module and put  $N = \prod_{\lambda \in \Lambda} N_\lambda$  Let  $\phi_\lambda: N_\lambda \rightarrow N$  and  $\pi_\lambda: N \rightarrow N_\lambda$  be injection and projection respectively.

Let each  $N_\lambda$  be  $M$  Annihilator Injective Module, let  $\alpha: l_M(R) \rightarrow N$  be a homomorphism and  $i: l_M(R) \rightarrow M$  be an injection map then for each  $\lambda$  there exist  $\beta_\lambda: M \rightarrow N_\lambda$  such that the diagram is commutative, therefore  $\pi_\lambda \alpha = \beta_\lambda i$  now define  $\beta: M \rightarrow N$  where  $\beta(m) = \{\beta_\lambda(m)\}_{\lambda \in \Lambda}$  ( $m \in M$ ) then  $\beta$  is an  $R$ -homomorphism and if  $x \in l_M(R)$  then  $\beta i(x) = \{\beta_\lambda i(x)\} = \{\pi_\lambda \alpha(x)\} = \alpha(x)$  which shows that  $N$  is Annihilator Injective.

#### Conversely:

Suppose that  $N$  is Annihilator Injective Module.

Consider  $i: l_M(R) \rightarrow M$  and  $\gamma: l_M(R) \rightarrow N$  be a homomorphism since  $N$  is Annihilator Injective then for any  $\text{Hom}(l_M(R), N)$  there exist a homomorphism  $\mu: M \rightarrow N$  and the map commute i.e.  $\mu i = \gamma$  now define a map

$\mu': M \rightarrow N_\lambda$  where  $\mu'(m) = \pi_\lambda \mu(m)$  then  $\mu'$  is an  $R$  homomorphism and if  $a \in l_M(R)$  then  $\mu'i(a) = \pi_\lambda \mu i(a) = \pi_\lambda \phi_\lambda \gamma(a) = \gamma(a)$  this shows that each  $N_\lambda$  is Annihilator Injective Module.

**Proposition: 2.4** let  $N_\lambda$  be a family of  $R$  -modules.

1. If  $\bigoplus_{\lambda \in \wedge} N_\lambda$  is Annihilator Injective then each  $N_\lambda$  is Annihilator Injective.
2. If the Index set  $\wedge$  is finite and each  $N_\lambda$  is Annihilator Injective then  $\bigoplus_{\lambda \in \wedge} N_\lambda$  is Annihilator Injective.

**Proof:** Proof is similar for the proposition (2.3).

**Note: 2.5** Every Injective Modules is Annihilator Injective Modules but converse is not true.

**Example: 2.6**  $Z(Z)$  is Annihilator Injective Module but it is not Injective Module.

**Proposition: 2.7** If  $N$  is Annihilator Injective Module then any monomorphism  $f: N \rightarrow M$  split.

**Proof:** Let  $f: N \rightarrow M$  be a monomorphism then  $f^{-1}: f(N) \rightarrow N$  be inverse of  $f$  since  $N$  is Annihilator Injective then there exist a homomorphism  $\beta: M \rightarrow N$  that extends  $f^{-1}$  set  $g = \beta f$  then  $g$  is clearly an identity map of  $N$  hence  $f$  splits in  $M$ .

**Proposition: 2.8** If  $N$  is Annihilator Injective module then  $N$  is Annihilator  $K$ -Injective module for any submodule  $K$  of  $M$ .

**Proof:** Let  $X$  be submodule of  $K$  and  $f: X \rightarrow N$  be a homomorphism then  $X$  be also a submodule of  $M$  and by Annihilator  $M$ -Injective module of  $N$ ,  $f$  extends to a homomorphism  $f^*: M \rightarrow N$  the restriction  $f^*/K$  of  $f^*$  on  $K$  is a homomorphism  $K \rightarrow N$  which extends  $f$  hence  $N$  is Annihilator  $K$ -Injective module.

**Proposition: 2.9** Every direct summand of Annihilator Injective Modules is also Annihilator Injective Modules.

**Proof:** Assume  $N$  is Annihilator Injective module and  $N = K \oplus K'$  let  $l_M(R)$  be a submodule of  $M$  and  $f: l_M(R) \rightarrow K$  be a homomorphism define  $g: l_M(R) \rightarrow N = K \oplus K'$  by  $g(m) = (f(m), 0)$  thus  $g$  is monomorphism and by Annihilator Injective Module of  $N$ ,  $g$  extends to a homomorphism  $\beta: M \rightarrow N$ , let  $\pi_K$  be the natural projection of  $N = K \oplus K'$  into  $K$  then  $\pi_K \beta: M \rightarrow K$  is a homomorphism which extends  $f$ , therefore  $K$  is Annihilator Injective.

**Definition:** - let  $M$  be an  $R$ -Module an element  $m$  of  $M$  is said to be divisible if for every  $r$  of  $R$  which is not a right zero divisor there exist  $m' \in M$  such that  $m = rm'$ , if every element of  $M$  is divisible then  $M$  is said to be a divisible module. An abelian group is said to be divisible if it is divisible as a  $Z$ -module. Alternatively  $M$  is divisible if  $M = rM$  whenever  $r$  is an element of  $R$  which is not a right zero divisor.

**Note 2.10** Every Injective module is divisible but every Annihilator Injective modules need not be divisible.

**Lemma 2.11** Let  $M$  be an  $R$ -Module such that every submodule of  $M$  and let  $M'$  be a submodule of  $M$  then every submodule of  $M'$  is a direct summand of  $M$ .

**Proof:** Let  $A$  be a submodule of  $M'$  then  $A$  is also a submodule of  $M$  so there is a submodule  $B$  of  $M$  such that  $M = A + (M' \cap B)$  (d. s) it follows that  $M' = A + (M' \cap B) = A \cap B = 0$ .

**Proposition: 2.12** Let  $M$  be an  $R$ -module then the following statements are equivalent.

1.  $M$  has a family  $\{S_i\}_{i \in I}$  of simple submodule such that  $M = \sum_{i \in I} S_i$  (d. s)

2.  $M$  has a family of simple submodules whose sum is  $M$  itself.
3. Every submodule of  $M$  is a direct summand of  $M$ .

**Proof:** [5, Proposition 3.2]

**Proposition: 2.13** Every simple Annihilator Injective modules is Injective Modules.

**Proof:** Simple

**ACKNOWLEDGEMENT:**

The first author would like to thank for the **AICTE** – India to provide the national doctoral fellowship.

**REFERENCE:**

- [1] Anderson, F. W., Fuller, K. R., Rings and Categories of Modules, Springer- Verlag New York, 1974.
- [2] Chaturvedi, A. K., Thesis subtitle (Variants of quasi-Injective Modules and there endomorphism Rings) 2008.
- [3] Dung, N. V., Huynh, D. V., Smith, P. F., Wisbauer, R., Extending Module, London: Pitman, 1994.
- [4] Pandeya, B. M., Koirala, S. P. (2001), Pseudo M-Injective Modules, In: Rizvi, S. T., Quadri, M. A., Ashraf, M., eds, Algebra and its Application, Narosa publishing House, p.p. 201-207.
- [5] Sharp, D. W., Vamos, P., Injective Module, Cambridge University Press, 1972.

\*\*\*\*\*