

Coefficient Bounds for Certain Subclasses of Meromorphic p-valent Functions

B. A. Frasin¹, *B. Srutha Keerthi², B. Adolf Stephen³, A. Gangadharan⁴

¹*Department of Mathematics, Al al-Bayt University, P.O.Box: 130095, Mafrqa, Jordan
E-mail: bafrasin@yahoo.com*

²*Department of Applied Mathematics, Sri Venkateswara College of Engineering,
Sriperumbudur, Chennai – 602105, India
E-mail : sruthilaya06@yahoo.co.in, laya@svce.ac.in*

³*Department of Mathematics, Madras Christian College, Chennai – 600059, India
E-mail : adolfmcc2003@yahoo.co.in*

⁴*Department of Applied Mathematics, Sri Venkateswara College of Engineering,
Sriperumbudur, Chennai - 602105, India
E-mail : ganga@svce.ac.in*

(Received on: 10-11-11; Accepted on: 24-11-11)

ABSTRACT

In this present investigation, the authors obtain sharp bounds for $|a_1 - \mu a_0^2|$ and $|a_2|$ for certain subclasses of meromorphic p-valent functions. As an application of these results, coefficients bounds for classes of functions defined through Ruscheweyh derivatives are also obtained.

2000 Mathematics Subject Classifications: Primary 30C45.

Keywords and Phrases: Meromorphic functions, starlike functions, p-valent functions, subordination, Ruscheweyh derivatives.

1. INTRODUCTION:

Let Σ_p denotes the class of functions $f(z)$ of the form

$$f(z) = z^{-p} + \sum_{k=0}^{\infty} a_k z^{k-p+1} \quad (p \in \mathbb{N} := \{1,2,3,\dots\}), \tag{1}$$

which are analytic and p-valent in the punctured unit disk

$$\Delta^* = \{z : z \in \mathbb{C} \text{ and } 0 < |z| < 1\} = \Delta - \{0\}.$$

For functions $f(z) \in \Sigma_p$ given by Eqn. (1) and $g \in \Sigma_p$ given by

$$g(z) = z^{-p} + \sum_{k=0}^{\infty} b_k z^{k-p+1} \quad (p \in \mathbb{N} := \{1,2,3,\dots\}).$$

we define the Hadamard product (or convolution) of f and g by

$$(f * g)(z) = z^{-p} + \sum_{k=0}^{\infty} a_k b_k z^{k-p+1} = (g * f)(z). \tag{2}$$

In terms of the Hadamard product (or convolution) of two functions, we define an analogue of the family Ruscheweyh derivative [3] by

$$D^{\lambda+p-1}f(z) = \frac{1}{z^p(1-z)^{\lambda+p}} * f(z), \quad (\lambda > -p, p \in \mathbb{N}, f \in \Sigma_p) \tag{3}$$

***Corresponding author: *B. Srutha Keerthi², *E-mail: sruthilaya06@yahoo.co.in**

or equivalently

$$D^{\lambda+p-1}f(z) = \frac{1}{z^p} \left(\frac{z^{\lambda+2p-1}f(z)}{(\lambda+p-1)!} \right)^{\lambda+p-1}, \quad (\lambda > -p, p \in \mathbb{N}, f \in \Sigma_p) \quad (4)$$

It follows readily from Eqn. (1), Eqn. (3), Eqn. (4) that

$$D^{\lambda+p-1}f(z) = z^{-p} + \sum_{k=0}^{\infty} \delta(\lambda, k) a_k z^{k-p+1}, \quad (\lambda > -p, p \in \mathbb{N}, f \in \Sigma_p) \quad (5)$$

where $f \in \Sigma_p$ is given by Eqn. (1) and

$$\delta(\lambda, k) = \binom{\lambda+p+k}{k+1} = \binom{\lambda+p+k}{\lambda+p-1}.$$

The function $f(z)$ is subordinate to the function $g(z)$, written $f \prec g$, provided there is an analytic function $w(z)$ defined on Δ with $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = g(w(z))$.

Definition: 1.1 Let $\phi(z)$ be an analytic function with positive real part in the unit disk Δ with $\phi(0) = 1$ and $\phi'(0) > 0$ that maps Δ onto a region starlike with respect to '1' and symmetric with respect to real axis. A function $f \in \Sigma_p$ is in the class $\Sigma_{p,\alpha}^*(\phi)$ if

$$-\frac{z^{1-\alpha}f'(z)}{p[f(z)]^{1+\alpha}} \prec \phi(z), \quad (z \in \Delta, \alpha > -\frac{1}{p}, p \in \mathbb{N}).$$

In the case of $\alpha = 0$ let us see the basic class as follows.

Definition: 1.2 Let $\phi(z)$ be an analytic function with positive real part in the unit disk Δ with $\phi(0) = 1$ and $\phi'(0) > 0$ that maps Δ onto a region starlike with respect to '1' and symmetric with respect to real axis. A function $f \in \Sigma_p$ is in the class $\Sigma_p^*(\phi)$ if

$$-\frac{zf'(z)}{pf(z)} \prec \phi(z), \quad (z \in \Delta, p \in \mathbb{N}).$$

The classes $\Sigma_{p,\alpha}^*(\phi)$, $\Sigma_p^*(\phi)$ were studied by Srutha Keerthi et al. [7].

Lemma: 1.1 If $p(z) = 1 + c_1z + c_2z^2 + \dots$ is a function with positive real part, then, for any complex number μ

$$|c_2 - \mu c_1^2| \leq 2 \max\{1, |2\mu - 1|\}$$

and the result is sharp for the functions given by

$$p(z) = \frac{1+z^2}{1-z^2}, \quad p(z) = \frac{1+z}{1-z}.$$

Lemma: 1.2 If $p_1(z) = 1 + c_1z + c_2z^2 + \dots$ is an analytic function with positive real part in Δ , then for r_1, r_2 real,

$$|c_3 + r_1c_1c_2 + r_2c_1^3| \leq 2H(s_1, s_2),$$

where $s_1 = 2(r_1 + 1)$, $s_2 = 2r_1 + 4r_2 + 1$ and

$$H(s_1, s_2) = \begin{cases} 1 & \text{for } (s_1, s_2) \in D_1 \cup D_2 \\ |s_2| & \text{for } (s_1, s_2) \in \cup_{k=3}^7 D_k \\ \frac{2}{3}(|s_1| + 1) \left(\frac{|s_1| + 1}{3|s_1| + s_2} \right)^{\frac{1}{2}} & \text{for } (s_1, s_2) \in D_8 \cup D_9 \\ \frac{1}{3} s_2 \left(\frac{s_1^2 - 4}{s_1^2 - 4s_2} \right) \left(\frac{s_1^2 - 4}{3(s_2 - 1)} \right)^{\frac{1}{2}} & \text{for } (s_1, s_2) \in D_{10} \cup D_{11} - \{\pm 2, 1\} \\ \frac{2}{3}(|s_1| - 1) \left(\frac{|s_1| - 1}{3(|s_1| - 1 - s_2)} \right)^{\frac{1}{2}} & \text{for } (s_1, s_2) \in D_{12} \end{cases}$$

the extremal functions up to the rotations are of the form

$$\begin{aligned}
 p_1(z) &= \frac{1+z^3}{1-z^3}, & p_1(z) &= \frac{1+z}{1-z}, \\
 p_1(z) = p_{1,0}(z) &= \frac{1+(1-2\lambda)(\varepsilon_1 - \varepsilon_2)z - \varepsilon_1\varepsilon_2z^2}{1-(\varepsilon_1 + \varepsilon_2)z + \varepsilon_1\varepsilon_2z^2}, \\
 p_1(z) = p_{1,1}(z) &= \frac{1-z^2}{1-2t_1z+z^2}, \\
 p_1(z) = p_{1,1}(z) &= \frac{1+2t_1z+z^2}{1-z^2}, \\
 \varepsilon_1 &= t_0 - e^{\frac{i\theta_0}{2}}(a \pm b), \\
 \varepsilon_2 &= -e^{-\frac{i\theta_0}{2}}(ia \pm b), \\
 a &= t_0 \cos\left(\frac{\theta_0}{2}\right), & b &= \sqrt{1-t_0^2 \sin^2\left(\frac{\theta_0}{2}\right)}, & \lambda &= \frac{b \pm a}{2b}, \\
 t_0 &= \left(\frac{2s_2(s_1^2+2)-3s_1^2}{3(s_2-1)(s_1^2-4s_2)}\right)^{\frac{1}{2}}, \\
 t_1 &= \left(\frac{|s_1|+1}{3(|s_1|+1+s_2)}\right)^{\frac{1}{2}}, \\
 t_2 &= \left(\frac{|s_1|-1}{3(|s_1|-1-s_2)}\right)^{\frac{1}{2}}, \\
 \cos\left(\frac{\theta_0}{2}\right) &= \frac{s_1}{2} \left[\frac{s_2(s_1^2+8)-2(s_1^2+2)}{2s_2(s_1^2+2)-3s_1^2} \right].
 \end{aligned}$$

The sets D_k , $k = 1, 2, \dots, 12$ are defined as follows.

$$D_1 = \left\{ (s_1, s_2) : |s_1| \leq \frac{1}{2}, |s_2| \leq 1 \right\},$$

$$D_2 = \left\{ (s_1, s_2) : \frac{1}{2}|s_1| \leq 2, \frac{4}{27}(|s_1|+1)^3 - (|s_1|+1) \leq |s_2| \leq 1 \right\},$$

$$D_3 = \left\{ (s_1, s_2) : |s_1| \leq \frac{1}{2}, |s_2| \leq -1 \right\},$$

$$D_4 = \left\{ (s_1, s_2) : |s_1| \geq \frac{1}{2}, |s_2| \leq -\frac{2}{3}(|s_1|+1) \right\},$$

$$D_5 = \left\{ (s_1, s_2) : |s_1| \leq 2, |s_2| \geq 1 \right\},$$

$$D_6 = \left\{ (s_1, s_2) : 2 \leq |s_1| \leq 4, |s_2| \geq \frac{1}{12}(s_1^2+8) \right\},$$

$$D_7 = \left\{ (s_1, s_2) : |s_1| \geq 4, |s_2| \geq \frac{2}{3}(|s_1|-1) \right\},$$

$$D_8 = \left\{ (s_1, s_2) : \frac{1}{2} \leq |s_1| \leq 2, -\frac{2}{3}(|s_1|+1) \leq |s_2| \leq \frac{4}{27}(|s_1|+1)^3 - (|s_1|+1) \right\},$$

$$D_9 = \left\{ (s_1, s_2) : |s_1| \geq 2, -\frac{2}{3}(|s_1|+1) \leq |s_2| \leq \frac{2|s_1|(|s_1|+1)}{s_1^2 + 2|s_1| + 4} \right\},$$

$$D_{10} = \left\{ (s_1, s_2) : 2 \leq |s_1| \leq 4, \frac{2|s_1|(|s_1|+1)}{s_1^2 + 2|s_1| + 4} \leq |s_2| \leq \frac{1}{12}(s_1^2 + 8) \right\},$$

$$D_{11} = \left\{ (s_1, s_2) : |s_1| \geq 4, \frac{2|s_1|(|s_1|+1)}{s_1^2 + 2|s_1| + 4} \leq |s_2| \leq \frac{2|s_1|(|s_1|-1)}{s_1^2 - 2|s_1| + 4} \right\},$$

$$D_{12} = \left\{ (s_1, s_2) : |s_1| \geq 4, \frac{2|s_1|(|s_1|-1)}{s_1^2 - 2|s_1| + 4} \leq |s_2| \leq \frac{2}{3}(|s_1|-1) \right\}.$$

Lemma 1.2 is a reformulation of a Lemma given by Prokhorov and Szyal [6]. For earlier works refer also [1, 2, 5].

2. COEFFICIENT BOUNDS:

Theorem: 2.1 Let $\phi(z) = 1 + B_1z + B_2z^2 + \dots$. If $f(z)$ given by (1) belongs to $\Sigma_{p,\alpha}^*(\phi)$, then for any complex number μ ,

$$|a_1 - \mu a_0^2| \leq \frac{B_1 p}{(2 + \alpha p)} \max \left\{ 1, \left| \frac{B_2}{B_1} - \left(\frac{(2 + \alpha p)}{2(1 + \alpha p)^2} (1 + \alpha - 2\mu) \right) B_1 p \right| \right\}. \quad (6)$$

This result is sharp. Further,

$$|a_2| \leq \frac{B_1 p}{2(3 + \alpha p)} H(s_1, s_2), \quad (7)$$

where s_1, s_2 and $H(s_1, s_2)$ is as defined in lemma 1.2, also

$$r_1 = \frac{B_2}{B_1} - \frac{(1 + \alpha)(3 + \alpha p) B_1 p}{2(2 + \alpha p)(1 + \alpha p)} - 1, \quad (8)$$

$$r_2 = \frac{B_3}{4B_1} - \frac{B_2}{2B_1} + \frac{(1 + \alpha)(3 + \alpha p)}{8(2 + \alpha p)(1 + \alpha p)} (2B_1 - 2B_2) p + \frac{(1 + \alpha)(2 + \alpha)(1 - \alpha p) B_1^2 p^2}{8(1 + \alpha p)^3} + \frac{1}{4}. \quad (9)$$

Proof: If $f(z) \in \Sigma_{p,\alpha}^*(\phi)$ let

$$p(z) = \frac{-z^{1-\alpha p} f'(z)}{p[f(z)]^{1+\alpha}} = 1 + b_1 z + b_2 z^2 + \dots \quad (10)$$

from (10)

$$a_0 = \frac{-b_1 p}{1 + \alpha p} \quad (11)$$

$$a_1 = -\frac{b_2 p}{(2 + \alpha p)} + \frac{b_1^2 (1 + \alpha) p^2}{2(1 + \alpha p)^2} \quad (12)$$

$$a_2 = b_3 - 3b_2 b_1 \left[\frac{(1 + \alpha) p}{(1 + \alpha p)(2 + \alpha p)} \right] + b_1^3 \left[\frac{(1 + \alpha)(1 + 2\alpha) p^2}{2(1 + \alpha p)^2} - \frac{2\alpha(1 + \alpha)(1 + 2\alpha) p^3}{6(1 + \alpha p)^3} \right] \quad (13)$$

Since $\phi(z)$ is univalent and $p(z) \prec \phi(z)$, then the function

$$p(z) = \frac{1 + \phi^{-1}(p(z))}{1 - \phi^{-1}(p(z))} = 1 + c_1 z + c_2 z^2 + \dots, \quad (14)$$

is analytic and has positive real part in Δ . Also we have

$$p(z) = \phi \left(\frac{p_1(z) - 1}{p_1(z) + 1} \right) \quad (15)$$

and from (15), we obtain

$$b_1 = \frac{B_1 c_1}{2} \quad (16)$$

$$b_2 = \frac{1}{2} \left[B_1 \left(c_2 - \frac{c_1^2}{2} \right) \right] + \frac{1}{4} B_2 c_1^2, \quad (17)$$

$$b_3 = \frac{B_1 c_3}{2} + (B_2 - B_1) \frac{c_2 c_1}{2} + (B_1 + B_3 - 2B_2) \frac{c_1^3}{8} \quad (18)$$

Substituting (16), (17), (18) in (11), (12), (13) we get

$$a_0 = \frac{-(B_1 c_1) p}{2(1 + \alpha p)}, \quad (19)$$

$$a_1 = \frac{B_1^2 c_1^2 (1 + \alpha) p^2}{8(1 + \alpha p)^2} + \frac{[B_1 c_1^2 - B_2 c_1^2 - 2B_1 c_2] p}{4(2 + \alpha p)} \quad (20)$$

$$a_2 = \frac{-B_1 p}{2(3 + \alpha p)} \left\{ c_3 + \left[\frac{B_2}{B_1} - \frac{(1 + \alpha)(3 + \alpha p) B_1 p}{2(2 + \alpha p)(1 + \alpha p)} - 1 \right] c_1 c_2 \right. \\ \left. + c_1^3 \left[\frac{(1 + \alpha)(1 + 2\alpha)(1 - \alpha p) B_1^2 p^2}{8(1 + \alpha p)^3} + \frac{2(1 + \alpha)(3 + \alpha p)(B_1 - B_2) p}{8(1 + \alpha p)(2 + \alpha p)} + \frac{B_3}{4B_1} - \frac{B_2}{2B_1} + \frac{1}{4} \right] \right\} \quad (21)$$

Therefore we have

$$a_1 - \mu a_0^2 = \frac{-B_1 p}{2(2 + \alpha p)} (c_2 - v c_1^2) \quad (22)$$

where

$$v = \frac{1}{2} \left[1 - \frac{B_2}{B_1} + \frac{B_1 p (2 + \alpha p) (1 + \alpha - 2\mu)}{2(1 + \alpha p)^2} \right]$$

Now we obtain inequality (6) by applying Lemma 1.1 in (22) and inequality (7) by applying Lemma 1.2 in (21). The result (6) is sharp for the function defined by

$$\frac{-z^{1-\alpha p} f'(z)}{p[f(z)]^{1+\alpha}} = \phi(z^2)$$

and

$$\frac{-z^{1-\alpha p} f'(z)}{p[f(z)]^{1+\alpha}} = \phi(z).$$

Corollary: 2.1 Let $\phi(z) = 1 + B_1 z + B_2 z^2 + \dots$. If $f(z)$ given by (1) belongs to $\Sigma_p^*(\phi)$, then for any complex number μ ,

$$|a_1 - \mu a_0^2| \leq \frac{B_1 p}{2} \max \left\{ 1, \left| \frac{B_2}{B_1} - (1 - 2\mu) B_1 p \right| \right\}. \quad (23)$$

Further,

$$|a_2| \leq \frac{B_1 p}{2} H(s_1, s_2) \quad (26)$$

where s_1, s_2 and $H(s_1, s_2)$ is as defined in lemma 1.2, also

$$r_1 = \frac{B_2}{B_1} - \frac{3B_1p}{4} - 1, \quad (25)$$

$$r_2 = \frac{B_3}{4B_1} - \frac{B_2}{2B_1} + \frac{3(B_1 - B_2)p}{8} + \frac{B_1^2p^2}{8} + \frac{1}{4}. \quad (26)$$

The result (23) is sharp for the function defined by

$$-\frac{zf'(z)}{pf(z)} = \phi(z^2)$$

and

$$-\frac{zf'(z)}{pf(z)} = \phi(z).$$

These are obtained by taking $\alpha = 0$ in equations (6) and (7).

3. APPLICATIONS TO FUNCTIONS DEFINED THROUGH RUSCHEWEYH DERIVATIVES:

The following classes $\Sigma_{p,\alpha}^\lambda(\phi)$ and $\Sigma_p^\lambda(\phi)$ were defined by Srutha Keerthi et al. [7].

Definition: 3.1 Let $\phi(z)$ be an analytic function with positive real part in the unit disk Δ with $\phi(0) = 1$, and $\phi'(0) > 0$ that maps Δ onto a region starlike with respect to 1 and symmetric with respect to the real axis. A function $f(z) \in \Sigma_p$ is in the class $\Sigma_{p,\alpha}^\lambda(\phi)$ if

$$\frac{-z^{1-\alpha p} [D^\lambda f(z)]'}{p[D^\lambda f(z)]^{1+\alpha}} < \phi(z), \quad (z \in \Delta, \alpha > -\frac{1}{p}, \lambda > -p, p \in \mathbb{N}).$$

By taking $\alpha = 0$, we set the class $\Sigma_p^\lambda(\phi)$.

Definition: 3.2 Let $\phi(z)$ be an analytic function with positive real part in the unit disk Δ with $\phi(0) = 1$ and $\phi'(0) > 0$ that maps Δ onto a region starlike with respect to 1 and symmetric with respect to the real axis. A function $f(z) \in \Sigma_p$ is in the class $\Sigma_p^\lambda(\phi)$ if

$$\frac{-z[D^\lambda f(z)]'}{p[D^\lambda f(z)]} < \phi(z), \quad (z \in \Delta, \lambda > -p, p \in \mathbb{N}).$$

Theorem: 3.1 Let the function $\phi(z) = 1 + B_1z + B_2z^2 + \dots$. If $f(z)$ given by (1) belongs to $\Sigma_{p,\alpha}^\lambda(\phi)$ then

$$|a_1 - \mu a_0^2| = \frac{2B_1p}{(\lambda + p + 1)(\lambda + p)(2 + \alpha p)} \max \left\{ 1, \left| \frac{B_2}{B_1} - \left[\frac{(2 + \alpha p)}{2(1 + \alpha p)^2} \left(1 + \alpha - \frac{(\lambda + p + 1)\mu}{(\lambda + p)} \right) \right] B_1p \right| \right\}. \quad (27)$$

This result is sharp.

Further

$$|a_2| \leq \frac{3B_1p}{(\lambda + p + 2)(\lambda + p + 1)(\lambda + p)(3 + \alpha p)} H(s_1, s_2),$$

where s_1, s_2 and $H(s_1, s_2)$ is as defined in lemma 1.2, also r_1, r_2 are given by equations (8), (9). The proof is similar to theorem 2.1, so the details are omitted.

Corollary: 3.1 Let the function $\phi(z) = 1 + B_1z + B_2z^2 + \dots$. If $f(z)$ given by (1) belongs to $\Sigma_p^\lambda(\phi)$ then

$$|a_1 - \mu a_0^2| = \frac{B_1p}{(\lambda + p + 1)(\lambda + p)} \max \left\{ 1, \left| \frac{B_2}{B_1} - \left[1 - \frac{(\lambda + p + 1)\mu}{(\lambda + p)} \right] B_1p \right| \right\}.$$

This result is sharp.

Further

$$|a_2| \leq \frac{B_1 p}{(\lambda + p + 2)(\lambda + p + 1)(\lambda + p)} H(s_1, s_2),$$

where s_1, s_2 and $H(s_1, s_2)$ is as defined in lemma 1.2, also r_1, r_2 are given by equations (25), (26).

Corollary 3.1 is obtained by taking $\alpha = 0$ in theorem 3.1.

REFERENCES:

- [1] Cho. N.E and Owa. S, (2003), On the Fekete-Szegő problem for strongly α -quasi convex functions, Tamkang J. Math., 34(1), 21–28.
- [2] Janteng. A and Darus. M, (2000), Coefficient problems for certain classes of analytic functions, J. Inst. Math. Comput. Sci. Math. Ser., 13(1), 91–96.
- [3] Joshi. S.B and Srivastava. H.M, (1999), A certain family of meromorphically multivalent functions, *Comp. Math. Appl.*, 38 (3–4), 201–211.
- [4] F.R. Keogh and E.P. Merkes, A coefficient inequality for certain classes of analytic functions, *Proc. Amer. Math. Soc.*, 20 (1969), 8–12.
- [5] Ma. W.C and Minda. D, (1992), *A unified treatment of some special classes of univalent functions*, Proceedings of the Conference on Complex Analysis, (Tianjin), 157–169, Conf. Proc. Lecture Notes, Anal., I, Int. Press, Cambridge, MA, 1994.
- [6] D.V. Prokhorov and J. Szynal, Inverse coefficients for (α, β) -qconvex functions, *Ann. Univ. Mariae Curie - Skłodowska Sect. A* 35 (1981), 125–143 (1984).
- [7] B. Srutha Keerthi, B. Adolf Stephen, A. Gangadharan and S. Sivasubramanian, (2008) A coefficient bounds for meromorphic p-valent functions, *Far East J. Math. Sci.*, 28(1), 161-172.
