

ON THE “ $\theta$ ” PARTIAL ORDERING OF s-UNITARY MATRICES

<sup>1</sup>S. Krishnamoorthy and <sup>2</sup>A. Govindarasu\*

<sup>1</sup>Professor and Head, Department of Mathematics, Ramanujan Research Centre, Government Arts College (Autonomous), Kumbakonam – 612 001, Tamil Nadu, INDIA

<sup>2</sup>Associate Professor in Mathematics, A. V. C. College (Autonomous), Mannampandal – 609 305, Tamil Nadu, INDIA

\*E-mail: [agavc@rediffmail.com](mailto:agavc@rediffmail.com)

(Received on: 24-11-11; Accepted on: 13-12-11)

## ABSTRACT

The concept of ‘ $\theta$ ’ partial ordering on s-unitary matrices is introduced. Some results relating to ‘ $\theta$ ’ partial ordering on s-unitary matrices are given.

**Mathematics Subject Classification:** 15A45, 15B57, 15A09.

**Keywords:** s-unitary matrix, Partial ordering, Lowener partial ordering, Star partial ordering, ‘ $\theta$ ’ partial ordering.

## 1. INTRODUCTION:

Some characterizations of the star partial ordering and rank subtractivity for matrices was discussed by R. E. Hartwig and G. P. H. Styan in [3]. Jurgen Grob observed some remarks on partial ordering of Hermitian Matrices in[5]. Xifu Liu and Hu yang discussed some results on the partial ordering of block matrices[7], In [4], Jorma K. Merikoski and Xiaogi Liu have developed star partial ordering on Normal Matrices. In this paper we introduced the concept of ‘ $\theta$ ’partial ordering on s-unitary matrices.

## 1.1. PRELIMINARIES AND NOTATIONS:

Let  $C_{n \times n}$  be the space of  $n \times n$  complex matrices of order  $n$ . For  $A \in C_{n \times n}$ ,

let  $A^T$ ,  $\bar{A}$ ,  $A^*$ ,  $A^s$ ,  $\bar{A}^s$  ( $= A^\theta$ ) denote transpose, conjugate, conjugate transpose, secondary transpose, conjugate secondary transpose of a matrix  $A$  respectively. Anna Lee[1] has initiated the study of secondary symmetric matrices. Also she has shown that for a complex matrix  $A$ , the usual transpose  $A^T$  and secondary transpose  $A^s$  are related as  $A^s = V A^T V$  where ‘V’ is the associated permutation matrix whose elements on the secondary diagonal are 1 and other elements are zero. Also  $\bar{A}^s$  denotes the conjugate secondary transpose of  $A$ .

i.e  $\bar{A}^s = (c_{ij})$  where  $c_{ij} = \overline{a_{n-j+1, n-i+1}}$  [2] and  $\bar{A}^s = V A^* V = A^\theta$ . Also ‘V’ satisfies the following properties.  $V^T = \bar{V} = V^* = V^\theta = V$  and  $V^2 = I$ .

A matrix  $A \in C_{n \times n}$  is said to be s-hermitian if  $A = A^\theta$  where  $A^\theta = \bar{A}^s$  or  $A^\theta = V A^* V$

A matrix  $A \in C_{n \times n}$  is said to be s-unitary if  $A A^\theta = A^\theta A = I$  or  $A^\theta = A^{-1}$  [6].

i.e.  $A V A^* V = V A^* V A = I$       i.e.  $V A^* V = A^{-1}$ .

In other words  $V A A^* = A^* V A = V$ .

**\*Corresponding author:** <sup>2</sup>A. Govindarasu\*, \*E-mail: [agavc@rediffmail.com](mailto:agavc@rediffmail.com)

## 2. MATRIX PARTIAL ORDERING ON MATRICES:

**Definition: 2.1** [Lowener Partial Ordering]

$$\text{For } A, B \in C_{n \times n} \quad A \underset{\text{L}}{\leq} B \text{ iff } B - A \geq 0$$

**Definition: 2.2** [Star Partial Ordering]

$$\text{For } A, B \in C_{n \times n}, A \underset{*}{\leq} B \text{ iff } A^*A = A^*B \text{ and } AA^* = BA^*$$

**Definition: 2.3** [Minus Partial Ordering]

$$\text{For } A, B \in C_{n \times n}, A \underset{\text{RS}}{\leq} B \text{ iff } \text{rank}(B - A) = \text{rank}B - \text{rank}A$$

**Remark: 2.4** If  $A \underset{\text{L}}{\leq} B$  for matrices A and B then  $B - A$  is s- hermitian.

## 3. “θ” PARTIAL ORDERING OF S-UNITARY MATRICES:

**Definition: 3.1** [‘θ’ Partial Ordering]

$$\text{For } A, B \in C_{n \times n}, A \underset{\theta}{\leq} B \text{ iff } A^\theta A = A^\theta B \text{ and } AA^\theta = BA^\theta$$

**Theorem: 3.2** Let A and B be s-hermitian matrices and nonnegative definite. Then  $A^2 \underset{\theta}{\leq} B^2$

$$\text{iff } A \underset{\theta}{\leq} B$$

**Proof:** Assuming  $A^2 = C, B^2 = D$

$$C \underset{\theta}{\leq} D \Rightarrow C^\theta C = C^\theta D$$

$$(A^2)^\theta A^2 = (A^2)^\theta B^2$$

$$(A^\theta)^2 A^2 = (A^\theta)^2 B^2$$

$$A^\theta A A = A^\theta B B$$

$$A^\theta A A = A^\theta B A \Rightarrow A^\theta A = A^\theta B$$

$$\text{Therefore } A \underset{\theta}{\leq} B$$

$$\text{Conversely, Assume, } A \underset{\theta}{\leq} B \Rightarrow A^\theta A = A^\theta B$$

Premultiplying bothsides by A

$$A^\theta A^2 = A^\theta B A$$

$$A^\theta A^2 = A^\theta A B \quad (\text{since } B A = A B)$$

$$A^\theta A^2 = A^\theta B A \quad (\text{since } A^\theta A = A^\theta B)$$

$$A^\theta A = A^\theta B$$

Premultiplying bothsides by  $A^\theta$

$$(A^\theta)^2 A^2 = (A^\theta)^2 B^2 \Rightarrow (A^2)^\theta A^2 = (A^2)^\theta B^2$$

$$\text{Therefore } A^2 \underset{\theta}{\leq} B^2$$

**Theorem: 3.3** Let  $AV \underset{\theta}{\leq} VA$ . If A is s-unitary then A is unitary.

$$\text{Proof: } AV \underset{\theta}{\leq} VA$$

$$\Rightarrow (AV)^\theta AV = (AV)^\theta VA$$

$$VA^\theta AV = VA^\theta VA$$

$$VIV = VA^\theta VA$$

$$I = VA^\theta VA$$

$$A^{-1} = V A^{\theta} V$$

$$\begin{aligned} A^{-1} &= V V A^{\theta} V V \\ A^{-1} &= A^{\theta} \end{aligned}$$

Therefore  $A A^{\theta} = I$

$$A V \xrightarrow{\theta} V A \Rightarrow A V (A V)^{\theta} = V A (A V)^{\theta}$$

$$A V V A^{\theta} = V A V A^{\theta}$$

$$I = V A V A^{\theta} V$$

$$V I V = A V^2 A^{\theta}$$

Therefore  $I = A A^{\theta}$

From (3.1) and (3.2)  $A A^{\theta} = A^{\theta} A = I$

Therefore  $A$  is unitary

**Theorem: 3.4** If  $A \xrightarrow[L]{\theta} B$  and  $A \xrightarrow{\theta} B$  then  $A = B$

**Proof:** Given,  $A \xrightarrow[\theta]{L} B \Rightarrow A^{\theta} A = A^{\theta} B$  and  $A A^{\theta} = B A^{\theta}$   
 $\Rightarrow A^{\theta} (B - A) = 0$

Taking secondary conjugate transpose on bothsides we get

$$(B - A)^{\theta} A = 0 \quad (3.3)$$

Since  $A \xrightarrow[L]{\theta} B$  we have  $(B - A)^{\theta} = V (B - A) V$

$$(3.3) \Rightarrow V (B - A)^{\theta} V A = 0$$

$$\Rightarrow V V (B - A) V V A = 0$$

$$\Rightarrow I (B - A) I A = 0$$

$$\Rightarrow (B - A) A = 0$$

$$\Rightarrow B A - A^2 = 0$$

$$\Rightarrow B A = A^2$$

$$\Rightarrow B = A$$

**Theorem: 3.5** If  $A$  and  $B$  are s-unitary matrices. Then  $A \xrightarrow[\theta]{L} B \Rightarrow A^{-1} \xrightarrow[\theta]{L} B^{-1}$

**Proof:** Given,  $A \xrightarrow[\theta]{L} B \Rightarrow A^{\theta} A = A^{\theta} B$  and  $A A^{\theta} = B A^{\theta}$

$$A^{\theta} A = A^{\theta} B$$

$$A^{-1} A = A^{-1} B$$

Taking secondary conjugate transpose on bothsides we get

$$(A^{-1} A)^{\theta} = (A^{-1} B)^{\theta}$$

$$A^{\theta} (A^{-1})^{\theta} = B^{\theta} (A^{-1})^{\theta}$$

$$A^{-1} (A^{-1})^{\theta} = B^{-1} (A^{-1})^{\theta}$$

$$A A^{\theta} = B A^{\theta} \Rightarrow A A^{-1} = B A^{-1} \quad [\text{Since } A \text{ and } B \text{ are s-unitary matrices}] \quad (3.4)$$

Taking secondary conjugate transpose on bothsides we get

$$(A A^{-1})^{\theta} = (B A^{-1})^{\theta}$$

$$(A^{-1})^{\theta} A^{\theta} = (A^{-1})^{\theta} B^{\theta} \Rightarrow (A^{-1})^{\theta} A^{-1} = (A^{-1})^{\theta} B^{-1} \quad (3.5)$$

From (3.4) and (3.5)  $A^{-1} \xrightarrow[\theta]{L} B^{-1}$

$$A \xrightarrow[\theta]{L} B \Rightarrow A^{-1} \xrightarrow[\theta]{L} B^{-1}$$

**Theorem: 3.6** If  $A$  and  $B$  are s-unitary matrices, then  $A \xrightarrow[\theta]{L} B \Rightarrow V A \xrightarrow[\theta]{*} V B$

**Proof:** Given,  $A \xrightarrow[\theta]{} B \Rightarrow A^\theta A = A^\theta B$  and  $AA^\theta = BA^\theta$

$$A^\theta A = A^\theta B \Rightarrow VA^* VA = VA^* VB$$

$$(VA)^*(VA) = (VA)^*(VB)$$

$$(AV)^*(VA) = (AV)^*(VB) \quad \text{Since } (VA)^* = (AV)^*$$

$$(VA)^*(VA) = (VA)^*(VB) \quad (3.6)$$

$$\begin{aligned} AA^\theta &= BA^\theta \\ AVA^* V &= BVA^* V \\ VAVA^* VV &= VBVA^* VV \\ VAVA^* &= VBVA^* \\ (VA)(AV)^* &= VB(VA)^* \\ (VA)(VA)^* &= VB(VA)^* \end{aligned} \quad (3.7)$$

From (3.6) and (3.7) we get  $VA \xrightarrow[\theta]{} VB$

$$A \xrightarrow[\theta]{} B \Rightarrow VA \xrightarrow[*]{} VB$$

**Theorem: 3.7** If  $A$  and  $B$  are s-unitary matrices. Then  $A \xrightarrow[*]{} B \Rightarrow VA \xrightarrow[\theta]{} VB$

**Proof:** Given,  $A \xrightarrow[\theta]{} B \Rightarrow A^* A = A^* B$  and  $AA^* = BA^*$

$$\begin{aligned} A^* A &= A^* B \\ (VA^* V)VA &= (VA^* V)VB \\ (A^\theta)VA &= (A^\theta)VB \\ V(A^\theta)VA &= V(A^\theta)VB \\ (AV)^\theta (VA) &= (AV)^\theta (VB) \\ (VA)^\theta (VA) &= (VA)^\theta (VB) \end{aligned} \quad (3.8)$$

$$\begin{aligned} AA^* &= BA^* \\ AVVA^* &= BVVA^* \\ AV(VA^* V) &= BV(VA^* V) \\ AV(A^\theta) &= BV(A^\theta) \\ AV(A^\theta V) &= BV(A^\theta V) \\ (AV)(VA)^\theta &= (BV)(VA)^\theta \\ (VA)(VA)^\theta &= (VB)(VA)^\theta \end{aligned} \quad (3.9)$$

From (3.8) and (3.9) we get  $VA \xrightarrow[\theta]{} VB$

## REFERENCES:

- [1]. Anna Lee., 1976, Secondary symmetric, secondary skew symmetric, secondary orthogonal matrices, Period. Math. Hungary, Vol.7 63-70.
- [2].Anna Lee. 1976, On s -symmetric, s-skew symmetric and s-orthogonal matrices, Period Math. Hungary Vol 7, 71–76.
- [3].Hartwig. R. E and Styani. G.P.H., 1986, On some characterization of the star partial ordering for matrices and rank substractivity, Lin. Alg. Appl., Vol. 82,145-161.
- [4].Jorma.K. Merikoski and Xiaoji Liu., 2006 ,On the star partial ordering of normal matrices, Journal of inequalities in Pure and Applied Mathematics, Vol. 7, issue 7, Art 17.
- [5].Jurgen Grob., 1997 ,Some remarks on partial ordering on hermitian matrices, Linear and Multilinear algebra, 42, 53-60.
- [6] Krishnamoorthy.S and Govindarasu. A., 2010, On secondary unitary Matrices, International Journal of computational science and Mathematics. Vol. 2 number 3, 247-253.
- [7] Liu and yang., 2011 “Some results on the partial ordering of block matrices” Vol. 54 1-9

\*\*\*\*\*