



COMPOSITION OPERATOR AND WEIGHTED COMPOSITION OPERATOR ON WEIGHTED SPACES OF ANALYTIC FUNCTIONS

D. Senthil Kumar¹ and P. Chandra Kala^{2*}

^{1,2}Department of Mathematics, Govt. Arts College (Autonomous), Coimbatore-641018, India
E-mail address: senthilsenkumhari@gmail.com, kala.mathchandra@gmail.com*

(Received on: 04-11-11; Accepted on: 18-11-11)

ABSTRACT

In this paper, we characterized invertible composition operator on weighted spaces of analytic functions and obtained dynamical system induced by weighted composition operator on weighted space of vector-valued analytic functions.

2010 Mathematics Subject Classification Primary: 47B37, 47B38, 47B07, 46E40, 46E10; **Secondary** 47D03, 37B05, 32A10, 30H05.

Keywords and phrases: System of weights, weighted composition operator, dynamical systems, seminorm, and operator valued mapping.

1. INTRODUCTION AND PRELIMINARIES:

The theory of weighted composition operators on spaces of analytic functions are gaining importance as it includes two nice classes of operators and very less information is known about weighted composition operators as compared to the theory of composition operators which has been presented in three monographs (see Cowen and MacCluer [3], Shapiro [8] and Singh and Manhas [9]). In [5] Manhas has made an application of multiplication operators can be found in the theory of dynamical systems. This paper is a continuation of the paper [6]. For more details about weighted composition operator on weighted spaces of continuous and analytic function spaces [1], [2], [4], [5], [6] and [9]. We have organized this paper into three sections. The preliminaries required for proving the results in the remaining sections are presented in section 1. In section two we have characterized the invertible composition operator on weighted spaces of analytic functions. In the last section we endowed with important theorems and obtained dynamical system induced by weighted composition operator on weighted spaces of analytic functions.

Let G be an open connected subset of C^N ($N \geq 1$) and $H(G, E)$ be the space of all vector-valued analytic functions from G into the Banach space E . Let V be a set of non-negative upper semi continuous functions on G . Then V is directed upward if for every pair $u, v \in V$ and $\lambda u > 0$, there exists $v \in V$ such that $\lambda u \leq v$ (point wise on G) for $i = 1, 2$. If V is directed upward and for each $z \in G$, there exists $v \in V$ such that $v(z) > 0$. Then we call V as an arbitrary system of weights on G . If U and V are two arbitrary systems of weights on G such that for each $u \in U$, there exists $v \in V$ for which $u \leq v$, then we write $U \leq V$. If $U \leq V$ and $V \leq U$, then we write $U \cong V$. Let V be an arbitrary system of weights on G .

Then we define $HV_b(G, E) = \{f \in H(G, E) : vf(G) \text{ is bounded in } E \text{ for each } v \in V\}$ and

$HV_o(G, E) = \{f \in H(G, E) : vf \text{ vanishes at } \infty \text{ on } G \text{ for each } v \in V\}$.

For $v \in V$ and $f \in H(G, E)$, we define $\|f\|_{v,E} = \sup\{v(z)\|f(z)\| : z \in G\}$. Clearly, the family $\|\cdot\|_{v,E} : v \in V$ of semi norms defines a Hausdorff locally convex topology on each of these spaces $HV_b(G, E)$ and $HV_o(G, E)$.

With this topology on each of these spaces $HV_b(G, E)$ and $HV_o(G, E)$ are called weighted locally convex spaces

Corresponding author: P. Chandra Kala^{2}, *E-mail: kala.mathchandra@gmail.com

of vector-valued analytic functions. These spaces have a basis of closed absolutely convex neighborhoods of the form

$$B_{v,E} = \left\{ f \in HV_b(G, E) : \|f\|_{v,E} \leq 1 \right\}.$$

Definitions of reasonable and essential system of weights: [5]:

An arbitrary system of weights V is called reasonable if it satisfies the following properties: for each $v \in V$, there exists $\tilde{v} \in \tilde{V}$ such that for each $v \in V, \|f\|_v \leq 1$ iff $\|f\|_{\tilde{v}} \leq 1$ such that for every $f \in H(G)$; if $v \in V$, then for every $z \in G \exists f_z \in B_v$ such that $|f_z(z)| = \frac{1}{\tilde{v}(z)}$. A weight $v \in V$ is called essential if there exists a constant $c > 0$ such that $v(z) \leq \tilde{v}(z) \leq cv(z)$, for each $z \in G$.

A reasonable system of weights V is called an essential system if each $v \in V$ is an essential weight.

Let G be a topological group with e as identity, let X be a topological space and $\pi : G \times X \rightarrow X$ be a continuous map such that

- (i) $\pi(e, x) = x$ for every $x \in X$
- (ii) $\pi(st, x) = \pi(s, \pi(t, x))$ for every $t, s \in G, x \in X$.

Then the triple (G, X, π) is called a transformation group, X is a state space. If $G = (R, +)$ the corresponding transformation group is called a dynamical system. The transformation group $(\mathbb{R}, X, \pi)$ is known as continuous dynamical system. If X is a Banach space and

$$\pi(t, \alpha x + \beta y) = \alpha \pi(t, x) + \beta \pi(t, y) \text{ for } t \in \mathbb{R}, \alpha, \beta \in \mathbb{C}, x, y \in X$$

then $(\mathbb{R}^+, X, \pi)$ is called a linear dynamical system.

2. INVERTIBLE COMPOSITION OPERATOR ON ANALYTIC FUNCTION SPACES:

Theorem: 2.1 Let V be an arbitrary system of weights on G . Let $\phi : G \rightarrow G$ be analytic map such that C_ϕ is a composition operator on $HV_b(G)$. Then C_ϕ is invertible if (i) for each $v \in V$, there exist $u \in V$ such that $v(\phi(z)) \leq u(z)$ for every $z \in G$. (ii) for each $z \in G$, ϕ is a conformal mapping of G .

Proof: It is enough to show that C_ϕ is bounded below and onto. For let $v \in V$ and B_v be a neighborhood of the origin in $HV_b(G)$. Then by condition (i), there exist $u \in V$ such that $v(\phi(z)) \leq u(z)$ for every $z \in G$.

We claim that $C_\phi(B_v^c) \subseteq B_u^c$. Let $f \in B_v^c$.

$$\begin{aligned} \text{Then } 1 < \|f\|_v &= \sup \{ v(z) \|f(\phi(z))\| : z \in G \} \leq \sup \{ u(z) \|f(\phi(z))\| : z \in G \} \\ &\leq \|C_\phi f\|_u. \end{aligned}$$

This shows that $C_\phi f \in B_u^c$ and hence C_ϕ is bounded below. Now we shall show that C_ϕ is onto. For let $g \in HV_b(G)$. Fix $z_0 \in G$ and let $v \in V$ be such that $v(\phi(z_0)) > 0$. Then by condition (i), there exists $u \in V$ such that $v(\phi(z_0)) \leq u(z_0)$ for every $z_0 \in G$. By condition (ii) define an analytic map $\phi^{-1} : G \rightarrow G$. Also define $h : G \rightarrow \mathbb{C}$ as $h = g \circ \phi^{-1}$.

Then we show that $h \in HV_b(G)$. For, let $v \in V$. Then by condition (i), there exists $v \in V$ such that $v(\phi(z)) \leq u(z)$ for every $z \in G$. Now we have

$$\begin{aligned}\|h\|_v &= \sup\{v(\phi(z))\|h(\phi(z))\| : z \in G\} \\ &\leq \sup\{u(z)\|g \circ \phi^{-1}\| : z \in G\} \\ &= \sup\{v(z)\|g(z)\| : z \in G\} \leq \|g\|_{u,E} < \infty.\end{aligned}$$

This proves that $h \in HV_b(G)$ and clearly $C_\phi(h) = g$. Hence C_ϕ is invertible.

Corollary: 2.2 Let V be an arbitrary system of weights on G . Let $\phi: G \rightarrow G$ be analytic map. Then C_ϕ is invertible composition operator if ϕ is a conformal mapping of G onto itself such that $V \circ \phi \equiv V$.

Proof: Since, $\phi: G \rightarrow G$ is a conformal mapping such that $V \leq V \circ \phi$ [5]. Implies that C_ϕ is a composition operator on $HV_b(G)$.

Let $v \in V$. Then, since $V \circ \phi \leq V$ such that $u \in V$ such that $v(\phi(z)) \leq u(z)$ for every $z \in G$.

Further, there exists $w \in V$ such that $\frac{1}{mv} \leq w$ and $v(\phi(z)) \leq w(z)$ for every $z \in G$. Thus according to the given condition and theorem: 3.1, it follows that C_ϕ is invertible composition operator on $HV_b(G)$.

Corollary: 2.3 Let G be a simply connected open subset of $\mathbb{C}$ and let $V = \{\lambda_{\chi_k} : \lambda > 0, K \subseteq G, K \text{ is compact}\}$. Let $\phi: G \rightarrow G$ be analytic map. Then C_ϕ is invertible composition operator on $HV_b(G)$ iff ϕ is a conformal mapping of G onto itself.

Theorem: 2.4 Let G be a simply connected open subset of $\mathbb{C}$ and let V be a reasonable system of weights on G . Let $\phi: G \rightarrow G$ be analytic map such that C_ϕ is composition operator on $\widetilde{HV_b(G)}$. Then C_ϕ is invertible iff

- (i) ϕ is a conformal mapping of G onto itself.
- (ii) for each $v \in V$ there exists $u \in V \ni \tilde{v}(\phi(z)) \leq \tilde{u}(z)$ for every $z \in G$.

Proof: Suppose, conditions (i) and (ii) hold. Let $z \in G$ and let $v \in V$ be such that $v(\phi(z_0)) > 0$. Then $\tilde{v}(\phi(z_0)) > 0$ and by condition (ii), there exists $u \in V$ such that $\tilde{v}(\phi(z_0)) \leq \tilde{u}(z_0)$. Further by condition (i), we have that $\tilde{v}(z) \leq \tilde{u}(\phi^{-1}(z))$ for every $z \in G$.

(i.e) $\tilde{v}(z)\|(\phi^{-1}(z))\| \leq \tilde{u}(\phi^{-1}(z))$ for every $z \in G$.

Therefore $C_{\phi^{-1}}$ is a composition operator on $\widetilde{HV_b(G)}$ such that $C_\phi C_{\phi^{-1}} = C_{\phi^{-1}} C_\phi = I$, the identity operator.

Hence C_ϕ is invertible. Conversely, suppose that C_ϕ is invertible on $\widetilde{HV_b(G)}$. First to show that ϕ is a conformal mapping. We need to prove that ϕ is bijective. Let $z_1, z_2 \in G$ be such that $z_1 \neq z_2$. Then by Riemann mapping theorem, there exists one-one analytic function $g \in H^\infty(G)$ such that $g(z_1) \neq g(z_2)$.

Define $h: G \rightarrow \mathbb{C}$ as $h(z) = g(z) - g(z_1)$ for every $z \in G$. Let $g_{z_2} \in \widetilde{HV_b(G)}$ be such that $g_{z_2}(z_2) \neq 0$ and $h_0 = h_{g_{z_2}}$. Since $\widetilde{HV_b(G)}$ is module over $H^\infty(G)$, it follows that $h_0 \in \widetilde{HV_b(G)}$ such that $h_0(z_1) = 0$ and $h_0(z_2) \neq 0$. Let $\mathcal{E} = \mathcal{E}_1 + \mathcal{E}_2$ and let $v \in V$ be such that $v(z_1) \geq 1$ and $v(z_2) \geq 1$. There exists $\tilde{v} \in \tilde{V}$ such that $\tilde{v}(z_1) \geq 1$ and $\tilde{v}(z_2) \geq 1$.

Since $C_\phi(\widetilde{HV_b(G)})$ is dense in $\widetilde{HV_b(G)}$ there exists $F \in \widetilde{HV_b(G)}$ such that $\|C_\phi F\| < \frac{1}{2\mathcal{E}}$.

$$(i.e) \tilde{v}_{z_1} \|F(\phi(z_1))\| < \frac{1}{2\mathcal{E}} \text{ and } \tilde{v}_{z_2} \|F(\phi(z_2))\| - \frac{1}{\mathcal{E}_2} < \frac{1}{2\mathcal{E}}.$$

Further it implies that $\|F(\phi(z_1))\| < \frac{\mathcal{E}_1}{2\mathcal{E}} < \frac{1}{2}$ and $|\|F(\phi(z_2))\| - 1| < \frac{\mathcal{E}_2}{2\mathcal{E}} < \frac{1}{2}$.

From the last two inequalities, it is easy to conclude that $\phi(z_1) \neq \phi(z_2)$. Thus ϕ is injective and also using the injectivity argument of C_ϕ , we can conclude that ϕ is onto. Now to prove condition (ii), we fix $v \in V$. Then there exists $\tilde{v} \in \tilde{V}$. According to theorem:2.1, C_ϕ is bounded below on $\widetilde{HV_b(G)}$. Thus there exists $\tilde{u} \in \tilde{V}$ with $u \in V$. We claim that $\tilde{v}(\phi(z)) \leq \tilde{u}(z)$ for every $z \in G$.

Let $z_0 \in G$. For $z = \phi(z_0)$. We have $\tilde{v}(\phi(z_0))g_0(\phi(z_0)) \leq 1$, where $g_0 \in \widetilde{HV_b(G)}$.

That is $\tilde{v}(\phi(z_0)) \leq \tilde{u}(z_0)$. This proves our claim.

Corollary: 2.5 Let G be a simply connected open subset of $\mathbb{C}$ and let V be an essential system of weights on G .

Let $\phi: G \rightarrow G$ be analytic map such that C_ϕ is composition operator on $HV_b(G)$. Then C_ϕ is invertible iff

- (i) ϕ is a conformal mapping of G onto itself for every $z \in G$.
- (ii) for each $v \in V$ there exists $u \in V \ni v(\phi(z)) \leq u(z)$ for every $z \in G$.

Proof: Follows from the above theorem, since each $v \in V$ is essential.

3. DYNAMICAL SYSTEM AND WEIGHTED COMPOSITION OPERATOR:

Let $g \in H^\infty(G, B(E))$ and $\|g\|_\infty = \sup\{\|g(z)\| : z \in G\}$. Then for each $t \in \mathbb{R}$. We define $\psi_t: G \rightarrow B(E)$ as $\psi_t(z) = e^{tg(z)} \forall z \in G$ and let $\phi_t: \mathbb{R} \rightarrow \mathbb{R}$ be the analytic map defined by $\phi_t(\omega) = t + \omega \forall t, \omega \in \mathbb{R}$.

Theorem: 3.1 Let $\varphi: X \rightarrow X$ and $\psi_t: X \rightarrow X$ be continuous functions. Then $W_{\psi_t, \varphi} f$ is bounded $\forall t \in \mathbb{R}, f \in H(X)$.

Proof: We shall show that $W_{\psi_t, \varphi} f$ is continuous at the origin. We claim that $W_{\psi_t, \varphi}(B) \subseteq B$. We have

$$\begin{aligned} \|W_{\psi_t, \varphi} f\|_v &= \sup\{v(x) \|\psi_t(x) f(\varphi(x))\| : x \in X \text{ and } t \in \mathbb{R}\} \\ &\leq \sup\{v(x) \|\psi_t(x)\| \|f(\varphi(x))\| : x \in X \text{ and } t \in \mathbb{R}\} \\ &\leq \sup\{v(x) \|e^{tg(x)}\| \|f(x)\| : x \in X \text{ and } t \in \mathbb{R}\} \\ &\leq e^{|t|M} \sup\{v(x) \|f(x)\| : x \in X\} \\ &\leq \|f\|_v \leq 1 \text{ as } t \rightarrow 0. \end{aligned}$$

Therefore $W_{\psi_t, \varphi} f$ is continuous at the origin. Hence proved.

Theorem: 3.2 Let $H^\infty(G, B(E))$ be the space of bounded analytic functions. Let $h_\alpha(\varphi_{t_\alpha})$ converges to $h(\varphi_t)$ in $H^\infty(G, B(E))$ and let f_α be a sequence converging to f in $HV_b(G, E)$. Then the product of $f_\alpha h_\alpha(\varphi_{t_\alpha})$ converges to $fh(\varphi_t)$ in $HV_b(G, E)$.

Proof: Let $v \in V$.

Then

$$\begin{aligned} \|f_n h_n(\varphi_{t_n}) - fh(\varphi_t)\|_{v,E} &= \sup\{v(z) \|f_n(z)h_n(\varphi_{t_n}(z)) - f(z)h(\varphi_t(z))\| : z \in G\} \\ &= \sup\{v(z) \|f_n(z)h_n(t_n + z) - f(z)h(t + z)\| : z \in G\} \\ &\leq \sup\{v(z) \|f_n(z)h_n(t_n)h_n(z) - f_n(z)h(t)h(z)\| : z \in G\} \\ &\quad + \sup\{v(z) \|f_n(z)h(t)h(z) - f(z)h(t)h(z)\| : z \in G\} \\ &\quad + \sup\{v(z) \|f_n(z) - f(z)\| \|h(t)h(z)\| : z \in G\} \\ &\leq \sup\{v(z) \|f_n(z)\| \|h_n(t_n)h_n(z) - h(t)h(z)\| : z \in G\} \\ &\leq \|h_n(\varphi_{t_n}) - h(\varphi_t)\|_\infty \|f_n\|_v + \|h(\varphi_t)\|_\infty \|f_n - f\|_v \rightarrow 0 \end{aligned}$$

as $\|h_n(\varphi_{t_n}) - h(\varphi_t)\|_\infty \rightarrow 0$ and $\|f_n - f\|_v \rightarrow 0$.

Theorem: 3.3 Let V be an arbitrary system of weights on G and let $\pi : \mathbb{R} \times HV_b(G, E) \rightarrow H(G, E)$ be the function defined by $\pi(t, f) = W_{\psi_t, \varphi_t} f$ for every $t \in \mathbb{R}$ and $f \in HV_b(G, E)$. Then π is linear dynamical system on $HV_b(G, E)$.

Proof: Since, for every $t \in \mathbb{R}$, W_{ψ_t, φ_t} is a weighted composition operator on $HV_b(G, E)$. Thus it follows that $\pi(t, f) \in HV_b(G, E)$ for every $t \in \mathbb{R}$ and $f \in HV_b(G, E)$.

Clearly π is linear and

$$\begin{aligned} \pi(0, f)(z) &= W_{\psi_0, \varphi_0} f(z) \text{ for all } z \in G \\ &= f(z) \text{ for all } z \in G \\ &= \psi_0(z) f(z). \end{aligned}$$

Therefore $\pi(0, f) = f$.

Similarly $\pi(t + s, f) = \pi(t, \pi(s, f))$ for all $t, s \in \mathbb{R}$.

Next to show that π is dynamical system, it sufficient to show that π is jointly continuous [7]. Let the sequence $\{(t_\alpha, f_\alpha)\}$ be a net in $\mathbb{R} \times HV_b(G, E)$ such that $(t_\alpha, f_\alpha) \rightarrow (t, f)$.

Let $v \in V$. Then

$$\begin{aligned} \|\pi(t_\alpha, f_\alpha) - \pi(t, f)\|_{v,E} &= \|W_{\psi_{t_\alpha}, \varphi_{t_\alpha}} f_\alpha - W_{\psi_t, \varphi_t} f\|_{v,E} \\ &= \sup\{v(z) \|\psi_{t_\alpha}(z) f_\alpha(\varphi_{t_\alpha}(z)) - \psi_t(z) f(\varphi_t(z))\| : z \in G\} \\ &\leq \sup\{v(z) \|\psi_{t_\alpha}(z) f_\alpha(\varphi_{t_\alpha}(z)) - \psi_t(z) f_\alpha(\varphi_{t_\alpha}(z))\| : z \in G\} \end{aligned}$$

$$\begin{aligned}
 & + \sup \{v(z) \|\psi_t(z) f_\alpha(\varphi_{t_\alpha}(z)) - \psi_t(z) f(\varphi_t(z))\| : z \in G\} \\
 & \leq \sup \{v(z) \|\psi_{t_\alpha}(z) - \psi_t(z)\| \|f_\alpha(\varphi_{t_\alpha}(z))\| : z \in G\} \\
 & + \sup \{v(z) \|\psi_t(z)\| \|f_\alpha(\varphi_{t_\alpha}(z)) - f(\varphi_t(z))\| : z \in G\} \\
 & \leq \sup \{ \|e^{(t_\alpha-t)g(z)}\| v(z) \|f_\alpha(t_\alpha + z)\| : z \in G\} \\
 & + \sup \{ \|e^{tg(z)}\| v(z) \|f_\alpha(t_\alpha + z) - f(t + z)\| : z \in G\} \\
 & \leq e^{\|t\|g\|_\infty} (e^{\|t_\alpha-t\|g\|_\infty} - 1) v(z) \|f_\alpha(\varphi_{t_\alpha})\|_{v,E} \\
 & + e^{\|t\|g\|_\infty} \|f_\alpha(\varphi_{t_\alpha}) - f(\varphi_t)\|_{v,E} \rightarrow 0.
 \end{aligned}$$

Since by theorem: 3.2. This shows that π is jointly continuous and hence π is a linear dynamical system on $HV_b(G, E)$.

ACKNOWLEDGEMENT:

The work is supported by Rajiv Gandhi National Fellowship under the University Grants Commission Fellowship and a bursary awarded to P. Chandra kala.

REFERENCES:

- [1] K. D. Bierstedt, J. Bonet and A. Galbis, Weight spaces of holomorphic function on balanced domains, Michigan Math J. 40(1993), no.2, 271-297.
- [2] M.D. Contreras and A.G.Hernandez-Diaz, Weighted composition operator on weighted Banach spaces of analytic function, J. Aust. Math. Soc. Ser A69 (2000), no. 1, 41-60.
- [3] C. Cowen and B. D. MacCluer, Composition operator on spaces of analytic functions, Studies in Advanced Mathematics, CRC press, Boca Raton, Fla, USA, 1995.
- [4] J. S. Manhas, Weighted composition operator on weighted locally convex spaces of analytic functions, Southeast Asian Bull. Math. 29. (2005), no.1, 127-140.
- [5] J.S. Manhas, Composition operators and Multiplication operator on weighted spaces of analytic function, Research article, International Journal of Mathematics and Mathematical Sciences, vol. 2007.
- [6] J.S. Manhas, Weighted composition operator on weighted spaces of vector-valued analytic function, J. Korean Math.soc. 45(2008), no.5, 1203-1220.
- [7] Ruess. W.M and Summers. W.H, Minimal sets of almost periodic motions, Math. Ann., 276(1986), 145-158.
- [8] J. H. Shpiiro, Composition operators and classical function theory, Universitext: Tracts in Mathematics, Springer, New York, NY, USA, 1993.
- [9] R.K. Singh and J.S. Manhas, Composition operators on function spaces, North Holland Mathematics Studies, 1993.
