



ON SOLVING PELL MEANS

Lexter R. Natividad*

Faculty of College of Education, Central Luzon State University, Science City of Muñoz 3120, Nueva Ecija, Philippines

E-mail: lexter_natividad@yahoo.com

(Received on: 06-12-11; Accepted on: 23-12-11)

ABSTRACT

Pell sequence is associated with Fibonacci sequence. Many identities have been already derived for Pell sequence in books and mathematical journals. This paper will present formulas in solving Pell means as well as solving the sequence itself.

2010 Mathematics Subject Classifications: 11B50, 11B99.

Key Words and Phrases: Pell sequence, Pell-like sequence, Pell mean, missing terms.

INTRODUCTION:

Pell Sequence is one of many sequences which share basic properties and identities of Fibonacci sequence as well as with Fibonacci's associated sequence, Lucas sequence [1]. By allowing an extension to the Pell sequence with negative subscripts, it can be defined as

$$\{P_n\}: \quad \dots \quad P_{-4} \quad P_{-3} \quad P_{-2} \quad P_{-1} \quad P_0 \quad P_1 \quad P_2 \quad P_3 \quad P_4 \quad P_5 \quad \dots$$

$$\quad \quad \quad \dots \quad -12 \quad 5 \quad -2 \quad 1 \quad 0 \quad 1 \quad 2 \quad 5 \quad 12 \quad 29 \quad \dots$$

in which

$$P_0 = 0, \quad P_1 = 1, \quad P_{n+2} = 2P_{n+1} + P_n$$

and

$$P_{-n} = (-1)^{n+1}P_n,$$

as shown by Horadam in [2].

In mathematical statement, Pell sequence is a succession of numbers that are obtained by getting the sum of twice the previous number and the number before that. On the other hand, Fibonacci sequence (1, 1, 2, 3, 5, 8, 13, 21....) is a succession of numbers that are obtained through adding the two preceding numbers.

All integer sequences that satisfy Fibonacci-like sequence, $g(n+2) = g(n) + g(n+1)$, can be solved given two initial values using the equation

$$g(n) = F(n)g(1) + F(n-1)g(0),$$

where $F(n)$ and $F(n-1)$ are corresponding terms of Fibonacci sequence.

The n^{th} term of Pell sequence can be solved using the explicit formula

$$P_n = \frac{(1+\sqrt{2})^n - (1-\sqrt{2})^n}{2\sqrt{2}}.$$

***Corresponding author: Lexter R. Natividad*, *E-mail: lexter_natividad@yahoo.com**

Also, it can be solved using the binomial sums

$$P_n = \sum_{k=0}^{\lfloor n-2/2 \rfloor} \binom{n}{2k+1} 2^k.$$

A well known counterpart of Pell sequence is called as Pell-Lucas sequence where the n^{th} Pell-Lucas number can be solved by the binomial sum

$$Q_n = 2 \sum_{k=0}^{n/2} \binom{n}{2k} 2^k$$

or by the Binet-type formula

$$Q_n = (1 - \sqrt{2})^n + (1 + \sqrt{2})^n.$$

As seen from the formula above, terms in Pell Sequence are proportional to silver ratio $(1 + \sqrt{2})$ which is analogous to the terms in Fibonacci sequence that are proportional to golden ratio $(1 + \sqrt{5})$. Many Pell sequence identities have been presented by Horadam in [2] and the author will add some properties by studying Pell means. Pell means are terms between any two terms of Pell or Pell-like sequence.

Definition: 1.1 If $a, x_1, x_2, x_3, \dots, x_{n-1}, x_n, b$ is a Pell or Pell-like sequence, then $x_1, x_2, x_3, \dots, x_{n-1}, x_n$ are called Pell means between a and b .

Inserting arithmetic means, geometric means and harmonic means has been very well solved in many college and high school mathematics books. Recently, a formula for solving Fibonacci means based on the number of missing terms has been proposed and derived in [3]. This formula is derived to find the first missing term of any Fibonacci-like sequence with n missing terms in any arbitrary constant first term a and last term b . It has the form of

$$x_1 = \frac{b - F_n a}{F_{n+1}}.$$

Since it is known that Pell sequence is a derivative of Fibonacci sequence, it is not farfetched to think that solving Pell means is related to that formula.

This paper will present deriving formulas in solving Pell means.

THE FORMULA:

Deriving the formula for the first missing term of any Pell-like sequence could be found using a recognizable pattern. Just like in Fibonacci sequence, it can be noted that all the missing terms in a Pell or Pell-like sequence could be solved by finding the first missing term x_1 given arbitrary constant a and b ; where a is the first term given and b is the last term given in this sequence.

From the definition of Pell sequence, the general formula for x_1 will make the solving easy for the other missing terms. Like the approach in [3], x_1 can be solved for Pell or Pell-like sequence with

- a. One missing term, the sequence is a, x_1, b

$$a + 2x_1 = b$$

then

$$x_1 = \frac{b - a}{2}.$$

- b. Two missing terms, the sequence is a, x_1, x_2, b

$$a + 2x_1 = x_2$$

$$x_1 + 2x_2 = b$$

then

$$x_1 = \frac{b - 2a}{5}.$$

c. Three missing terms, the sequence is a, x_1, x_2, x_3, b

$$a + 2x_1 = x_2$$

$$x_1 + 2x_2 = x_3$$

$$x_2 + 2x_3 = b$$

then

$$x_1 = \frac{b - 5a}{12}$$

d. Four missing terms, the Pell like sequence is a, x_1, x_2, x_3, x_4, b

$$a + 2x_1 = x_2$$

$$x_1 + 2x_2 = x_3$$

$$x_2 + 2x_3 = x_4$$

$$x_3 + 2x_4 = b$$

then

$$x_1 = \frac{b - 12a}{29}$$

A clear observation could be seen in the formula for x_1 that the numerical coefficient of a in numerator and the denominator of the formulas were following the Pell sequence as shown in Table 1. The formula can be illustrated as

$$x_1 = \frac{b - P_n a}{P_{n+1}}.$$

Table- 1: Relationship of number of missing terms with numerator and denominator of formulas.

Number of Missing Term	Coefficient of a in Numerator	Coefficient of Denominator
1	1	2
2	2	5
3	5	12
4	12	29
⋮	⋮	⋮
⋮	⋮	⋮
⋮	⋮	⋮
N	$\frac{(1 + \sqrt{2})^n - (1 - \sqrt{2})^n}{2\sqrt{2}}$	$\frac{(1 + \sqrt{2})^{n+1} - (1 - \sqrt{2})^{n+1}}{2\sqrt{2}}$

Therefore for Pell mean, the formula for x_1 is

$$x_1 = \frac{b - \left[\frac{(1 + \sqrt{2})^n - (1 - \sqrt{2})^n}{2\sqrt{2}} \right] a}{\frac{(1 + \sqrt{2})^{n+1} - (1 - \sqrt{2})^{n+1}}{2\sqrt{2}}}$$

It can also be written as

$$x_1 = \frac{2b\sqrt{2} - \left[(1 + \sqrt{2})^n - (1 - \sqrt{2})^n \right] a}{(1 + \sqrt{2})^{n+1} - (1 - \sqrt{2})^{n+1}}$$

where x_1 is the first missing term in Pell like sequence
 a is the first term given
 b is the last term given
 n is the number of missing terms .

These formulas could now be used to find all the Pell means in the sequence.

ACKNOWLEDGMENT:

The author would like to thank Central Luzon State University Science High School Math Enthusiasts' Circle (CLSUSHS-MEC) for the support they gave to him and many thanks to Ms. Cynthia Lyn Pagaduan-Curada, for being his grammar editor for this paper.

REFERENCES:

- [1] Bicknell, M., A Primer on the Pell Sequence and Related Sequences, *The Fibonacci Quarterly* 13 (1975), #4 345-349.
- [2] Horadam, A.F., Pell Identities, *The Fibonacci Quarterly* 9 (1971), #3 245-252, 263.
- [3] Natividad, L.R., Deriving a formula in solving Fibonacci-like sequence, *International Journal of Mathematics and Scientific Computing* 1 (2011), #1 19-21.
