

FIXED POINT THEOREM OF MULTIVALUED MAPPINGS IN CONE METRIC SPACES

Saurabh Jain[§] and Dr. V. H. Badshah*

[§]Indore Institute of Science and Technology, Indore (M. P.), India

E-mail: sj5129@gmail.com

*School of Studies in Mathematics Vikram University Ujjain-456010 (M. P.), India

E-mail: vhbadshah@yahoo.co.in

(Received on: 19-10-11; Accepted on: 03-11-11)

ABSTRACT

Let P be a subset of a Banach space E and P is normal and regular cone on E , we prove the existence of the fixed point for multivalued maps in cone metric spaces and these theorems generalize the recent results of various authors.

Mathematics Subject Classification: Primary 47H10; Secondary 54H25.

Keywords: Cone metric space, Multivalued mappings, Fixed point.

1. INTRODUCTION AND PRELIMINARIES:

In recent years, several authors (see [1-6]) have studied the strong convergence to a fixed point with contractive constant in cone metric spaces. Seong Hoon Cho and Mi Sun Kim [6] have proved certain fixed point theorems by using Multivalued mapping in the setting of contractive constant in metric spaces. We first recall definitions and known results that are needed in the sequel. Let E be a Banach space and a subset P of E is said to be a cone if it satisfies the following conditions,

- (i) $P \neq \emptyset$ and P is closed
- (ii) $ax + by \in P$ for all $x, y \in P$ and a, b are non-negative real numbers;
- (iii) $P \cap (-P) = \emptyset$.

The partial ordering \leq with respect to the cone P by $x \leq y$ if and only if $y - x \in P$. If $y - x \in$ interior of P , then it is denoted by $x \ll y$. The cone P is said to be a Normal if a number $K > 0$ such that for all $x, y \in E, 0 \leq x \leq y$ implies $\|x\| \leq K \|y\|$. The cone P is called regular if every increasing sequence which is bounded above is convergent and every decreasing sequence which is bounded below is convergent.

Definition: 1.1 Let X be a non-empty set, and suppose the mapping $d : X \times X \rightarrow E$ is said to be a Cone metric space if it satisfies

- (i) $0 \leq d(x, y)$ for all $x, y \in X$ and $d(x, y) = 0$ if and only if $x = y$.
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$.
- (iii) $d(x, y) = d(x, z) + d(z, y)$ for all $x, y \in X$.

Example: 1.2 Let $E = \mathbb{R}^2, P = \{(x, y) \in E : x, y \geq 0\}, X = \mathbb{R}$ and $d : X \times X \rightarrow E$ defined by

$$d(x, y) = (|x - y|, \alpha|x - y|)$$

where $\alpha \geq 0$ is a constant. Then (X, d) is a cone metric space [1].

Definition: 1.3 Let (X, d) be a cone metric space, $x \in X$ and $\{x_n\}$ a sequence in X . Then

- (i) $\{x_n\}$ converges to x whenever for every $c \in E$ with $0 \ll c$ there is a natural number N such that $d(x_n, x) \ll c$ for all $n \geq N$.

*Corresponding author: Saurabh Jain[§], *E-mail: sj5129@gmail.com

(ii) $\{x_n\}$ is a Cauchy sequence whenever for every $c \in E$ with $0 \ll c$ there is a natural number N such that $d(x_n, x_m) \ll c$ for all $n, m \geq N$.

Definition: 1.4 Let (X, d) is said to be a complete cone metric space, if every Cauchy sequence is convergent in X .

Let (X, d) be a metric space. We denote by $CB(X)$ the family of non-empty closed bounded subset of X . Let $H(.,.)$ be the Hausdorff distance on $CB(X)$ That is, for,

$$H(A, B) = \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(A, b) \right\}$$

where $d(a, B) = \inf \{d(a, b) : b \in B\}$ is the distance from the point a to the subset B . An element $x \in X$ is said to be a fixed point of a multi-valued mapping $T : X \rightarrow 2^X$ if $x \in T(x)$.

2. MAIN RESULTS:

Theorem: 2.1 Let (X, d) be a complete cone metric space and the mapping $T : X \rightarrow CB(X)$ be multivalued map satisfying for each $x, y \in X$,

$$H(Tx, Ty) \leq a[d(x, Tx) + d(y, Ty)] + b[d(x, Ty) + d(y, Tx)] + c[d(x, y) + d(Tx, Ty)] \text{ for all } x, y \in X,$$

and $a + b + c < \frac{1}{2}$, $a, b, c \in \left[0, \frac{1}{2}\right)$. Then T has a fixed point in X .

Proof: For every $x_0 \in X$ and $n \geq 1$, $x_1 \in T(x_0)$ and $x_{n+1} \in T(x_n)$

$$\begin{aligned} d(x_{n+1}, x_n) &\leq H(Tx_n, Tx_{n+1}) \\ &\leq a[d(x_n, Tx_n) + d(x_{n-1}, Tx_{n-1})] + b[d(x_n, Tx_{n-1}) + d(Tx_n, x_{n-1})] + c[d(x_n, x_{n-1}) + d(Tx_n, Tx_{n-1})] \\ &\leq a[d(x_n, x_{n+1}) + d(x_{n-1}, x_n)] + b[d(x_n, x_n) + d(x_{n+1}, x_{n-1})] + c[d(x_n, x_{n-1}) + d(x_{n+1}, x_n)] \\ &\leq a[d(x_n, x_{n+1}) + d(x_{n-1}, x_n)] + b[d(x_{n+1}, x_{n-1})] + c[d(x_n, x_{n-1}) + d(x_{n+1}, x_n)] \\ &\leq a[d(x_n, x_{n+1}) + d(x_{n-1}, x_n)] + b[d(x_{n+1}, x_n) + d(x_n, x_{n-1})] + c[d(x_n, x_{n-1}) + d(x_{n+1}, x_n)] \\ &\leq (a + b + c)[d(x_n, x_{n+1}) + d(x_{n-1}, x_n)] \end{aligned}$$

$$d(x_{n+1}, x_n) \leq Ld(x_n, x_{n-1}) \text{ where } L = \frac{(a + b + c)}{(1 - (a + b + c))}$$

$$d(x_{n+1}, x_n) \leq L^n d(x_1, x_0)$$

For $n > m$ we have

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_{n-1}) + d(x_{n-1}, x_{n-2}) + \dots + d(x_{m+1}, x_m) \\ &\leq [L^{n-1} + L^{n-2} + \dots + L^m] d(x_1, x_0) \\ &\leq \frac{L^m}{(1 - L)} d(x_1, x_0) \end{aligned}$$

Let $0 \ll h$ be given, choose a natural number N_1 such that $\frac{L^m}{(1 - L)} d(x_0, x_1) \ll h$ for all $m \geq N_1$ this implies $d(x_n, x_m) \ll h$. For $n > m$, $\{x_n\}$ is a Cauchy sequence in (X, d) is a complete cone metric space, there exists

$p \in X$ such that $x_n \rightarrow p$. Choose a natural number N_2 such that $d(x_n, p) \ll \frac{h(1-L)}{3}$, for all $n \geq N_2$. Hence

for $n \geq N_2$ we have $d(x_n, p) \ll \frac{h(1-k)}{3}$ where $k = a + b + c$

$$\begin{aligned} d(Tp, p) &\leq H(Tx_n, Tp) + d(Tx_n, p) \\ &\leq a[d(x_n, Tx_n) + d(p, Tp)] + b[d(x_n, Tp) + d(Tx_n, p)] + c[d(x_n, p) + d(Tx_n, Tp)] + d(x_{n+1}, p) \\ &\leq a[d(x_n, x_{n+1}) + d(p, Tp)] + b[d(x_n, Tp) + d(x_{n+1}, p)] + c[d(x_n, p) + d(x_{n+1}, Tp)] + d(x_{n+1}, p) \\ &\leq a[d(x_n, x_{n+1}) + d(p, Tp)] + b[d(x_n, p) + d(p, Tp) + d(x_{n+1}, p)] \\ &\quad + c[d(x_n, Tp) + d(p, Tp) + d(x_{n+1}, p)] + d(x_{n+1}, p) \end{aligned}$$

$$\begin{aligned} (1-k)d(Tp, p) &\leq kd(x_n, p) + kd(x_{n+1}, p) + d(x_{n+1}, p) \\ &\leq d(x_n, p) + d(x_{n+1}, p) + d(x_{n+1}, p) \end{aligned}$$

$$d(Tp, p) \leq \frac{d(x_n, p) + d(x_{n+1}, p) + d(x_{n+1}, p)}{(1-k)}$$

$$d(Tp, p) \ll \frac{h}{3} + \frac{h}{3} + \frac{h}{3}$$

$$d(Tp, p) \ll h$$

for all $n \geq N_2$, $d(Tp, p) \ll \frac{h}{m}$ for all $m \geq 1$, we get $\frac{h}{m} - d(Tp, p) \in P$, and as $m \rightarrow \infty$ we get $\frac{h}{m} \rightarrow 0$ and P is closed $-d(Tp, p) \in P$, but $d(Tp, p) \in P$, $\therefore d(Tp, p) = 0$ and so $p \in Tp$.

Corollary: 2.1 Let (X, d) be a complete cone metric space and the mapping $T : X \rightarrow CB(X)$ be multivalued map satisfies condition $d(Tx, Ty) \leq a[d(x, Tx) + d(y, Ty)] + b[d(x, Ty) + d(y, Tx)]$ for all $x, y \in X$, where

$a, b \in \left[0, \frac{1}{2}\right)$ are constant. Then T has a fixed point in X .

Proof: The proof of the corollary immediately follows by putting $c = 0$ in the previous theorem.

Corollary: 2.2 Let (X, d) be a complete cone metric space and the mapping $T : X \rightarrow CB(X)$ be multivalued map satisfies condition $d(Tx, Ty) \leq b[d(x, Ty) + d(y, Tx)]$ for all $x, y \in X$, where $b \in \left[0, \frac{1}{2}\right)$ is a constant. Then T has a fixed point in X .

Proof: The proof of the corollary immediately follows by putting a and $c = 0$ in the previous theorem.

REFERENCES:

- [1] Huang Gaung, Zhang Xian, "Cone metric spaces and fixed point theorems of contractive mappings" J. math. Anal. Appl.332 (2007)1468-1476.
- [2] H. Mohebi, Topical functions and their properties in a class of ordered Banach spaces, in continuous Optimization, Current Trends and Modern Applications, Part II, Springer, 2005.pp.343-361.
- [3] H. Mohebi, H. Sadeghi, A.M. Rubinov, Best approximation in a class of normed spaces with star-shaped cone, Numer. Funct. Anal. Optim.27(3-4)(2006)411-436.

[4] R. Krishnakumar, M. Marudai, "Fixed Point Theorems of Multivalued Mappings in Cone Metric Spaces", Int. J. Contemp. Math. Sciences, Vol. 5, 2010, no. 31, 1533 - 1540

[5] Sh. Rezapour, R. Hambarani, "Some notes on the Paper-Cone metric Spaces and fixed point theorems of contractive mappings,"-J.Math.Anal.Appl.345 (2008)719-724.

[6] Seong Hoon Cho, Mi Sun Kim, "Fixed point theorems for general contractive multivalued mappings,"-J. Appl. Math. Informatics Vol.27 (2009)343-350.
