

**A SINGLE SERVER M/G/1 QUEUE WITH SERVICE INTERRUPTION
AND BERNOULLI SCHEDULE SERVER VACATION HAVING GENERAL VACATION
TIME DISTRIBUTION**

N. Balamani*

*Department of Mathematics, Avinashilingam Institute for Home Science and Higher Education
for Women University, Coimbatore -43, Tamil Nadu, India*

E-mail: nbalamani77@gmail.com

(Received on: 10-01-12; Accepted on: 13-02-12)

ABSTRACT

This paper studies an M/G/1 queue with service interruption and Bernoulli schedule server vacation. The server provides essential service to all arriving customers with service time following general distribution. However after a service completion, the server may take a vacation with probability p , or to continue to stay in the system with probability $1 - p$. We consider a general distribution for vacation time. The server is interrupted at random and the duration of attending interruption follows exponential distribution. The steady state solutions have been found by using Supplementary variables technique. Also the mean queue length and the mean waiting time are computed.

Keywords: *M/G/1 queue, Bernoulli Schedule Server Vacation, Steady state, the Mean number in the system, Mean waiting time.*

1. INTRODUCTION:

Vacation queues have been studied by numerous researchers since Levy and Yechiali [5] and Doshi [3, 4] due to their wide applications in manufacturing and telecommunication systems. Vacation queues with c servers have been studied by Tian et al [7]. Choudhury and Borthakur [2] have studied vacation queues with batch arrivals. Multiple vacations have been studied by Tian and Zhang [8].

Madan and Maraghi [6] have studied batch arrival queueing system with random breakdowns and Bernoulli schedule server vacation having general vacation time. They have obtained steady state results in terms of the probability generating functions for the number of customers in the queue. Queues with interruptions were considered by many others. To mention White and Christie [10] have studied queue with interruption. Times of interruptions and services generally distributed are considered by Avi – Itzhak and Naor [1] and Thiruvengadan [9].

In this paper we consider queue with interruption and interrupt it through a different angle. The vacations follow a Bernoulli distribution, that is, after a service completion, the server may go for a vacation with probability p ($0 \leq p \leq 1$) or may continue to serve the next customer, if any, with probability $1 - p$. The vacation time and service time are assumed to be general, while the time of attending interruption follows exponential distribution. Also we assume the customer whose service is interrupted goes back to the head of the queue where the arrivals are Poisson. The customer arrives to the system one by one and are served on a first come first served basis.

2. MATHEMATICAL MODEL:

- Customers arrive at the system one by one according to a Poisson stream with arrival rate λ (>0).
- There is a single server and the service time follows a general (arbitrary) distribution with distribution function $G(s)$ and density function $g(s)$. While serving the customers we assume interruptions arrive at random and assumed to occur according to a Poisson process with mean rate $\alpha > 0$. Let β be the server rate of attending interruption. Further we assume that once the interruption arrives the customer whose service is interrupted comes back to the head of the queue.

***Corresponding author: N. Balamani*, *E-mail: nbalamani77@gmail.com**

- Let $\mu(x)dx$ be the conditional probability of service completion during the interval $(x, x + dx]$, given that the elapsed service time is x , so that

$$\mu(x) = \frac{g(x)}{1-G(x)} \tag{1}$$

and therefore,

$$g(s) = \mu(s) e^{-\int_0^s \mu(x)dx} \tag{2}$$

- As soon as the essential service is completed, then with probability p the server may take vacation of random length, or with probability $1 - p$ he may stay in the system providing service, where $0 \leq p \leq 1$.
- The server's vacation time follows a general (arbitrary) distribution with distribution function $H(r)$ and density function $h(r)$. Let $\gamma(x)dx$ be the conditional probability of a completion of a vacation during the interval $(x, x + dx]$ given that the elapsed vacation time is x , so that

$$\gamma(x) = \frac{h(x)}{1-H(x)} \tag{3}$$

and, therefore

$$h(r) = \gamma(r) e^{-\int_0^r \gamma(x)dx} \tag{4}$$

- On returning from vacation the server instantly starts serving the customer at the head of the queue if any.
- The customers are provided service one by one on a first come first served rule.
- Various stochastic processes involved in the system are assumed to be independent of each other.

3. DEFINITIONS AND NOTATIONS:

We define $P_n(x,t)$ = Probability that at time t , the server is active providing essential service and there are $n (\geq 0)$ customers in the queue excluding one customer in service and the elapsed service time for this customer is x .

Consequently $P_n(t) = \int_0^\infty P_n(x,t) dx$ denotes the probability that at time t there are n customers in the queue excluding the one customer in essential service irrespective of the value of x .

$V_n(x,t)$ = Probability that at time t , there are $n (\geq 0)$ customers in the queue and the server is on vacation with elapsed vacation time is x . Consequently $V_n(t) = \int_0^\infty V_n(x,t) dx$ denotes the probability that at time t there are $n (\geq 0)$ customers in the queue and the server is on vacation irrespective of the value of x .

$R_n(t)$ = Probability that at time t , the server is inactive due to the arrival of interruption

$Q(t)$ = Probability that at time t , there are no customers in the queue or in service and the server is idle but available in the system.

4. STEADY STATE CONDITION:

Let

$$\lim_{t \rightarrow \infty} A_n(x,t) = A_n(x) \qquad \lim_{t \rightarrow \infty} A_n(t) = \lim_{t \rightarrow \infty} \int_0^{\infty} A_n(x,t) dx = A_n$$

$$\lim_{t \rightarrow \infty} \frac{dA_n(t)}{dt} = 0 \qquad n \geq 0$$

Where A= P, V.

$$\lim_{t \rightarrow \infty} Q(t)=Q \qquad \lim_{t \rightarrow \infty} R(t)=R$$

denote the corresponding steady state probabilities.

5. EQUATIONS GOVERNING THE SYSTEM:

According to the mathematical model mentioned above, the system has the following set of differential-difference equations

$$\frac{\partial}{\partial x} P_n(x) + (\lambda + \mu(x) + \alpha) P_n(x) = \lambda P_{n-1}(x), \qquad n = 1, 2, 3 \dots \tag{5}$$

$$\frac{\partial}{\partial x} P_0(x) + (\lambda + \mu(x) + \alpha) P_0(x) = 0 \tag{6}$$

$$\frac{\partial}{\partial x} V_n(x) + (\lambda + \gamma(x)) V_n(x) = \lambda V_{n-1}(x), \qquad n = 1, 2, 3 \dots \tag{7}$$

$$\frac{\partial}{\partial x} V_0(x) + (\lambda + \gamma(x)) V_0(x) = 0 \tag{8}$$

$$(\lambda + \beta) R_n = \lambda R_{n+1} + \alpha \int_0^{\infty} P_{n-1}(x) dx \qquad n=1, 2, 3 \dots \tag{9}$$

$$(\lambda + \beta) R_0 = 0 \tag{10}$$

$$\lambda Q = R_0 \beta + \int_0^{\infty} V_0(x) \gamma(x) dx + (1-p) \int_0^{\infty} P_0(x) \mu(x) dx \tag{11}$$

The above equations are to be solved subject to the following boundary conditions

$$P_n(0) = (1-p) \int_0^{\infty} P_{n+1}(x) \mu(x) dx + \int_0^{\infty} V_{n+1}(x) \gamma(x) dx + \beta R_{n+1} \qquad n=1,2,3 \dots \tag{12}$$

$$P_0(0) = (1-p) \int_0^{\infty} P_1(x) \mu(x) dx + \int_0^{\infty} V_1(x) \gamma(x) dx + \lambda Q + \beta R_1 \tag{13}$$

$$V_n(0) = p \int_0^{\infty} P_n(x) \mu(x) dx \qquad n=0,1,2 \dots \tag{14}$$

6. QUEUE SIZE DISTRIBUTION AT A RANDOM EPOCH:

Defining the following probability generating functions

$$A_q(x, z) = \sum_{n=0}^{\infty} z^n A_n(x) \qquad A_q(z) = \sum_{n=0}^{\infty} z^n A_n$$

Where A=P,V $R_q(z) = \sum_{n=0}^{\infty} z^n R_n$ (15)

Now multiplying the equation (5) by z^n and summing over n from 1 to ∞ , adding to equation (6) and using the generating functions defined in (15) we obtain

$$\frac{\partial}{\partial x} P_q(x, z) + (\lambda - \lambda z + \mu(x) + \alpha) P_q(x, z) = 0 \tag{16}$$

Performing similar operations to equations (7) and (8) and (9) and (10) we get

$$\frac{\partial}{\partial x} V_q(x, z) + (\lambda - \lambda z + \gamma(x)) V_q(x, z) = 0 \tag{17}$$

$$(\lambda - \lambda z + \beta) R_q(z) = \alpha z \int_0^{\infty} P_q(x, z) dx \tag{18}$$

For the boundary conditions we multiply both sides of equation (12) by z^{n+1} , sum over n from 1 to ∞ , multiply both sides of (13) by z , add the two results and using the generating function (15) we obtain

$$\begin{aligned} z P_q(0, z) = & (1-p) \int_0^{\infty} P_q(x, z) \mu(x) dx + \int_0^{\infty} V_q(x, z) \gamma(x) dx + \beta R_q(z) + \lambda Q z \\ & - ((1-p) \int_0^{\infty} P_0(x) \mu(x) dx + \int_0^{\infty} V_0(x) \beta(x) dx + \beta R_0) \end{aligned} \tag{19}$$

using equation (11), equation (19) becomes

$$z P_q(0, z) = (1-p) \int_0^{\infty} P_q(x, z) \mu(x) dx + \int_0^{\infty} V_q(x, z) \gamma(x) dx + \beta R_q(z) + \lambda Q (z-1) \tag{20}$$

Now multiply equation (14) by z^n and sum over n from 0 to ∞ , we get

$$V_q(0, z) = p \int_0^{\infty} P_q(x, z) \mu(x) dx \tag{21}$$

Integrating equation (16) from 0 to x we get

$$P_q(x, z) = P_q(0, z) e^{-(\lambda - \lambda z + \alpha)x - \int_0^x \mu(t) dt} \tag{22}$$

Where $P_q(0, z)$ is given by equation (20). Integrating equation (22) by parts with respect to x yields

$$P_q(z) = P_q(0, z) \left(\frac{1 - G^*(\lambda - \lambda z + \alpha)}{\lambda - \lambda z + \alpha} \right) \tag{23}$$

where $G^*(\lambda - \lambda z + \alpha) = \int_0^{\infty} e^{-(\lambda - \lambda z + \alpha)x} dG(x)$ is the Laplace-stieltjes transform of service time $G(x)$.

Multiplying both sides of equation (22) by $\mu(x)$ and integrating with respect to x we obtain

$$\int_0^{\infty} P_q(x, z) \mu(x) dx = P_q(0, z) G^*(\lambda - \lambda z + \alpha) \tag{24}$$

using equation (24), equation (21) becomes

$$V_q(0, z) = p P_q(0, z) G^*(\lambda - \lambda z + \alpha) \tag{25}$$

Integrating equation (17) from 0 to x we get

$$V_q(x, z) = V_q(0, z) e^{-(\lambda - \lambda z)x - \int_0^x \gamma(t) dt} \quad (26)$$

Substituting for $V_q(0, z)$ from equation (25) in (26) we obtain

$$V_q(x, z) = p P_q(0, z) G^*(\lambda - \lambda z + \alpha) e^{-(\lambda - \lambda z)x - \int_0^x \gamma(t) dt} \quad (27)$$

Integrating equation (27) by parts with respect to x yields

$$V_q(z) = p P_q(0, z) G^*(\lambda - \lambda z + \alpha) \left(\frac{1 - H^*(\lambda - \lambda z)}{\lambda - \lambda z} \right) \quad (28)$$

where $H^*(\lambda - \lambda z) = \int_0^\infty e^{-(\lambda - \lambda z)x} dH(x)$ is the Laplace-stieltjes transform of the vacation time $H(x)$

Multiplying both sides of equation (27) by $\gamma(x)$ and integrating with respect to x we obtain

$$\int_0^\infty V_q(x, z) \gamma(x) dx = p P_q(0, z) G^*(\lambda - \lambda z + \alpha) H^*(\lambda - \lambda z) \quad (29)$$

Using equation (22) in equation (18) we get

$$R_q(z) = \alpha z P_q(0, z) \left(\frac{1 - G^*(\lambda - \lambda z + \alpha)}{\lambda - \lambda z + \alpha} \right) \left(\frac{1}{\lambda + \beta - \lambda z} \right) \quad (30)$$

Assuming $a = (\lambda - \lambda z + \alpha)$, $b = (\lambda - \lambda z + \beta)$ and $c = (\lambda - \lambda z)$

Now using equations (24), (29) and (30) in equation (20) yields

$$P_q(0, z) = \frac{-abcQ}{ab \{z - (1-p)G^*(a) - pG^*(a)H^*(c)\} - \beta\alpha z[1 - G^*(a)]} \quad (31)$$

Substituting $P_q(0, z)$ in equations (23), (28) and (30) we obtain

$$P_q(z) = \frac{-bcQ(1 - G^*(a))}{ab \{z - (1-p)G^*(a) - pG^*(a)H^*(c)\} - \beta\alpha z[1 - G^*(a)]} \quad (32)$$

$$V_q(z) = \frac{abpQG^*(a)[H^*(c) - 1]}{ab \{z - (1-p)G^*(a) - pG^*(a)H^*(c)\} - \beta\alpha z[1 - G^*(a)]} \quad (33)$$

$$R_q(z) = \frac{-\alpha zcQ(1 - G^*(a))}{ab \{z - (1-p)G^*(a) - pG^*(a)H^*(c)\} - \beta\alpha z[1 - G^*(a)]} \quad (34)$$

Let $S_q(z)$ denote the probability generating function of the queue size irrespective of the state of the system. Then adding equations (32), (33) and (34), we obtain

$$S_q(z) = Pq(z) + Vq(z) + Rq(z) = \frac{-Qc[1-G^*(a)][b+\alpha z] + abpQG^*(a)[H^*(c)-1]}{ab\{z-(1-p)G^*(a) - pG^*(a)H^*(c)\} - \beta\alpha z[1-G^*(a)]} \quad (35)$$

It is easy to verify that for $z = 1$, $S_q(z)$ is indeterminate of the form 0/0 Hence we apply L'Hopitals rule on equation (35), we obtain

$$S_q(1) = \frac{\lambda Q[(\alpha + \beta)(1-G^*(\alpha)) + \alpha p \beta G^*(\alpha) E(V)]}{\alpha \beta G^*(\alpha) - \lambda[1-G^*(\alpha)](\alpha + \beta) - p \alpha \beta \lambda G^*(\alpha) E(V)} \quad (36)$$

Therefore adding Q to equation (36) and equating to 1 and on simplifying, we get

$$Q = 1 - \lambda \left[\frac{1}{\alpha G^*(\alpha)} + \frac{1}{G^*(\alpha)\beta} - \frac{1}{\alpha} - \frac{1}{\beta} + pE(V) \right] \quad (37)$$

And hence the utilization factor ρ of the system is given by

$$\rho = \lambda \left[\frac{1}{\alpha G^*(\alpha)} + \frac{1}{G^*(\alpha)\beta} - \frac{1}{\alpha} - \frac{1}{\beta} + pE(V) \right] \quad (38)$$

where $\rho < 1$ is the stability condition under which the steady state exists. Equation (37) gives the probability that the server is idle. Substituting for Q from (37) into (35), we have completely and explicitly determined $S_q(z)$, the probability generating function of the queue size.

7. THE MEAN NUMBER IN THE SYSTEM:

Let L_q denote the mean number of customers in the queue under the steady state. Then

$$L_q = \frac{d}{dz} S_q(z) \Big|_{z=1} \quad (39)$$

Since this formula gives 0/0 form, then we write $S_q(z)$ given in (35) as $S_q(z) = N(z) / D(z)$ where $N(z)$ and $D(z)$ are the numerator and denominator of the right hand side of (35) respectively. Then using the L'Hopital's rule twice we obtain

$$L_q = \lim_{z \rightarrow 1} \frac{D'(z)N''(z) - N'(z)D''(z)}{2(D'(z))^2} = \lim_{z \rightarrow 1} \frac{D'(1)N''(1) - N'(1)D''(1)}{2(D'(1))^2} \quad (40)$$

where, primes and double primes in (40) denote first and second derivatives at $z = 1$. Carrying out the derivatives at $z = 1$, using the fact that $H^*(0) = 1$, $H^{*'}(0) = -E(V)$ and $H^{*''}(0) = E(V^2)$, the second moment of the vacation time.

After simplification we get

$$N'(1) = \lambda Q \{(\alpha + \beta) + G^*(\alpha) [E(V)p\alpha\beta - \alpha - \beta]\} \quad (41)$$

$$N''(1) = 2\lambda^2 Q [(-1 + \alpha/\lambda) + G^*(\alpha) [1 - \alpha/\lambda - p\alpha E(V) - p\beta E(V) + 1/2 p\alpha\beta E(V^2)]] + G^{*'}(\alpha)(\alpha + \beta - \alpha\beta E(V)) \quad (42)$$

$$D'(1) = -\lambda(\alpha + \beta) + G^*(\alpha) [\alpha\beta + \lambda(\alpha + \beta - p\alpha\beta E(V))] \quad (43)$$

$$D''(1) = 2\lambda^2 \left\{ \left(1 - \frac{\alpha + \beta}{\lambda} \right) + G^*(\alpha) [-1 + p\alpha E(V) + p\beta E(V) - 1/2 p\alpha\beta E(V^2)] + G^{**}(\alpha) \left[-\frac{\alpha\beta}{\lambda} - \alpha - \beta + \alpha\beta pE(V) \right] \right\} \quad (44)$$

Using equations (41) to (44) into (40), we obtain L_q in closed form where Q has been found in equation (37). Further we find the average system size L using Little's formula. Thus we have

$$L = L_q + \rho \quad (45)$$

where L_q has been found in equation (40) and ρ is obtained from equation (38) as ρ

$$\rho = 1 - Q \quad (46)$$

8. THE MEAN WAITING TIME:

Let W_q and W denote the mean waiting time in the queue and in the system respectively. Then using Little's formula, we obtain

$$W_q = \frac{L_q}{\lambda} \quad (47)$$

$$W = \frac{L}{\lambda} \quad (48)$$

Where L_q and L have been found in equations (40) and (45)

REFERENCES:

- [1] B.Avi – Itzhak, P.Naor, Some queueing problems with the service station subject to breakdowns, *Operations Research* 11(3) (1963), 303 – 319.
- [2] A. Borthakur and G. Choudhury, On a batch arrival Poisson queue with generalised vacation, *Sankhya Ser. B*, 59 (1997), 369-383.
- [3] B. T. Doshi, A note on stochastic decomposition in a GI/G/1 queue with vacations or setup times, *Journal of Applied Probability*, 22(1985), 419-428.
- [4] B. T. Doshi, Queueing systems with vacations - A survey, *Queueing Systems*, 1(1986), 29-66.
- [5] Y.Levy and U. Yechiali, Utilization of idle time in an M/G/1 queueing system, *Management Science*, 22(1975), 202-211.
- [6] F.A. Maraghi, K.C. Madan and K.D. Dowman, Batch arrival queueing system with random breakdowns and Bernoulli schedule server vacations having general time distribution, *International Journal of Information and Management Sciences*, 20 (2009), 55-70
- [7] N. Tian, Q. Li and J. Cao, Conditional stochastic decomposition in M/M/c queue with server vacations, *Stochastic Models*, 15 (1999), 367-377.
- [8] N. Tian and Z.G. Zhang, The discrete-time GI/Geo/1 queue with multiple vacations, *Queueing Systems*, 40 (2002), 283-294.
- [9] K. Thiruvengadam, Queueing with breakdowns, *Operations Research* 11(1) (1963), 62 – 71.
- [10] H. White, L. Christie, Queueing with preemptive priorities or with breakdowns, *Operations Research* 6 (1) (1958) 79 – 95.
