

**ORDERED INTUITIONISTIC FUZZY  
PRE SEMI BASICALLY DISCONNECTED SPACES**

**D. Amsaveni\*, M. K. Uma and E. Roja**

*Sri Sarada College for Women, Salem-636016, Tamilnadu, India*

*E-mail: [d\\_amsaveni@rediffmail.com](mailto:d_amsaveni@rediffmail.com)*

*(Received on: 13-01-12; Accepted on: 17-02-12)*

**ABSTRACT**

*In this paper, a new class of intuitionistic fuzzy topological spaces called ordered intuitionistic fuzzy pre semi basically disconnected spaces is introduced. Tietze extension theorem for ordered intuitionistic fuzzy pre semi basically disconnected spaces has been discussed as in [15] and several other properties are also discussed.*

**Key words:** *Ordered intuitionistic fuzzy pre semi basically disconnected spaces, ordered intuitionistic fuzzy pre semi continuous functions, and lower  $F_{\sigma}$  (resp. upper  $F_{\sigma}$ ) intuitionistic fuzzy pre semi continuous functions.*

**1. INTRODUCTION:**

After the introduction of the concept of fuzzy sets by Zadeh [18], several researches were conducted on the generalizations of the notion of fuzzy set. The concept of ‘‘Intuitionistic fuzzy sets’’ was first published by Atanassov [2] and many works by the same author and his colleagues appeared in the literature [3-5]. Later this concept was generalized to ‘‘Intuitionistic L-fuzzy sets’’ by Atanassov and stoeva [6]. An introduction to intuitionistic fuzzy topological space was introduced by Dogan Coker [9]. Several types of fuzzy connectedness in intuitionistic fuzzy topological spaces were defined by Coker (1997). The construction is based on the idea of intuitionistic fuzzy set developed by Atanassov (1983, 1986; Atanassov and Stoeva, 1983). In this paper a new class of intuitionistic fuzzy topological spaces namely, ordered intuitionistic fuzzy pre semi basically disconnected spaces is introduced by using the concepts of [11,13,15]. ‘Intuitionistic fuzzy pre semi closed sets’ was introduced by [1]. Tietze extension theorem for ordered intuitionistic fuzzy pre semi basically disconnected spaces has been discussed as in [15].

**2. PRELIMINARIES:**

Throughout this paper let  $X$  be a non empty set and  $I = [0, 1]$ .

**Definition: 2.1[4]** Let  $X$  be a non empty fixed set. An intuitionistic fuzzy set (IFS for short)  $A$  is an object having the form  $A = \{ \langle x, \mu_A(x), \gamma_A(x) \rangle : x \in X \}$  where the functions  $\mu_A : X \rightarrow I$  and  $\gamma_A : X \rightarrow I$  denote the degree of membership (namely  $\mu_A(x)$ ) and the degree of non membership (namely  $\gamma_A(x)$ ) of each element  $x \in X$  to the set  $A$ , respectively, and  $0 \leq \mu_A(x) + \gamma_A(x) \leq 1$  for each  $x \in X$ .

**Remark: 2.1[4]** For the sake of simplicity, we shall use the symbol  $A = \langle x, \mu_A, \gamma_A \rangle$ .

**Definition: 2.2[9]** Let  $X$  be a non empty fixed set. Then,  $0_{\sim} = \{ \langle x, 0, 1 \rangle : x \in X \}$  and  $1_{\sim} = \{ \langle x, 1, 0 \rangle : x \in X \}$ .

**Definition: 2.3[9]** Let  $X$  be a non empty set. An intuitionistic fuzzy topology (IFT for short) on a non empty set  $X$  is a family  $\tau$  of intuitionistic fuzzy sets (IFSs for short) in  $X$  satisfying the following axioms:

(T<sub>1</sub>)  $0_{\sim}, 1_{\sim} \in \tau$ ,

(T<sub>2</sub>)  $G_1 \cap G_2 \in \tau$  for any  $G_1, G_2 \in \tau$ ,

**\*Corresponding author: D. Amsaveni\*, \*E-mail: [d\\_amsaveni@rediffmail.com](mailto:d_amsaveni@rediffmail.com)**

$$(T_3) \bigcup_{G_i \in \tau} G_i \in \tau \text{ for any arbitrary family } \{G_i : i \in J\}.$$

In this case the pair  $(X, \tau)$  is called an intuitionistic fuzzy topological space (IFTS for short) and any  $IFS$  in  $\tau$  is known as an intuitionistic fuzzy open set (IFOS for short) in  $X$ .

**Definition: 2.4[9]** Let  $(X, \tau)$  be an IFTS and  $A = \langle x, \mu_A, \gamma_A \rangle$  be an IFS in  $X$ . Then the fuzzy interior and fuzzy closure of  $A$  are defined by

$$cl(A) = \bigcap \{K : K \text{ is an IFCS in } X \text{ and } A \subseteq K\}, \text{ int}(A) = \bigcup \{G : G \text{ is an IFOS in } X \text{ and } G \subseteq A\}.$$

**NOTATION: 2.1** An IFTS  $(X, T)$  represent intuitionistic fuzzy topological spaces and for a subset  $A$  of a space  $(X, T)$ ,  $IFcl(A)$ ,  $IFint(A)$ ,  $IFPScI(A)$ ,  $IFPSint(A)$ ,  $IFspint(A)$ ,  $IFspcl(A)$  and  $\bar{A}$  denote an intuitionistic fuzzy closure of  $A$ , an intuitionistic fuzzy interior of  $A$ , intuitionistic fuzzy pre semi closure of  $A$ , an intuitionistic fuzzy semi pre interior of  $A$ , an intuitionistic fuzzy pre semi interior of  $A$ , an intuitionistic fuzzy semi pre closure of  $A$  and the complement of  $A$  in  $X$  respectively.

**Definition: 2.5 [17]** A subset  $A$  of an IFTS  $(X, T)$  is called an IF semi pre open set if  $A \subseteq IFcl(IFint(IFcl(A)))$  and an IF semi pre closed set if  $IFint(IFcl(IFint(A))) \subseteq A$ ;

**Definition: 2.6[14]** A subset  $A$  of an IFTS  $(X, T)$  is called an IF generalized closed (briefly IF g-closed) set if  $IFcl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is IF open in  $(X, T)$ . The complement of an IF g-closed set is called an IF g-open set;

**Definition: 2.7[1]** A subset  $A$  of an IFTS  $(X, T)$  is called intuitionistic fuzzy pre semi closed (IF pre semi closed for short) if  $IFspcl(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is IF g-open in  $(X, T)$ .

**Definition: 2.8[1]** A subset  $A$  of an IFTS  $(X, T)$  is called intuitionistic fuzzy pre semi open (IF pre semi open for short) if  $\bar{A}$  is IF pre semi closed.

**Definition: 2.9[1]** A function  $f : (X, T) \rightarrow (Y, S)$  is called intuitionistic fuzzy pre semi continuous (IF pre semi continuous for short) if  $f^{-1}(V)$  is an IF pre semi closed set of  $(X, T)$  for every IF closed set  $V$  of  $(Y, S)$ .

**Definition: 2.10** Let  $(X, T)$  be an IFTS. Let  $A$  be any IF pre semi open  $F_G$  in  $(X, T)$ . If IF pre semi closure of  $A$  is IF pre semi open, then  $(X, T)$  is said to be intuitionistic fuzzy pre semi basically disconnected (for short, IF pre semi basically disconnected).

**Definition: 2.11[11]** An ordered set on which there is given a fuzzy topology is called an ordered fuzzy topological space.

### 3. ORDERED INTUITIONISTIC FUZZY PRE SEMI BASICALLY DISCONNECTED SPACES:

In this section, the concept of ordered intuitionistic fuzzy pre semi basically disconnected spaces is introduced. Some of its characterizations and properties are studied.

**Definition: 3.1** An ordered set on which there is given an intuitionistic fuzzy topology is called an ordered intuitionistic fuzzy topological space (for short ordered IFTS).

**Definition: 3.2** An IFS  $A$  in a partially ordered set  $(X, T, \leq)$  is said to be an

(a) Increasing IFS if  $x \leq y$  implies  $A(x) \subseteq A(y)$ .

That is,  $\mu_A(x) \leq \mu_A(y)$  and  $\gamma_A(x) \geq \gamma_A(y)$ .

(b) Decreasing IFS if  $x \leq y$  implies  $A(x) \supseteq A(y)$ .

That is,  $\mu_A(x) \geq \mu_A(y)$  and  $\gamma_A(x) \leq \gamma_A(y)$ .

**Definition: 3.3** Let  $(X, T, \leq)$  be an intuitionistic fuzzy topological space (for short IFTS) and  $A$  be an intuitionistic fuzzy set (for short IFS) in  $(X, T)$ .  $A$  is called an intuitionistic fuzzy  $G_\delta$  (for short IF  $G_\delta$ ) if  $A = \bigcap_{i=1}^{\infty} A_i$  where each  $A_i$  is IF open.

**Definition: 3.4** Let  $(X, T, \leq)$  be an intuitionistic fuzzy topological space (for short IFTS) and  $A$  be an intuitionistic fuzzy set (for short IFS) in  $(X, T, \leq)$ .  $A$  is called an intuitionistic fuzzy  $F_\sigma$  (IF  $F_\sigma$ ) if  $A = \bigcup_{i=1}^{\infty} \overline{A_i}$  where each  $\overline{A_i}$  is IF open.

**Definition: 3.5** Let  $(X, T, \leq)$  be an ordered IFTS and let  $A$  be any IFS in  $(X, T, \leq)$ ,  $A$  is called increasing IF pre semi open if  $IF\ sp\ int(A) \supseteq U$  whenever  $A \supseteq U$  and  $U$  is IF  $g$ -closed in  $(X, T, \leq)$ . The complement of an increasing IF pre semi open set is called decreasing IF pre semi closed.

**Note: 3.1** (a) Let  $(X, T, \leq)$  be an ordered IFTS. An IFS  $A$  in  $(X, T, \leq)$  which is both intuitionistic fuzzy pre semi open and IF  $F_\sigma$  is denoted by IF pre semi open  $F_\sigma$ .

(b) Let  $(X, T, \leq)$  be an IFTS. An IFS  $A$  in  $(X, T, \leq)$  which is both intuitionistic fuzzy pre semi closed (for short IF pre semi closed) and IF  $G_\delta$  is denoted by IF pre semi closed  $G_\delta$ .

(c) An IFS  $A$  which is both IF pre semi open  $F_\sigma$  and IF pre semi closed  $G_\delta$  is denoted by intuitionistic fuzzy pre semi COGF (IF pre semi COGF).

**Definition: 3.6** Let  $(X, T, \leq)$  be an IFTS. For any IFS  $A$  in  $(X, T, \leq)$ ,

$$I^{IFPSG_\delta}(A) = \text{increasing intuitionistic fuzzy pre semi closure } G_\delta \text{ of } A \\ = \bigcap \{ B / B \text{ is an increasing intuitionistic fuzzy pre semi closed } G_\delta \text{ set and } B \supseteq A \},$$

$$D^{IFPSG_\delta}(A) = \text{decreasing intuitionistic fuzzy pre semi } G_\delta \text{ closure of } A \\ = \bigcap \{ B / B \text{ is a decreasing intuitionistic fuzzy pre semi closed } G_\delta \text{ set and } B \supseteq A \},$$

$$I^{0IFPSF_\sigma}(A) = \text{increasing intuitionistic fuzzy pre semi } F_\sigma \text{ interior of } A \\ = \bigcup \{ B / B \text{ is an increasing intuitionistic fuzzy pre semi open } F_\sigma \text{ set and } B \subseteq A \},$$

$$D^{0IFPSF_\sigma}(A) = \text{decreasing intuitionistic fuzzy pre semi } F_\sigma \text{ interior of } A \\ = \bigcup \{ B / B \text{ is a decreasing intuitionistic fuzzy pre semi open } F_\sigma \text{ set and } B \subseteq A \}.$$

Clearly,  $I^{IFPSG_\delta}(A)$  (resp.  $D^{IFPSG_\delta}(A)$ ) is the smallest increasing (resp. decreasing) intuitionistic fuzzy pre semi closed  $G_\delta$  set containing  $A$  and  $I^{0IFPSF_\sigma}(A)$  (resp.  $D^{0IFPSF_\sigma}(A)$ ) is the largest increasing (resp. decreasing) intuitionistic fuzzy pre semi open  $F_\sigma$  set contained in  $A$ .

**Definition: 3.7** Let  $(X, T, \leq)$  be an *IFTS*. Let  $A$  be any *IF* pre semi open  $F_\sigma$  in  $(X, T, \leq)$ . If  $\overline{A}$  is *IF* pre semi open  $F_\sigma$ , then  $(X, T, \leq)$  is said to be intuitionistic fuzzy pre semi basically disconnected (for short, *IF* pre semi basically disconnected).

**Proposition: 3.1** For any *IFS*  $A$  of an ordered intuitionistic fuzzy topological space  $(X, T, \leq)$ , the following statements are hold.

- (a)  $\overline{I^{IFPSG_\delta}(A)} = D^{0IFPSF_\sigma}(\overline{A})$ .
- (b)  $\overline{D^{IFPSG_\delta}(A)} = I^{0IFPSF_\sigma}(\overline{A})$ .
- (c)  $\overline{I^{0IFPSF_\sigma}(A)} = D^{IFPSG_\delta}(\overline{A})$ .
- (d)  $\overline{D^{0IFPSF_\sigma}(A)} = I^{IFPSG_\delta}(\overline{A})$ .

**Proof: (a):** Since  $I^{IFPSG_\delta}(A)$  is an increasing *IF* pre semi closed  $G_\delta$  set containing  $A$ ,  $\overline{I^{IFPSG_\delta}(A)}$  is a decreasing *IF* pre semi open  $F_\sigma$  set such that  $\overline{I^{IFPSG_\delta}(A)} \subseteq \overline{A}$ . Let  $B$  be another decreasing *IF* pre semi open  $F_\sigma$  set such that  $B \subseteq \overline{A}$ . Then  $\overline{B}$  is an increasing *IF* pre semi closed  $G_\delta$  set such that  $\overline{B} \supseteq A$ . It follows that  $I^{IFPSG_\delta}(A) \subseteq \overline{B}$ . That is,  $B \subseteq \overline{I^{IFPSG_\delta}(A)}$ . Thus,  $\overline{I^{IFPSG_\delta}(A)}$  is the largest decreasing *IF* pre semi open  $F_\sigma$  set such that  $\overline{I^{IFPSG_\delta}(A)} \subseteq \overline{A}$ . That is,  $\overline{I^{IFPSG_\delta}(A)} = D^{0IFPSF_\sigma}(\overline{A})$ .

The proofs of (b), (c) and (d) can be proved in a similar manner.

**Definition: 3.8** Let  $(X, T, \leq)$  be an ordered *IFTS*. Let  $A$  be any increasing (resp. decreasing) *IF* pre semi open  $F_\sigma$  (resp. *IF* pre semi closed  $G_\delta$ ) set in  $(X, T, \leq)$ . If  $I^{IFPSG_\delta}(A)$  (resp.  $D^{IFPSG_\delta}(A)$ ) is increasing (resp. decreasing) *IF* pre semi open  $F_\sigma$  in  $(X, T, \leq)$ , then  $(X, T, \leq)$  is said to be upper  $F_\sigma$  (resp. lower  $F_\sigma$ ) intuitionistic fuzzy pre semi basically disconnected (for short *IF* pre semi basically disconnected). An *IFTS*  $(X, T, \leq)$  is said to be ordered *IF* pre semi basically disconnected if it is both upper  $F_\sigma$  and lower  $F_\sigma$  intuitionistic fuzzy pre semi basically disconnected.

**Proposition: 3.2** For an *IFTS*  $(X, T, \leq)$ , the followings are equivalent:

- (a)  $(X, T, \leq)$  is upper  $F_\sigma$  *IF* pre semi basically disconnected.
- (b) For each decreasing *IF* pre semi closed  $G_\delta$  set  $A$ ,  $D^{0IFPSF_\sigma}(A)$  is decreasing *IF* pre semi closed  $G_\delta$ .
- (c) For each increasing *IF* pre semi open  $F_\sigma$  set  $A$ ,  $\overline{D^{IFPSG_\delta}(I^{IFPSG_\delta}(A))} = I^{IFPSG_\delta}(A)$ .
- (d) For each pair of increasing *IF* pre semi open  $F_\sigma$  set  $A$  and decreasing *IF* pre semi open  $F_\sigma$  set  $B$  in  $(X, T, \leq)$  with  $\overline{I^{IFPSG_\delta}A} = B$ ,  $\overline{D^{IFPSG_\delta}B} = I^{IFPSG_\delta}A$ .

**Proof: (a)  $\Rightarrow$  (b):** Let  $A$  be any decreasing *IF* pre semi closed  $G_\delta$  set. We claim that  $D^{0IFPSF_\sigma}(A)$  is decreasing *IF* pre semi closed  $G_\delta$ . Now,  $\overline{A}$  is increasing *IF* pre semi open  $F_\sigma$ . So by assumption (a),  $\overline{I^{IFPSG_\delta}(\overline{A})}$  is increasing *IF* pre semi open  $F_\sigma$ . That is,  $\overline{D^{0IFPSF_\sigma}(A)}$  is decreasing *IF* pre semi closed  $G_\delta$ .

**(b)  $\Rightarrow$  (c):** Let  $A$  be an increasing *IF* pre semi open  $F_\sigma$  set. Then,  $\overline{I^{IFPSG_\delta}(A)} = D^{0IFPSF_\sigma}(\overline{A})$ . Consider  $\overline{I^{IFPSG_\delta}(A) + D^{IFPSG_\delta}(\overline{I^{IFPSG_\delta}(A)})} = I^{IFPSG_\delta}(A) + D^{IFPSG_\delta}(D^{0IFPSF_\sigma}(\overline{A}))$ . Since  $A$  is increasing *IF* pre

semi open  $F_\sigma$ ,  $\bar{A}$  is a decreasing IF pre semi closed  $G_\delta$  set and by (b),  $D^{0IFPSF_\sigma}(\bar{A})$  is a decreasing IF pre semi closed  $G_\delta$  set. Therefore,  $D^{IFPSG_\delta}(D^{0IFPSF_\sigma}(\bar{A})) = D^{0IFPSF_\sigma}(\bar{A})$ . Now,  $I^{IFPSG_\delta}(A) + D^{IFPSG_\delta}(D^{0IFPSF_\sigma}(\bar{A})) = I^{IFPSG_\delta}(A) + (D^{0IFPSF_\sigma}(\bar{A})) = I^{IFPSG_\delta}(A) + \overline{I^{IFPSG_\delta}(A)}$ .

Hence,  $D^{IFPSG_\delta}(D^{0IFPSF_\sigma}(\bar{A})) = \overline{I^{IFPSG_\delta}(A)}$ . By Proposition 3.1, (c) holds.

(c)  $\Rightarrow$  (d): Let  $A$  be any increasing IF pre semi open  $F_\sigma$  set and  $B$  be any decreasing IF pre semi open  $F_\sigma$  set such that  $\overline{I^{IFPSG_\delta}(A)} = B$ . By Proposition 3.1,  $\bar{B} = D^{0IFPSF_\sigma}(\bar{A})$ .

$$\text{By (c), } D^{IFPSG_\delta}(\overline{I^{IFPSG_\delta}(A)}) = D^{IFPSG_\delta}(D^{0IFPSF_\sigma}(\bar{A})) = B \tag{3.1}$$

$$\text{Therefore, } D^{IFPSG_\delta}(\overline{I^{IFPSG_\delta}(A)}) = D^{IFPSG_\delta}(B) \tag{3.2}$$

From (3.1) and (3.2), we have,  $\overline{I^{IFPSG_\delta}(A)} = D^{IFPSG_\delta}(B)$ .

(d)  $\Rightarrow$  (a): Let  $A$  be any increasing IF pre semi open  $F_\sigma$  set. Let  $\overline{I^{IFPSG_\delta}(A)} = B$ . From (d), it follows that  $\overline{D^{IFPSG_\delta}(B)} = I^{IFPSG_\delta}(A)$ . Hence,  $(X, T, \leq)$  is upper  $F_\sigma$  IF pre semi basically disconnected space.

**Proposition: 3.3** Let  $(X, T, \leq)$  be an ordered IFTS. Then  $(X, T, \leq)$  is an upper  $F_\sigma$  IF pre semi basically disconnected space if and only if for any decreasing IF pre semi open  $F_\sigma$  set  $A$  and decreasing IF pre semi closed  $G_\delta$  set  $B$  such that  $A \subseteq B$ ,  $D^{IFPSG_\delta}(A) \subseteq D^{0IFPSF_\sigma}(B)$ .

**Notation: 3.1** An ordered IFS which is both decreasing (resp. increasing) IF pre semi open  $F_\sigma$  and IF pre semi closed  $G_\delta$  is called a decreasing (resp. increasing) IF pre semi clopen set (for short IF pre semi COGF).

**Remark: 3.1** Let  $(X, T, \leq)$  be an upper  $F_\sigma$  IF pre semi basically disconnected space. Let  $\{A_i, \bar{B}_i / i \in N\}$  be a collection such that  $A_i$ 's are decreasing IF pre semi open  $F_\sigma$  sets,  $B_i$ 's are decreasing IF pre semi closed  $G_\delta$  sets and let  $A, \bar{B}$  be decreasing IF pre semi open  $F_\sigma$  and increasing IF pre semi open  $F_\sigma$  sets respectively. If  $A_i \subseteq A \subseteq B_j$  and  $A_i \subseteq B \subseteq B_j$  for all  $i, j \in N$ , then there exists a decreasing IF pre semi COGF set  $C$  such that  $D^{IFPSG_\delta}(A_i) \subseteq C \subseteq D^{0IFPSF_\sigma}(B_j)$  for all  $i, j \in N$ .

**Proposition: 3.4** Let  $(X, T, \leq)$  be an upper  $F_\sigma$  IF pre semi basically disconnected space. Let  $(A_q)_{q \in Q}$  and  $(B_q)_{q \in Q}$  be the monotone increasing collections of decreasing IF pre semi open  $F_\sigma$  sets and decreasing IF pre semi closed  $G_\delta$  sets of  $(X, T, \leq)$  respectively and suppose that  $A_{q_1} \subseteq B_{q_2}$  whenever  $q_1 < q_2$  ( $Q$  is the set of rational numbers). Then there exists a monotone increasing collection  $\{C_q\}_{q \in Q}$  of decreasing IF pre semi COGF sets of  $(X, T, \leq)$  such that  $D^{IFPSG_\delta}(A_{q_1}) \subseteq C_{q_2}$  and  $C_{q_1} \subseteq D^{0IFPSF_\sigma}(B_{q_2})$  whenever  $q_1 < q_2$ .

#### 4. PROPERTIES AND CHARACTERIZATIONS OF INTUITIONISTIC FUZZY PRE SEMI BASICALLY DISCONNECTED SPACES:

In this section various properties and characterizations of intuitionistic fuzzy pre semi basically disconnected spaces are discussed.

**Definition: 4.1** Let  $(X, T, \leq)$  be an ordered *IFTS*. A function  $f : X \rightarrow R(I)$  is called lower  $F_{\sigma}$  (resp. upper  $F_{\sigma}$ ) *IF* pre semi continuous, if  $f^{-1}(R_t)$  (resp.  $f^{-1}(L_t)$ ) is an increasing or decreasing *IF* pre semi open  $F_{\sigma}$  set (resp. *IF* pre semi open  $F_{\sigma}$  / *IF* pre semi closed  $G_{\delta}$ ) for each  $t \in R$ .

**Proposition: 4.1** Let  $(X, T, \leq)$  be an ordered *IFTS*. Let  $A$  be an *IFS* in  $X$ , and let  $f : X \rightarrow R(I)$  be such that

$$f(x)(t) = \begin{cases} 1 & \text{if } t < 0, \\ A(x) & \text{if } 0 \leq t \leq 1, \\ 0 & \text{if } t > 1, \end{cases}$$

for all  $x \in X$ . Then  $f$  is lower  $F_{\sigma}$  (resp. upper  $F_{\sigma}$ ) *IF* pre semi continuous iff  $A$  is an increasing or decreasing *IF* pre semi open  $F_{\sigma}$  (resp. *IF* pre semi open  $F_{\sigma}$  / *IF* pre semi closed  $G_{\delta}$ ) set.

**Definition: 4.2** The characteristic function of any *IFS*  $A$  in  $X$  is the function  $\chi_A : X \rightarrow I(L)$  defined by  $\chi_A(x) = (A(x))$ ,  $x \in X$ .

**Proposition: 4.2** Let  $(X, T, \leq)$  be an ordered *IFTS*, and let  $A$  be an *IFS* in  $X$ . Then  $\chi_A$  is lower  $F_{\sigma}$  (resp. upper  $F_{\sigma}$ ) *IF* pre semi continuous iff  $A$  is an increasing or decreasing *IF* pre semi open  $F_{\sigma}$  (resp. *IF* pre semi open  $F_{\sigma}$  / *IF* pre semi closed  $G_{\delta}$ ) set.

**Proof:** The proof follows from Proposition 4.1.

**Definition: 4.3** Let  $(X, T, \leq)$  and  $(Y, S, \leq)$  be *IFTS*s. A function  $f : (X, T, \leq) \rightarrow (Y, S, \leq)$  is called increasing  $F_{\sigma}$  (resp. decreasing  $G_{\delta}$ ) intuitionistic fuzzy strongly pre semi continuous (for short, increasing  $F_{\sigma}$  (resp. decreasing  $G_{\delta}$ ) *IF* strongly pre semi continuous) if  $f^{-1}(A)$  is increasing  $F_{\sigma}$  (resp. decreasing  $G_{\delta}$ ) *IF* pre semi clopen in  $(X, T, \leq)$  for every *IF* pre semi open  $F_{\sigma}$  set in  $(Y, S, \leq)$ . If  $f$  is both increasing  $F_{\sigma}$  and decreasing  $G_{\delta}$  *IF* strongly pre semi continuous, then it is called ordered *IF* strongly pre semi continuous.

**Proposition: 4.3** Let  $(X, T, \leq)$  be an ordered *IFTS*. Then the following statements are equivalent:

- $(X, T, \leq)$  is upper  $F_{\sigma}$  *IF* pre semi basically disconnected,
- If  $g, h : X \rightarrow R(I)$ ,  $g$  is lower  $F_{\sigma}$  *IF* pre semi continuous,  $h$  is upper  $F_{\sigma}$  *IF* pre semi continuous and  $g \subseteq h$  then there exists an increasing  $F_{\sigma}$  *IF* strongly pre semi continuous function,  $f : (X, T, \leq) \rightarrow R(I)$  such that  $g \subseteq f \subseteq h$ .
- If  $\overline{A}$  is increasing *IF* pre semi open  $F_{\sigma}$  set and  $B$  is decreasing *IF* pre semi open  $F_{\sigma}$  set such that  $B \subseteq A$ , then there exists an increasing  $F_{\sigma}$  *IF* strongly pre semi continuous function  $f : (X, T, \leq) \rightarrow [0, 1](I)$  such that  $B \subseteq \overline{L_1}f \subseteq R_0f \subseteq A$ .

**Proof:** (a)  $\Rightarrow$  (b) can be established by the concepts of increasing  $F_{\sigma}$  (resp. decreasing  $G_{\delta}$ ) IF pre semi clopen set in  $(X, T, \leq)$  and the theorem 3.7 of Kubiak [15] with some slight suitable modifications.

(b)  $\Rightarrow$  (c): Suppose  $\overline{A}$  is an increasing IF pre semi open  $F_{\sigma}$  set and  $B$  is an decreasing IF pre semi closed  $G_{\delta}$  set, such that  $B \subseteq A$ . Then  $\chi_B \subseteq \chi_A$  and  $\chi_B, \chi_A$  are lower  $F_{\sigma}$  and upper  $F_{\sigma}$  IF pre semi continuous functions respectively. Hence by (b), there exists an increasing  $F_{\sigma}$  IF strongly pre semi continuous function  $f : (X, T, \leq) \rightarrow R(I)$  such that  $\chi_B \subseteq f \subseteq \chi_A$ . Clearly,  $f(x) \in [0, 1](I)$  for all  $x \in X$  and  $B = (\overline{L_1})\chi_B \subseteq (\overline{L_1})f \subseteq R_0f \subseteq R_0\chi_A = A$ .

(c)  $\Rightarrow$  (a): This follows from Proposition 3.2 and the fact that  $(\overline{L_1})f$  and  $R_0f$  are decreasing IF pre semi closed  $G_{\delta}$  and decreasing IF pre semi open  $F_{\sigma}$  sets respectively.

### 5. TIETZE EXTENSION THEOREM FOR ORDERED IF PRE SEMI BASICALLY DISCONNECTED SPACES:

In this section Tietze extension theorem for ordered IF pre semi basically disconnected spaces is studied.

**Proposition: 5.1** Let  $(X, T, \leq)$  be an upper  $F_{\sigma}$  IF pre semi basically disconnected space and let  $A \subset X$  be such that  $\chi_A$  is increasing IF pre semi open  $F_{\sigma}$  set in  $(X, T, \leq)$ . Let  $f : (A, T/A) \rightarrow [0, 1](I)$  be an increasing  $F_{\sigma}$  IF strongly pre semi continuous function. Then  $f$  has an increasing  $F_{\sigma}$  IF strongly pre semi continuous extension over  $(X, T, \leq)$ .

**Proof:** Let  $g, h : X \rightarrow [0, 1](I)$  be such that  $g = f = h$  on  $A$ , and  $g(x) = \langle 0 \rangle, h(x) = \langle 1 \rangle$  if  $x \notin A$ .

We now have,

$$R_t g = \begin{cases} B_t \cap \chi_A & \text{if } t \geq 0, \\ 1 & \text{if } t < 0, \end{cases}$$

Where  $B_t$  is increasing IF pre semi open  $F_{\sigma}$  set such that  $B_t/A = R_t f$  and

$$L_t h = \begin{cases} A_t \cap \chi_A & \text{if } t \leq 1, \\ 1 & \text{if } t > 1, \end{cases}$$

where  $A_t$  is increasing IF pre semi open  $F_{\sigma}$  such that  $A_t/A = L_t f$ . Thus,  $g$  is lower  $F_{\sigma}$  IF pre semi continuous,  $h$  is upper  $F_{\sigma}$  IF pre semi continuous and  $g \subseteq h$ . By Proposition 4.2, there is an increasing  $F_{\sigma}$  IF strongly pre semi continuous function  $F : (X, T, \leq) \rightarrow [0, 1](I)$  such that  $g \subseteq F \subseteq h$ ; hence  $F \equiv f$  on  $A$ .

### ACKNOWLEDGEMENT:

The authors express their sincere thanks to the referee for his valuable comments regarding the improvement of the paper.

### REFERENCES:

- [1] D. Amsaveni, M. K. Uma and E. Roja, Intuitionistic Fuzzy Pre Semi Closed Sets, *International Journal of Mathematical Sciences and applications*. (Accepted)
- [2] K. Atanassov, Intuitionistic fuzzy sets, in: V. Sgurev, Ed., VII ITKR's Session, Sofia (une 1983 Central Sci. and Techn. Library, Bulg. Academy of sciences, 1984).
- [3] K. Atanssov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 20 (1986), 87-96.

- [4] K Atanassov, Review and new results on intuitionistic fuzzy sets, Preprint IM-MFAIS 1-88, Sofia, 1988.
- [5] K. Atanassov and S. Stoeva, Intuitionistic fuzzy sets, in: *Polish symp. On Internal & Fuzzy Mathematics, Poznan* (August 1983), 23-26.
- [6] K. Atanassov and S. Stoeva, Intuitionistic L-fuzzy sets, in: R. Trappl, Ed., *Cybernetics and system Research*, Vol. 2 (Elsevier, Amsterdam, 1984), 539-540.
- [7] A.S. Bin Shahna, On fuzzy strong semi continuity and fuzzy pre - continuity, *Fuzzy Sets and Systems*, 44 (1991), 303-308.
- [8] D. Coker, An introduction to intuitionistic fuzzy topological spaces, *Fuzzy Sets and Systems*, 88 (1997), 81-89.
- [9] Gurcay, H., Coker, D., and Haydar, Es. A., On fuzzy continuity in intuitionistic fuzzy topological spaces, *The J. fuzzy mathematics*, 1997, 365-378.
- [10] Joung Kon Jeon, Young Bae Jun, and Jin Han Park, Intuitionistic fuzzy alpha-continuity and intuitionistic fuzzy pre continuity, *International Journal of Mathematics and Mathematical Sciences*, 2005, 3091-3101.
- [11] Katsaras A.K., Ordered fuzzy topological spaces, *J. Math. Anal. Appl.* 84(1981), 44-58.
- [12] Necla Turanli, Dogan Coker, Fuzzy connectedness in intuitionistic fuzzy topological spaces, *Fuzzy sets and systems* 116(2000), 369-375.
- [13] G. Thangaraj and G. Balasubramanian, *On fuzzy basically disconnected spaces*, *The Journal of Fuzzy Mathematics*, 9(2001), 103-110.
- [14] Thakur, S. S. and Singh, S., On fuzzy semi-pre open sets and fuzzy semi-precontinuity, *Fuzzy sets and systems*, 1998, 383-391.
- [15] Tomasz Kubiak, L-fuzzy normal space and Tietze extension theorem, *J. Math. Anal. Appl.*, 25(1987), 141-153.
- [16] M.K. Uma, E. Roja and G. Balasubramanian, Tietze extension theorem for ordered fuzzy pre extremally disconnected spaces, *East Asian Math J.*, 24 (2008), 213-221.
- [17] Young Bae Jun and Seok- Zun Song, Intuitionistic fuzzy semi-pre open sets and Intuitionistic semi-pre continuous mappings, *jour. of Appl. Math & computing*, 2005, 467-474.
- [18] L. A. Zadeh, Fuzzy sets, *Inform and control*, 8 (1965), 338-353.

\*\*\*\*\*