

## DUAL PSUEDO-COMPLEMENTED ALMOST DISTRIBUTIVE LATTICES

G. C. Rao\* & Naveen Kumar Kakumanu<sup>1</sup>

Department of Mathematics, Andhra University, Visakhapatnam, Andhra Pradesh – 530003, India

E-mail: [gcrmaths@yahoo.co.in](mailto:gcrmaths@yahoo.co.in); [ramanawinmaths@gmail.com](mailto:ramanawinmaths@gmail.com)

(Received on: 16-12-11; Accepted on: 05-01-12)

### ABSTRACT

The concept of a dual pseudo-complemented Almost Distributive Lattice is introduced. Necessary and sufficient conditions for an Almost Distributive Lattice to become a dual pseudo-complemented Almost Distributive Lattice are derived. It is proved that a dual pseudo-complemented Almost Distributive Lattice is equationally definable. A one to one correspondence between the set of all dual pseudo-complementations on an ADL  $A$  and the set of all maximal elements of  $A$  is obtained. Also proved that the set  $A_* = \{x_* \wedge m \mid x \in A\}$  is a Boolean algebra.

**Keywords:** Almost Distributive Lattice; Maximal element; Principal ideal; Dual pseudo-complementation.

**AMS Subject Classification:** 06D15, 06D99.

### INTRODUCTION

A pseudo-complemented lattice is a lattice  $A$  such that to each  $x \in A$ , there exists an element  $x^* \in A$  such that  $x \wedge y = 0$  for any  $y \in A$  if and only if  $y \leq x^*$ . In [6], Swamy, U.M., Rao, G.C. and Nanaji Rao, G. introduced the concept of Pseudo-Complementation on Almost Distributive Lattice and studied its properties. They observed that an Almost Distributive Lattice (ADL)  $A$  can have several pseudo-complementations and they discussed the relation between the maximal elements and the pseudo-complementations on  $A$ . Unlike in lattices, the dual of an ADL is not an ADL, in general. For this reason, in this paper, the concept of a dual pseudo-complementation in an ADL is introduced and important properties of a dual pseudo-complementation in an ADL are derived. It is proved that dual pseudo-complementation on an ADL is equationally definable. A number of characterizations for an ADL to become a dual pseudo-complemented ADL are obtained.

### 1. PRELIMINARIES

In this section, we give the necessary definitions and important properties of an ADL taken from [5]

**Definition 1.1:** [5] An algebra  $(A, \vee, \wedge, 0)$  of type  $(2, 2, 0)$  is called an Almost Distributive Lattice (ADL) if it satisfies the following axioms:

- (i)  $x \vee 0 = x$
- (ii)  $0 \wedge x = 0$
- (iii)  $(x \vee y) \wedge z = (x \wedge z) \vee (y \wedge z)$
- (iv)  $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
- (v)  $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$
- (vi)  $(x \vee y) \wedge y = y$  for all  $x, y, z \in A$ .

The binary relation  $\leq$  defined on an ADL  $A$  by  $x \leq y$  if and only if  $x \wedge y = x$  or equivalently  $x \vee y = y$ , is a partial ordering on  $A$ . A non-empty subset  $I$  of an ADL  $A$  is called an ideal of  $A$  if  $x \vee y \in I$  and  $x \wedge a \in I$  for any  $x, y \in I$  and  $a \in A$ . The principal ideal of  $A$  generated by  $x$  is denoted by  $[x]$ . The set  $PI(A)$  of all

\*Corresponding author: G. C. Rao \*, \*E-mail: [gcrmaths@yahoo.co.in](mailto:gcrmaths@yahoo.co.in)

principal ideals of  $A$  forms a distributive lattice under the operations  $\vee, \wedge$  defined by  $(x] \vee (y] = (x \vee y]$  and  $(x] \wedge (y] = (x \wedge y]$  in which  $(0]$  is the least element. If  $A$  has a maximal element  $m$ , then  $(m]$  is the greatest element of  $PI(A)$ .

**Theorem 1.2:** [5] Let  $A$  be an ADL and  $x, y \in A$ . Then the following are equivalent:

- (i)  $(x] \subseteq (y]$
- (ii)  $y \wedge x = x$
- (iii)  $y \vee x = y$
- (iv)  $[y) \subseteq [x)$

For other properties of an ADL, we refer to [5].

## 2. DUAL PSEUDO-COMPLEMENTATION ON ADLS

We begin with the following definition of a dually pseudo-complementation in an ADL.

**Definition 2.1:** Let  $(A, V, \Lambda, 0)$  be an ADL. Then a unary operation  $*$  on  $A$  is called a dual pseudo-complementation on  $A$  if, for any  $x, y \in A$ , it satisfies the following conditions:

**d<sub>1</sub>:** if  $x \vee y = m$ , then  $(x_* \vee y) \wedge m = y \wedge m$ .

**d<sub>2</sub>:**  $x \vee x_*$  is a maximal element of  $A$ .

**d<sub>3</sub>:**  $(x \wedge y)_* = x_* \vee y_*$ .

An ADL  $A$  with a dual pseudo-complementation is called a dually Pseudo-Complemented Almost Distributive Lattice (or simply dual PCADL). Here afterwards,  $A$  stands for a dual PCADL  $(A, V, \Lambda, *, 0, m)$  with a maximal element  $m$ .

In the following theorem, some important fundamental properties of  $*$  which will be frequently used are given and they can be proved directly from the definition.

**Theorem 2.2:** For any  $x, y \in A$ , we have the following:

- (i)  $m_* = 0$ .
- (ii) If  $x$  is maximal, then  $x_* = 0$ .
- (iii)  $(x \wedge m)_* = x_*$ .
- (iv)  $(x \wedge y)_* = (y \wedge x)_*$  and  $(x \vee y)_* = (y \vee x)_*$ .
- (v)  $0_* \wedge m = m$ .
- (vi)  $m_{**} \wedge m = m$ .
- (vii)  $x_{**} \wedge m \leq x \wedge m$  and  $x \wedge x_{**} = x_{**}$ .

**Theorem 2.3:** For any  $x, y \in A$ , we have the following:

- (i) If  $x \leq y$ , then  $y_* \leq x_*$  and  $x_{**} \leq y_{**}$ .
- (ii)  $x_* = x_{***}$ .
- (iii)  $x_* = 0 \Leftrightarrow x_{**} \wedge m = m$ .
- (iv)  $x \wedge m = m \Leftrightarrow x_{**} \wedge m = m$ .

**Proof:** Suppose  $x \leq y$ . Then  $x = x \wedge y$ . Thus  $x_* = (x \wedge y)_* = (y \wedge x)_* = y_* \vee x_*$  and hence  $y_* \leq x_*$ . Similarly, we get  $x_{**} \leq y_{**}$ . Thus we get (i). Since  $x_{**} = x \wedge x_{**}$ , we get  $x_{***} = (x \wedge x_{**})_* = x_* \vee x_{***} = x_*$  (by Theorem 2.2 (vii)). We get (iii), by using the facts that  $0_* \wedge m = m, m_* = 0$  and  $x_* = x_{***}$ . Now we prove (iv). Now if  $x \wedge m = m$ , then  $0 = m_* = (x \wedge m)_* = x_*$  and hence  $x_{**} \wedge m = m$ . Converse follows from Theorem 2.2 (vii).

**Corollary 2.4:** For any  $x, y \in A$ , we have the following:

- (i)  $x_{**} = (x \wedge y)_{**} \vee x_{**}$ .
- (ii)  $x_{**} = x_{**} \wedge (x \vee y)_{**}$ .

In a distributive lattice, the dual pseudo-complementation is unique (if it exists). But, in an ADL, there can be several dual pseudo-complementations. Now we prove the following.

**Lemma 2.5:** Let  $A$  be an ADL and  $*$  and  $\perp$  be dual pseudo-complementations on  $A$ . Then, for any  $x, y \in A$ , we have the following:

- (i)  $x_{\perp} \wedge x_* = x_*$  and  $x_{\perp} \vee x_* = x_{\perp}$ .
- (ii)  $x_{*\perp} = x_{\perp\perp}$ .
- (iii)  $x_* = y_* \Leftrightarrow x_{\perp} = y_{\perp}$ .
- (iv)  $x_*$  is maximal  $\Leftrightarrow x_{\perp}$  is maximal.
- (v)  $x_{\perp} = x_* \wedge 0_{\perp}$ .
- (vi)  $x_* \wedge x_{**} = 0 \Leftrightarrow x_{\perp} \wedge x_{\perp\perp} = 0$ .

**Proof:** Since  $(x \vee x_{\perp}) \wedge m = m$ , we get  $(x_* \vee x_{\perp}) \wedge m = x_{\perp} \wedge m$ . Now

$x_{\perp} \wedge x_* = x_{\perp} \wedge m \wedge x_* = (x_* \vee x_{\perp}) \wedge m \wedge x_* = (x_{\perp} \vee x_*) \wedge x_* = x_*$  and hence  $x_{\perp} \vee x_* = x_{\perp}$ . Thus we get (i).

Now  $x_{*\perp} = (x_{\perp} \wedge x_*)_{\perp} = (x_* \wedge x_{\perp})_{\perp} = x_{\perp\perp}$ . To prove (iii), suppose  $x_* = y_*$ .

Then  $x_{\perp} = x_{\perp\perp\perp} = x_{*\perp\perp} = y_{*\perp\perp} = y_{\perp\perp\perp} = y_{\perp}$ . By symmetry, we get the converse. (iv) follows from (i). Now  $x_* \wedge 0_{\perp} = x_{\perp} \wedge x_* \wedge 0_{\perp} = x_{\perp} \wedge 0_{\perp} = x_{\perp}$  (since  $x_{\perp} \leq 0_{\perp}$ ). Thus we get (v). Suppose  $x_* \wedge x_{**} = 0$ .

Then  $x_{\perp} \wedge x_{\perp\perp} = x_{\perp} \wedge x_{*\perp} = (x_* \wedge 0_{\perp}) \wedge (x_{**} \wedge 0_{\perp}) = 0$ .

Hence  $x_{\perp} \wedge x_{\perp\perp} = 0$ . By symmetry, we get the converse.

If  $(A, \vee, \wedge)$  is a finite distributive lattice and if we define, for any  $x \in A$ ,

$x_* = \Lambda \{y \in A \mid x \vee y = 1\}$ . Then  $*$  is a dual pseudo-complementation on  $A$ . Using this, we prove the following.

**Theorem 2.6:** If  $A$  is a finite ADL, then  $A$  is a dual PCADL.

**Proof:** Let  $A$  be a finite ADL and  $m$  be a maximal element of  $A$  with respect to  $\leq$ . Then  $([0, m], \vee, \wedge)$  is a distributive lattice and hence a dual pseudo-complemented lattices. For any  $x \in A$ , define  $x_{\perp} = (x \wedge m)_*$  where  $(x \wedge m)_*$  is the dual pseudo-complement of  $x \wedge m$  in  $[0, m]$ . Let  $x, y \in A$ .

Now

$$(x \vee x_{\perp}) \wedge m = (x \wedge m) \vee (x_{\perp} \wedge m)$$

$$\begin{aligned} &= (x \wedge m) \vee ((x \wedge m)_* \wedge m) \\ &= (x \wedge m) \vee (x \wedge m)_* \\ &= m. \end{aligned}$$

Suppose  $x \vee y = m$ . Then  $(x_{\perp} \vee y) \wedge m = ((x \wedge m)_* \wedge m) \vee (y \wedge m)$

$$\begin{aligned} &= (x \wedge m)_* \vee (y \wedge m) \\ &= (y \wedge m) \text{ (since } (x \wedge m) \vee (y \wedge m) = m \text{)}. \end{aligned}$$

Finally,  $(x \wedge y)_{\perp} = ((x \wedge y) \wedge m)_* = (x \wedge m)_* \vee (y \wedge m)_* = x_{\perp} \vee y_{\perp}$ . Hence  $A$  is a dual PCADL.

**Theorem 2.7:** For any  $x, y \in A$ , the following are equivalent:

- (i)  $(x \vee y) \wedge m = m$ .
- (ii)  $(x_{**} \vee y) \wedge m = m$ .
- (iii)  $(x_{**} \vee y_{**}) \wedge m = m$ .
- (iv)  $(x \vee y_{**}) \wedge m = m$ .

**Proof:** (i)  $\Rightarrow$  (ii): Suppose  $(x \vee y) \wedge m = m$ . Then  $(x \wedge m) \vee (y \wedge m) = m$  and hence, by definition

$$((x \wedge m)_* \vee (y \wedge m)) \wedge m = y \wedge m. \text{ Thus, we get } (x_* \vee y) \wedge m = y \wedge m.$$

Now  $(x_{**} \vee y) \wedge m = (x_{**} \wedge m) \vee [(x_* \vee y) \wedge m] = (x_{**} \vee x_* \vee y) \wedge m = m$ . (ii)  $\Rightarrow$  (iii) follows from

(i)  $\Rightarrow$  (ii). Since  $x_{**} \wedge m \leq x \wedge m$ , we get (iii)  $\Rightarrow$  (iv). Similarly, since  $y_{**} \wedge m \leq y \wedge m$ , we get (iv)  $\Rightarrow$  (i).

**Theorem 2.8:** For any  $x, y \in A$ ,  $(x \vee y)_{**} = x_{**} \vee y_{**}$ .

**Proof:** From theorem 2.3 (i), we get  $x_{**} \vee y_{**} \leq (x \vee y)_{**}$ . Since for any  $x, y \in A$ ,

$$((x \vee y) \vee (x \vee y)_*) \wedge m = m, \text{ by Theorem 2.7, we get } [x_{**} \vee y_{**} \vee (x \vee y)_*] \wedge m = m.$$

Thus  $((x \vee y)_{**} \vee x_{**} \vee y_{**}) \wedge m = (x_{**} \vee y_{**}) \wedge m$  and hence  $(x \vee y)_{**} = x_{**} \vee y_{**}$ .

In the following two theorems we prove that dual PCADL is equationally definable.

**Theorem 2.9:** Let  $A$  be an ADL with a maximal element  $m$ . A unary operation  $*$  on  $A$  is a dual pseudo-complementation on  $A$  if and only if, for any  $x, y \in A$ , the following conditions hold:

- (i)  $(x \vee x_*) \wedge m = m$ .
- (ii)  $(x_{**} \vee x) \wedge m = x \wedge m$ .
- (iii)  $(x \vee y)_{**} = x_{**} \vee y_{**}$ .
- (iv)  $(m_* \vee x) \wedge m = x \wedge m$ .
- (v)  $(x \wedge y)_* = x_* \vee y_*$ .

**Proof:** Suppose  $A$  satisfies conditions (i) to (v). It is enough to prove **d**<sub>1</sub> of definition 2.1. For this, first we prove that

(a).  $(x \wedge m)_* \wedge m = (x_* \vee m_*) \wedge m = (m_* \vee x_*) \wedge m = x_* \wedge m$  (by (iv)).

(b).  $x_{***} \wedge m = (x_{**} \vee m_*) \wedge m$  (by (iv))

$$= (x_{**} \wedge m)_* \wedge m \text{ (by (v))}$$

$$\begin{aligned} &= (x_{**} \wedge x \wedge m)_* \wedge m \text{ (by (ii))} \\ &= (x_{***} \vee x_* \vee m_*) \wedge m = x_* \wedge m \text{ (by (a) and (iv)).} \end{aligned}$$

Let  $x, y \in A$  such that  $x \vee y = m$ .

$$\begin{aligned} \text{Then } y \wedge m &= (m_* \vee y) \wedge m \text{ (by (iv))} \\ &= \left\{ \left[ (x_* \vee x_{**}) \wedge m \right]_* \vee y \right\} \wedge m \text{ (by (i))} \\ &= \left[ \left( (x_* \vee x_{**}) \wedge m \right)_* \wedge m \right] \vee (y \wedge m) \\ &= \left[ (x_* \vee x_{**})_* \wedge m \right] \vee \left( (y_{**} \vee y) \wedge m \right) \text{ (by (a) and (ii))} \\ &= \left( (x \wedge x_*)_{**} \vee y_{**} \vee y \right) \wedge m \text{ (by (v))} \\ &= \left[ \left( (x \wedge x_*) \vee y \right)_{**} \vee y \right] \wedge m \text{ (by (iii))} \\ &= \left[ \left( (x \vee y) \wedge (x_* \vee y) \right)_{**} \vee y \right] \wedge m \\ &= \left[ \left( m \wedge (x_* \vee y) \right)_{**} \vee y \right] \wedge m \\ &= \left[ (x_* \vee y)_{**} \vee y \right] \wedge m \\ &= (x_{***} \vee y_{**} \vee y) \wedge m \text{ (by (iii))} \\ &= (x_* \vee y) \wedge m \text{ (by (b) and (ii)).} \end{aligned}$$

Hence  $A$  is a dual PCADL. Conversely, if  $A$  is a dual PCADL, then conditions (i) to (v) are already proved.

**Theorem 2.10:** Let  $A$  be an ADL with a maximal element  $m$ . A unary operation  $*$  on  $A$  is a dual pseudo-complementation if and only if, for any  $x, y \in A$ , it satisfies the following conditions:

- (i)  $(x_* \vee y) \wedge m = \left( (x \vee y)_* \vee y \right) \wedge m$ .
- (ii)  $(m_* \vee x) \wedge m = x \wedge m$ .
- (iii)  $m_{**} \wedge m = m$ .
- (iv)  $(x \wedge y)_* = x_* \vee y_*$ .

**Proof:** Suppose  $*$  is a dual pseudo-complementation on  $A$ . Clearly, we have (ii), (iii) and (iv). Since  $(x \vee y \vee (x \vee y)_*) \wedge m = m$ , we get  $(y \vee (x \vee y)_*) \wedge m = (x_* \vee y \vee (x \vee y)_*) \wedge m = (x_* \vee y) \wedge m$ .

Conversely, suppose that  $*$  satisfies conditions (i) to (iv). Let  $x, y \in A$  such that  $x \vee y = m$ .

$$\begin{aligned} \text{Then } (x \vee y)_* \wedge m &= \left[ (x \vee y)_* \vee y \right] \wedge m \text{ (by (i))} \\ &= (m_* \vee y) \wedge m = y \wedge m \text{ (by (ii)). Thus we get } \mathbf{d}_1 \text{ of def 2.1.} \end{aligned}$$

$$\begin{aligned} \text{Now } (x_* \vee x) \wedge m &= \left( (x_* \wedge m) \vee x \right) \wedge m \\ &= \left\{ \left[ (m_* \vee x) \wedge m \right]_* \vee x \right\} \wedge m \text{ (by (ii))} \\ &= \left( (m_* \vee x)_* \vee x \right) \wedge m \text{ (since } (x \wedge m)_* \wedge m = x_* \wedge m \text{)} \\ &= (m_{**} \vee x) \wedge m \text{ (by (i))} \\ &= m. \end{aligned}$$

Hence  $A$  is a dual PCADL.

**Theorem 2.11:** Let  $A$  be an ADL with a maximal element  $m$ . Then the following are equivalent:

- (i)  $A$  is a dual PCADL.
- (ii)  $[a, a \vee m]$  is a dual pseudo-complemented lattice for all  $a \in A$ .
- (iii)  $[0, m]$  is a dual pseudo-complemented lattice.

**Proof:** (i)  $\Rightarrow$  (ii): Suppose  $A$  is a dual PCADL and  $a \in A$ . For  $x \in A$ , define

$x_{\perp} = a \vee (x_{*} \wedge m)$ . Let  $x, y \in [a, a \vee m]$  such that  $x \vee y = a \vee m$ . Then  $(x \vee y) \wedge m = m$  and hence  $(x_{*} \vee y) \wedge m = y \wedge m$ . This implies  $(x_{*} \vee y) \wedge m_1 = y \wedge m_1 = y$  where  $m_1 = a \vee m$ . Therefore  $x_{*} \wedge m_1 \leq y$  and hence  $x_{\perp} = a \vee (x_{*} \wedge m) \leq y$ . Conversely, suppose  $x_{\perp} \leq y$ .

Then  $x \vee x_{\perp} = (x \vee x_{\perp}) \wedge m_1 = [x \vee a \vee (x_{*} \wedge m)] \wedge m_1 = m_1 = a \vee m$ . Hence  $[a, a \vee m]$  is a dual pseudo-complemented lattice. (ii)  $\Rightarrow$  (iii) is trivial. Now we show that (iii)  $\Rightarrow$  (i). Suppose  $[0, m]$  is a dual pseudo-complemented lattice under the unary operation  $\perp$ . For  $x \in A$ , define  $x_{*} = (x \wedge m)_{\perp}$ .

Then  $(x \vee x_{*}) \wedge m = [(x \wedge m) \vee (x_{*} \wedge m)] = (x \wedge m) \vee (x \wedge m)_{\perp} = m$ . Suppose  $x, y \in A$  such that  $x \vee y = m$ . Then  $(x \wedge m) \vee (y \wedge m) = m$ .

Thus  $(x \wedge m)_{\perp} \leq y \wedge m$  and hence  $(x_{*} \vee y) \wedge m = (x_{*} \wedge m) \vee (y \wedge m) = (x \wedge m)_{\perp} \vee (y \wedge m) = y \wedge m$ .

Finally,  $(x \wedge y)_{*} = (x \wedge y \wedge m)_{\perp} = [(x \wedge m) \wedge (y \wedge m)]_{\perp} = (x \wedge m)_{\perp} \vee (y \wedge m)_{\perp} = x_{*} \vee y_{*}$ . Hence  $A$  is a dual PCADL. If  $A$  is an ADL, then the set  $PI(A)$  of all principal ideals of  $A$  forms a distributive lattice [5]. Now, we prove the following.

**Theorem 2.12:** Let  $A$  be an ADL. Then  $A$  is a dual PCADL if and only if  $PI(A)$  is a dual pseudo-complemented lattice.

**Proof:** Suppose  $(A, \vee, \wedge, *, 0, m)$  is a dual PCADL. For any  $a \in A$ , define  $(x]_{+} = (x_{*}]$ . Let  $x, y \in A$  such that  $(x] \vee (y] = A$ . Then  $m = (x \vee y) \wedge m$ , so that  $y \wedge m = (x_{*} \wedge m) \vee (y \wedge m)$  and hence  $(x_{*}] \subseteq (y]$ . Also  $(x] \vee (x_{*}] = (x \vee x_{*}) \wedge m = m = A$ . Hence  $PI(A)$  is a dual pseudo-complemented lattice.

Conversely, suppose  $(PI(A), \vee, \wedge, +)$  is a dual pseudo-complemented lattice. For  $x \in A$ , define  $x_{*} = a \wedge m$  where  $(x]_{+} = (a]$ . Since  $(a] = (b]$  if and only if  $a \wedge m = b \wedge m$ , we get that  $*$  is well defined. We also get that  $(x]_{+} = (x_{*}]$ . Let  $x, y \in A$  such that  $x \vee y = m$ . Then  $(x] \vee (y] = A$  and hence  $(x]_{+} \subseteq (y]$ . Therefore  $x_{*} \wedge m \leq y \wedge m$ . Now,  $(m] = (x] \vee (x]_{+} = (x \vee x_{*}) \wedge m = m$ . Therefore  $x \vee x_{*}$  is a maximal. Finally, let  $x, y \in A$  suppose  $(x]_{+} = (a]$  and  $(y]_{+} = (b]$ .

Then  $(x \cap y]_{+} = ((x] \cap (y])_{+} = (x]_{+} \vee (y]_{+} = (x_{*}] \vee (y_{*}] = (x_{*} \vee y_{*}]$ .

Hence by definition,  $(x \wedge y)_{*} = x_{*} \vee y_{*}$ .

Thus  $A$  is a dual PCADL.

**Theorem 2.13:** Let  $N$  be the set of all maximal elements in  $A$  and  $DPC(A)$  be the set of all dual pseudo-complementations on  $A$ . For any  $n \in N$ , define  $*_n : A \rightarrow A$  by  $x_{*n} = x_{*} \wedge n$  for all  $x \in A$ . Then  $(A, \vee, \wedge, *_n, 0, n)$  is a dual PCADL and the map  $\varphi : N \rightarrow DPC(A)$  defined by  $\phi(x) = x_{*n}$  is a bijection.

**Proof:** First we prove that  $\varphi$  is well-defined. Let  $x, y \in A$  such that  $x \vee y$  is maximal.

Then  $(x_n \vee y) \wedge n = ((x_n \wedge n) \vee y) \wedge m \wedge n = y \wedge m \wedge n = y \wedge n$   
and  $(x \vee x_n) \wedge n = (x \wedge n) \vee (x_n \wedge n) = (x \vee x_n) \wedge n = m \wedge n = n$ .

Therefore  $x \vee x_n$  is maximal.

Now  $(x \wedge y)_n = (x \wedge y)_* \wedge n = (x_* \vee y_*) \wedge n = (x_* \wedge n) \vee (y_* \wedge n) = x_n \wedge y_n$ . Therefore  $*_n$  is a dual pseudo-complementation on  $A$ . Let  $n_1$  and  $n_2$  be two maximal elements such that  $*_{n_1} = *_{n_2}$ .

Then  $n_1 = 0_* \wedge n_1 = 0_{*_{n_1}} = 0_{*_{n_2}} = 0_* \wedge n_2 = n_2$ .

Finally, we prove  $\varphi$  is onto. Let  $\perp \in DPC(A)$ .

Then  $n_0 = 0_\perp$  and for any  $x \in A$ ,  $x_{*_{n_0}} = x_* \wedge 0_\perp = x_\perp$  ( by lemma 2.5 (v) ). Thus  $\varphi$  is a bijection.

Let  $A$  be an ADL. It may be recalled that  $(A, \leq)$  is a partial ordered set if we define  $x \leq y$  if and only if  $x \wedge y = x$  or equivalently  $x \vee y = y$ , for any  $x, y \in A$ . Now we prove the following.

**Theorem 2.14:**  $(A_*, \leq)$  is a Boolean algebra, where  $A_* = \{a_* \wedge m \mid a \in A\}$ .

**Proof:** For any  $x \in A$ , we have  $0 \leq x$  and hence  $x_* \leq 0_*$ . So that, for any  $x, y \in A$ ,  $(x \wedge y)_* = x_* \vee y_* = y_* \vee x_*$ , and hence  $(x_* \vee y_*) \wedge m$  is the l.u.b of  $x_* \wedge m, y_* \wedge m$  in  $(A_*, \leq)$ . Also, since  $x \wedge m \leq (x \vee y) \wedge m$  and  $y \wedge m \leq (x \vee y) \wedge m$ , we get  $x_* \geq (x \vee y)_*$  and  $y_* \geq (x \vee y)_*$ . Thus  $(x \vee y)_* \wedge m$  is lower bound of  $x_* \wedge m, y_* \wedge m$  in  $(A_*, \leq)$ . Suppose  $t_* \wedge m \in A_*$  such that  $t_* \wedge m \leq x_* \wedge m$  and  $t_* \wedge m \leq y_* \wedge m$ .

Then  $x_{**} \leq t_{**}$ ,  $y_{**} \leq t_{**}$  and hence  $x_{**} \vee y_{**} \leq t_{**}$ . Thus  $t_* \wedge m \leq (x_{**} \vee y_{**})_* \wedge m$ . If we write  $(x_* \wedge m) \bar{\wedge} (y_* \wedge m) = (x_{**} \vee y_{**})_* \wedge m$ , then  $(x_* \wedge m) \bar{\wedge} (y_* \wedge m)$  is the g.l.b of  $x_{**} \wedge m, y_{**} \wedge m$  in the poset  $(A_*, \leq)$ .

Hence  $(A_*, V, \bar{\wedge}, 0, 0_* \wedge m)$  is a bounded lattice. For any  $x \in A_*$ ,  $(x_* \vee x_{**}) \wedge m = m = 0_* \wedge m$  and  $(x_* \wedge m) \bar{\wedge} (x_{**} \wedge m) = (x_{**} \vee x_{***})_* \wedge m = 0$ . Thus  $(A_*, V, \bar{\wedge}, 0, 0_* \wedge m)$  is a complemented lattice.

Finally, we prove the distributivity. Let  $x_* \wedge m, y_* \wedge m, z_* \wedge m \in A_*$ . Then

$$\begin{aligned} ((x_* \wedge m) \bar{\wedge} (y_* \wedge m)) \vee ((x_* \wedge m) \bar{\wedge} (z_* \wedge m)) &= ((x_{**} \vee y_{**})_* \wedge m) \vee ((x_{**} \vee z_{**})_* \wedge m) \\ &= [(x_{**} \vee y_{**}) \wedge (x_{**} \vee z_{**})]_* \wedge m \\ &= [x_{***} \vee (y_{**} \wedge z_{**})]_{***} \wedge m \\ &= [x_{***} \vee (y_{**} \wedge z_{**})]_* \wedge m \\ &= [x_{**} \vee (y_{***} \vee z_{***})]_* \wedge m \\ &= [x_* \vee (y_* \vee z_*)]_* \wedge m \\ &= (x_* \wedge m) \bar{\wedge} ((y_* \wedge m) \vee (z_* \wedge m)). \end{aligned}$$

Hence  $(A_*, V, \bar{\wedge}, 0, 0_* \wedge m)$  is a Boolean algebra.

**REFERENCES:**

- [1] Frink, O.: Pseudo-complements in semi-lattice, Duke Math J. 29, 505 -514(1962).
- [2] G. Epstein and A. Horn : P-algebras, an abstraction from Post algebras, Vol. 4, Number 1, 195-206,1974, Algebra Universalis.
- [3] Gratzer, G.: Lattice Theory: First concepts and Distributive Lattices, W. H. Freeman and Company, San Fransisco, 1971.
- [4] Lee, K.B.: Equational class of distributive pseudo-complemented lattices, Cand. J. Math. 22, 881-891(1970).
- [5] Swamy, U.M. and Rao, G.C., Almost Distributive Lattices, J. Aust. Math. Soc. (Series A), Vol.31 (1981), 77-91.
- [6] Swamy, U.M., Rao, G.C. and Rao, G.N., Pseudo-complementation on Almost Distributive Lattices, Southeast Asian Bulletin of Mathematics, Vol.24(2000), 95-104.

***[<sup>1</sup>The author research is supported by U.G.C under XI Plan]***

\*\*\*\*\*