

FORMATION OF SOME SUMMATION FORMULAE INVOLVING  
HYPERGEOMETRIC FUNCTION

SALAHUDDIN

P.D.M College of Engineering,Bahadurgarh ,Haryana,India  
E-mails: sludn@yahoo.com ;vsludn@gmail.com

M.P. CHAUDHARY

American Mathematical Society , U.S.A  
E-mail: mpchaudhary\_2000@yahoo.com  
(Received on: 18-12-10; Accepted on: 29-12-10)

## ABSTRACT

The main object of this paper is to establish some summation formulae involving Gauss second summation theorem .The results derived in this paper are of general character.

**Key words and Phrases:** Contiguous relation,Recurrence relation, Gauss second summation theorem .

**A.M.S. Subject Classification (2000):** 33C60 , 33C70

## A. Introduction:

The Pochhammer's symbol is defined by

$$(\alpha, k) = (\alpha)_k = \frac{\Gamma(\alpha + k)}{\Gamma(\alpha)} = \begin{cases} \alpha(\alpha + 1)(\alpha + 2) \cdots (\alpha + k - 1); & \text{if } k = 1, 2, 3, \dots \\ 1 & ; \text{ if } k = 0 \\ k! & ; \text{ if } \alpha = 1 \end{cases} \quad (1)$$

\*Corresponding author: M. P. Chaudhary

E-mail: mpchaudhary\_2000@yahoo.com

*International Journal of Mathematical Archive- 2 (1), Jan. 2011* 124

Generalized Gaussian Hypergeometric function of one variable is defined by

$${}_A F_B \left[ \begin{matrix} a_1, a_2, \dots, a_A ; \\ b_1, b_2, \dots, b_B ; \end{matrix} z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_A)_k z^k}{(b_1)_k (b_2)_k \dots (b_B)_k k!}$$

or

$${}_A F_B \left[ \begin{matrix} (a_A) ; \\ (b_B) ; \end{matrix} z \right] \equiv {}_A F_B \left[ \begin{matrix} (a_j)_{j=1}^A ; \\ (b_j)_{j=1}^B ; \end{matrix} z \right] = \sum_{k=0}^{\infty} \frac{((a_A))_k z^k}{((b_B))_k k!} \quad (2)$$

where the parameters  $b_1, b_2, \dots, b_B$  are neither zero nor negative integers and  $A, B$  are non-negative integers.

**Contiguous Relation is defined by**

[ Andrews p.363(9.16), E. D. p.51(10), H.T. F. I p.103(32)]

$$(a-b) {}_2F_1 \left[ \begin{matrix} a, b ; \\ c ; \end{matrix} z \right] = a {}_2F_1 \left[ \begin{matrix} a+1, b ; \\ c ; \end{matrix} z \right] - b {}_2F_1 \left[ \begin{matrix} a, b+1 ; \\ c ; \end{matrix} z \right] \quad (3)$$

**Legendre's duplication formula**

$$\sqrt{\pi} \Gamma(2z) = 2^{(2z-1)} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right) \quad (4)$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} = \frac{2^{(b-1)} \Gamma\left(\frac{b}{2}\right) \Gamma\left(\frac{b+1}{2}\right)}{\Gamma(b)} \quad (5)$$

$$= \frac{2^{(a-1)} \Gamma\left(\frac{a}{2}\right) \Gamma\left(\frac{a+1}{2}\right)}{\Gamma(a)} \quad (6)$$

**Recurrence relation is defined by**

$$\Gamma(z+1) = z \Gamma(z) \quad (7)$$

Gauss second summation theorem is defined by [Prud., 491(7.3.7.5)]

$${}_2F_1 \left[ \begin{matrix} a, b ; & 1 \\ \frac{a+b+1}{2} ; & 2 \end{matrix} \right] = \frac{\Gamma(\frac{a+b+1}{2}) \Gamma(\frac{1}{2})}{\Gamma(\frac{a+1}{2}) \Gamma(\frac{b+1}{2})} \quad (8)$$

$$= \frac{2^{(b-1)} \Gamma(\frac{b}{2}) \Gamma(\frac{a+b+1}{2})}{\Gamma(b) \Gamma(\frac{a+1}{2})} \quad (9)$$

In a monograph of Prudnikov et al., a summation theorem is given in the form [Prud., p.491(7.3.7.3)]

$${}_2F_1 \left[ \begin{matrix} a, b ; & 1 \\ \frac{a+b-1}{2} ; & 2 \end{matrix} \right] = \sqrt{\pi} \left[ \frac{\Gamma(\frac{a+b+1}{2})}{\Gamma(\frac{a+1}{2}) \Gamma(\frac{b+1}{2})} + \frac{2 \Gamma(\frac{a+b-1}{2})}{\Gamma(a) \Gamma(b)} \right] \quad (10)$$

Now using Legendre's duplication formula and Recurrence relation for Gamma function, the above theorem can be written in the form

$${}_2F_1 \left[ \begin{matrix} a, b ; & 1 \\ \frac{a+b-1}{2} ; & 2 \end{matrix} \right] = \frac{2^{(b-1)} \Gamma(\frac{a+b-1}{2})}{\Gamma(b)} \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a-1}{2})} + \frac{2^{(a-b+1)} \Gamma(\frac{a}{2}) \Gamma(\frac{a+1}{2})}{\{\Gamma(a)\}^2} + \frac{\Gamma(\frac{b+2}{2})}{\Gamma(\frac{a+1}{2})} \right] \quad (11)$$

## B. MAIN RESULTS OF SUMMATION FORMULAE

$$\begin{aligned} {}_2F_1 \left[ \begin{matrix} a, b ; & 1 \\ \frac{a+b+6}{2} ; & 2 \end{matrix} \right] &= \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b)^2 \Gamma(b)} \times \\ &\times \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(-8a - 4a^2 + 4a^3 + 8b + 40ab + 20a^2b - 4b^2 - 20ab^2 - 4b^3)}{(a-b+4)(a-b+2)(a-b+1)(a-b-2)} + \right. \right. \\ &\quad \left. \left. + \frac{(-16ab + 16a^2b - 16b^2 - 16b^3)}{(a-b+2)(a-b-1)(a-b-2)(a-b-4)} \right\} - \right. \\ &\quad \left. - \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(16a^2 + 16a^3 + 16ab - 16ab^2)}{(a-b+4)(a-b+2)(a-b+1)(a-b-2)} + \right. \right. \\ &\quad \left. \left. + \frac{(-8a + 4a^2 + 4a^3 + 8b - 40ab + 20a^2b + 4b^2 - 20ab^2 - 4b^3)}{(a-b+2)(a-b-1)(a-b-2)(a-b-4)} \right\} \right] \quad (12) \end{aligned}$$

$$\begin{aligned}
 {}_2F_1 \left[ \begin{matrix} a, b ; \\ \frac{a+b+7}{2} ; \end{matrix} \frac{1}{2} \right] &= \frac{2^b \Gamma(\frac{a+b+7}{2})}{(a-b) \Gamma(b)} \times \\
 &\times \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(4a(a^2 + 5b^2 + 10ab - 4a + 3))}{(a-b+5)(a-b+3)(a-b+1)(a-b-1)(a-b-3)} + \right. \right. \\
 &\quad \left. \left. + \frac{4b(b^2 + 5a^2 + 10ab - 4b + 3)}{(a-b+3)(a-b+1)(a-b-1)(a-b-3)(a-b-5)} \right\} - \right. \\
 &\quad \left. - \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{8(5a^2 + b^2 + 10ab + 4b + 3)}{(a-b+5)(a-b+3)(a-b+1)(a-b-1)(a-b-3)} + \right. \right. \\
 &\quad \left. \left. + \frac{8(5b^2 + a^2 + 10ab + 4a + 3)}{(a-b+3)(a-b+1)(a-b-1)(a-b-3)(a-b-5)} \right\} \right] \quad (13)
 \end{aligned}$$

$$\begin{aligned}
 {}_2F_1 \left[ \begin{matrix} a, b ; \\ \frac{a+b+8}{2} ; \end{matrix} \frac{1}{2} \right] &= \frac{2^b \Gamma(\frac{a+b+8}{2})}{(a-b) \Gamma(b)} \times \\
 &\times \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(8(8a - 6a^2 + a^3 + 8b + 15a^2b + 6b^2 + 15ab^2 + b^3))}{(a-b+6)(a-b+4)(a-b+2)(a-b)(a-b-2)(a-b-4)} + \right. \right. \\
 &\quad \left. \left. + \frac{16b(8 + 2a + 3a^2 - 2b + 10ab + 3b^2)}{(a-b+4)(a-b+2)(a-b)(a-b-2)(a-b-4)(a-b-6)} \right\} - \right. \\
 &\quad \left. - \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{16a(8 - 2a + 3a^2 + 2b + 10ab + 3b^2)}{(a-b+6)(a-b+4)(a-b+2)(a-b)(a-b-2)(a-b-4)} + \right. \right. \\
 &\quad \left. \left. + \frac{8(8a + 6a^2 + a^3 + 8b + 15a^2b - 6b^2 + 15ab^2 + b^3)}{(a-b+4)(a-b+2)(a-b)(a-b-2)(a-b-4)(a-b-6)} \right\} \right] \quad (14)
 \end{aligned}$$

**C. DERIVATIONS OF SUMMATION FORMULAE (12) TO (14):**

**Derivation of (12):** Substituting  $c = \frac{a+b+6}{2}$  and  $z = \frac{1}{2}$  in equation (2), we get

$$(a-b) {}_2F_1 \left[ \begin{matrix} a, b ; \\ \frac{a+b+6}{2} ; \end{matrix} \frac{1}{2} \right] = a {}_2F_1 \left[ \begin{matrix} a+1, b ; \\ \frac{a+b+6}{2} ; \end{matrix} \frac{1}{2} \right] - b {}_2F_1 \left[ \begin{matrix} a, b+1 ; \\ \frac{a+b+6}{2} ; \end{matrix} \frac{1}{2} \right]$$

Now with the help of Gauss second summation theorem, we get

$$\begin{aligned}
 L.H.S &= a \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b+1)\Gamma(b)} \left[ \frac{4 \Gamma(\frac{b}{2})}{a \Gamma(\frac{a}{2})} \left\{ \frac{(a^2 + 3ab + a + 3b)}{(a-b+4)(a-b+2)(a-b)} + \right. \right. \\
 &+ \left. \left. \frac{(b^2 + 3ab + 2b)}{(a-b+2)(a-b)(a-b-2)} \right\} - \frac{2 \Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(6a + 2b + 8)}{(a-b+4)(a-b+2)(a-b)} + \right. \right. \\
 &\left. \left. + \frac{(2a + 6b + 4)}{(a-b+2)(a-b)(a-b-2)} \right\} \right] - \\
 &- \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b-1)\Gamma(b)} \left[ \frac{4 \Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(a^2 + 3ab + 2a)}{(a-b+2)(a-b)(a-b-2)} + \frac{(b^2 + 3ab + b + 3a)}{(a-b)(a-b-2)(a-b-4)} \right\} - \right. \\
 &\left. - \frac{2b \Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(6a + 2b + 4)}{(a-b+2)(a-b)(a-b-2)} + \frac{(2a + 6b + 8)}{(a-b)(a-b-2)(a-b-4)} \right\} \right] \\
 &= \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b+1)\Gamma(b)} \left[ \frac{4 \Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(a^2 + 3ab + a + 3b)}{(a-b+4)(a-b+2)(a-b)} + \right. \right. \\
 &+ \left. \left. \frac{(b^2 + 3ab + 2b)}{(a-b+2)(a-b)(a-b-2)} \right\} - \frac{2 \Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(6a^2 + 2ab + 8a)}{(a-b+4)(a-b+2)(a-b)} + \right. \right. \\
 &\left. \left. + \frac{(2a^2 + 6ab + 4a)}{(a-b+2)(a-b)(a-b-2)} \right\} \right] - \\
 &- \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b-1)\Gamma(b)} \left[ \frac{4 \Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(a^2 + 3ab + 2a)}{(a-b+2)(a-b)(a-b-2)} + \frac{(b^2 + 3ab + b + 3a)}{(a-b)(a-b-2)(a-b-4)} \right\} - \right. \\
 &\left. - \frac{2 \Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(6ab + 2b^2 + 4b)}{(a-b+2)(a-b)(a-b-2)} + \frac{(2ab + 6b^2 + 8b)}{(a-b)(a-b-2)(a-b-4)} \right\} \right] \\
 &= \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b+1)\Gamma(b)} \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(4a^2 + 12ab + 4a + 12b)}{(a-b+4)(a-b+2)(a-b)} + \right. \right. \\
 &+ \left. \left. \frac{(4b^2 + 12ab + 8b)}{(a-b+2)(a-b)(a-b-2)} \right\} - \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(12a^2 + 4ab + 16a)}{(a-b+4)(a-b+2)(a-b)} + \right. \right.
 \end{aligned}$$

$$\left. + \frac{(4a^2 + 12ab + 8a)}{(a-b+2)(a-b)(a-b-2)} \right\} - \\
 - \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b-1)\Gamma(b)} \left[ \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(4a^2 + 12ab + 8a)}{(a-b+2)(a-b)(a-b-2)} + \frac{(4b^2 + 12ab + 4b + 12a)}{(a-b)(a-b-2)(a-b-4)} \right\} - \right. \\
 \left. - \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(12ab + 4b^2 + 8b)}{(a-b+2)(a-b)(a-b-2)} + \frac{(4ab + 12b^2 + 16b)}{(a-b)(a-b-2)(a-b-4)} \right\} \right]$$

On simplification ,we get

$${}_2F_1 \left[ \begin{matrix} a, b ; & 1 \\ \frac{a+b+6}{2} ; & 2 \end{matrix} \right] = \frac{2^b \Gamma(\frac{a+b+6}{2})}{(a-b)^2 \Gamma(b)} \times \\
 \times \left[ \frac{\Gamma(\frac{b}{2})}{\Gamma(\frac{a}{2})} \left\{ \frac{(-8a - 4a^2 + 4a^3 + 8b + 40ab + 20a^2b - 4b^2 - 20ab^2 - 4b^3)}{(a-b+4)(a-b+2)(a-b+1)(a-b-2)} + \right. \right. \\
 \left. \left. + \frac{(-16ab + 16a^2b - 16b^2 - 16b^3)}{(a-b+2)(a-b-1)(a-b-2)(a-b-4)} \right\} - \right. \\
 \left. - \frac{\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+1}{2})} \left\{ \frac{(16a^2 + 16a^3 + 16ab - 16ab^2)}{(a-b+4)(a-b+2)(a-b+1)(a-b-2)} + \right. \right. \\
 \left. \left. + \frac{(-8a + 4a^2 + 4a^3 + 8b - 40ab + 20a^2b + 4b^2 - 20ab^2 - 4b^3)}{(a-b+2)(a-b-1)(a-b-2)(a-b-4)} \right\} \right]$$

Thus , we prove the result (12).

Similarly, we can prove the other results.

### REFERENCES

1. Arora, Asish, Singh, Rahul , Salahuddin. ; Development of a family of summation formulae of half argument using Gauss and Bailey theorems *Journal of Rajasthan Academy of Physical Sciences.*, 7(2008), 335-342.
2. Lavoie, J. L.; Some summation formulae for the series  ${}_3F_2$ , *Math. Comput.*, 49(1987), 269-274.
3. Lavoie, J. L., Grondin, F. and Rathie, A.K.; Generalizations of Watson's theorem on the sum of a  ${}_3F_2$ , *Indian J. Math.*, 34(1992), 23-32.
4. Lavoie, J. L., Grondin, F. and Rathie, A.K.; Generalizations of Whipple's theorem on the sum of a  ${}_3F_2$ , *J. Comput. Appl. Math.*, 72(1996), 293-300.
5. Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O.I.; *Integrals and Series Vol. 3: More Special Functions*. Nauka, Moscow, 1986. Translated from the Russian by G.G. Gould, Gordon and Breach Science Publishers, New York, Philadelphia, London, Paris, Montreux, Tokyo, Melbourne, 1990.
6. Rainville, E. D.; The contiguous function relations for  ${}_pF_q$  with applications to Bateman's  $J_n^{u,v}$  and Rice's  $H_n(\zeta, p, \nu)$ , *Bull. Amer. Math. Soc.*, 51(1945), 714-723.
7. Salahuddin, Chaudhary, M.P ; Development of some summation formulae using Hypergeometric function, *Global journal of Science Frontier Research*, 10(2010),36-48.
8. Salahuddin, Chaudhary, M.P ; Certain summation formulae associated to Gauss second summation theorem, *Global journal of Science Frontier Research*, 10(2010),30-35.