



DUAL STONE ALMOST DISTRIBUTIVE LATTICES

G. C. Rao* & Naveen Kumar Kakumanu¹

Department of Mathematics, Andhra University, Visakhapatnam, Andhra Pradesh, India - 530003.

E-mail: gcaomaths@yahoo.co.in; ramanawinmaths@gmail.com

(Received on: 09-01-12; Accepted on: 13-02-12)

ABSTRACT

In this paper, the dual Stonity of a dual pseudo-complemented ADL is defined and proved that it is a sub ADL. The concept of a dual Stone Almost Distributive Lattice is introduced. Several necessary and sufficient conditions for an Almost Distributive Lattice (ADL) to become a dual Stone ADL are obtained.

Keywords: Almost Distributive Lattice (ADL); Birkhoff center; Principal ideal; Dual pseudo-complementation; Dual Stonity; Dual Stone ADL.

AMS 2000 Subject Classification: 06D15; 06D99.

1. INTRODUCTION

Boolean algebra is a homomorphic image of the algebra $C(X, 2)$ of all continuous functions mappings of a compact Hausdorff and totally disconnected topological space X onto the discrete two element Boolean algebra 2 . In place of the algebra 2 , if we take a dense bounded distributive lattice D , then $C(X, D)$ and its homomorphic images represent a special class of pseudo-complemented distributive lattices, namely Stone lattices. A pseudo-complemented distributive lattice A is said to be a Stone lattice if $x^* \vee x^{**} = 0^*$ for all $x \in A$. The concept of Almost Distributive Lattice (ADL) was introduced in [5] as a common abstraction of the most of the existing ring theoretic generalization of a Boolean algebra on one hand and the class of distributive lattice on the other. Later, in [6], pseudo-complementation in an ADL was introduced and the properties of Stone ADL were studied in [7]. Unlike in lattices, the dual of an ADL is not an ADL in general. For this reason, we introduced the concept of dual pseudo-complemented ADL (Dual PCADL) in [3] and studied its properties. In this paper, we define the concept of dual Stonity is a dual PCADL and derive the inter-relation between different dual pseudo-complementation in an ADL A and observe that the dual Stonity of A under different pseudo-complementations is same. We derive necessary and sufficient conditions for an ADL become a dual Stone ADL.

2. PRELIMINARIES

In this section, we give the necessary definitions and important properties of an ADL taken from [5] for ready reference.

Definition 2.1: [5] An algebra $(A, \vee, \wedge, 0)$ of type $(2, 2, 0)$ is called an Almost Distributive Lattice (ADL) if it satisfies the following axioms:

- (i) $x \vee 0 = x$
- (ii) $0 \wedge x = 0$
- (iii) $(x \vee y) \wedge z = (x \wedge z) \vee (y \wedge z)$
- (iv) $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
- (v) $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$
- (vi) $(x \vee y) \wedge y = y$ for all $x, y, z \in A$.

Corresponding author: G. C. Rao, *E-mail: gcaomaths@yahoo.co.in

A non-empty subset I of an ADL A is called an ideal of A if $x \vee y \in I$ and $x \wedge a \in I$ for any $x, y \in I$ and $a \in A$. The principal ideal of A generated by x is denoted by $(x]$ and it is given by $(x] = \{x \wedge a \mid a \in A\}$. The set $PI(A)$ of all principal ideals of A forms a distributive lattice under the operations \vee, \wedge defined by $(x] \vee (y] = (x \vee y]$ and $(x] \wedge (y] = (x \wedge y]$ in which $(0]$ is the least element. If A has a maximal element m , then $(m]$ is the greatest element of $PI(A)$. We extended many existing concept in the class of distributive lattices to the class of ADL, through the principal ideal lattice $PI(A)$. For other properties of ADLs we refer to [5]. In [6], the concept of Pseudo-Complementation in an ADL was defined as follows.

Definition 2.2: [6] Let $(A, \vee, \wedge, 0)$ be an ADL. Then a unary operation $x \rightarrow x^*$ on A is called a pseudo-complementation on A if, for any $x, y \in A$, it satisfies the following conditions:

- (i) $x \wedge y = 0 \Rightarrow x^* \wedge y = y$.
- (ii) $x \wedge x^* = 0$.
- (iii) $(x \vee y)^* = x^* \wedge y^*$.

An ADL $(A, \vee, \wedge, 0)$ together with a pseudo-complement is called a pseudo-complementation on A . The concept of a Stone ADL is defined in [7] as follows.

Definition 2.3: [7] Let $(A, \vee, \wedge, 0)$ be an ADL with a pseudo-complement $*$. Then A is called a Stone Almost Distributive Lattice (Stone ADL) if, for any $x \in A$, $x^* \vee x^{**} = 0^*$.

For other properties of PCADL, Stone ADL we refer to [6], [7].

3. DUAL STONITY DS (A)

The concept of pseudo-complementation in an ADL A was introduced in [6] and it was observed that, the set A^* of all pseudo-complementation of elements of A forms a Boolean algebra with respect to the same partial ordering of A . A pseudo-complemented ADL $(A, \vee, \wedge, *, 0)$ is called a Stone ADL if $x^* \vee x^{**} = 0^*$ for all $x \in A$. We begin with the following.

Definition 3.1: [3] Let $(A, \vee, \wedge, 0)$ be an ADL. Then a unary operation $*$ on A is called a dual pseudo-complementation on A if, for any $x, y \in A$, it satisfies the following conditions:

- d₁:** if $x \vee y = m$, then $(x_* \vee y) \wedge m = y \wedge m$.
- d₂:** $x \vee x_*$ is a maximal element of A .
- d₃:** $(x \wedge y)_* = x_* \vee y_*$.

An ADL A with a dual pseudo-complementation is called a dual Pseudo-Complemented Almost Distributive Lattice (or simply dual PCADL). We state the important properties of dual PCADL, taken from [3] in the following theorem.

Theorem 3.2: [3] Let A be an ADL and $*$ be a dual pseudo-complementation on A . For any $x, y \in A$, we have the following:

- (i) $x_* = x_{***}$.
- (ii) $(x \vee y)_{**} = x_{**} \vee y_{**}$.
- (iii) $(x \wedge m)_* = x_*$.
- (iv) If $x \leq y$, then $y_* \leq x_*$.

Unlike in distributive lattices, an ADL can have more than one dual pseudo-complementation. The following theorem is taken from [3] will be used in this paper.

Theorem 3.3: [3] Let A be an ADL and $*$ and \perp be a dual pseudo-complementation on A . For any $x, y \in A$, we have the following:

- (i) $x_{*\perp} = x_{\perp\perp}$.
- (ii) $x_* = y_* \Leftrightarrow x_{\perp} = y_{\perp}$.
- (iii) $x_* \wedge x_{**} = 0 \Leftrightarrow x_{\perp} \wedge x_{\perp\perp} = 0$.

For other properties of a dual PCADL, we refer to [3]. The following definition of dual annihilator of a subset of A is taken from [4].

Definition 3.4: [4] Let A be an ADL with a maximal element m and S is any non-empty subset of A . Write $S^+ = \{a \in A \mid s \vee a \text{ is maximal for all } s \in S\}$. Then S^+ is called the dual annihilator of S in A and it can be verified that S^+ is a filter of A . We write s^+ for S^+ if $S = \{s\}$.

Now we prove the following theorem.

Theorem 3.5: Let A be a dual PCADL. Then, for any $x \in A, x^+ = [x_*]$.

Proof: Since, for any $x \in A, x \vee x_*$ is maximal, we get $x_* \in x^+$ and hence $[x_*] \subseteq x^+$. On the other hand, if $y \in x^+$, then $x \vee y$ is maximal and hence $(x_* \vee y) \wedge m = y \wedge m$.

Now $y \wedge x_* = y \wedge m \wedge x_* = (x_* \vee y) \wedge m \wedge x_* = x_*$ and hence $y \vee x_* = y$. Thus $y \in [x_*]$. Hence $x^+ \subseteq [x_*]$.

Corollary 3.6: Let A be an ADL with a maximal element m and $*, \perp$ be dual pseudo-complementations on A . Then, for any $x \in A, [x_*] = [x_{\perp}]$.

If $*$ is a dual pseudo-complementation on A , then it was proved in [3], that $A_* = \{x_* \wedge m \mid x \in A\}$ is a Boolean algebra under operations $\vee, \bar{\wedge}$ where for any $x_* \wedge m, y_* \wedge m \in A_*, (x_* \wedge m) \bar{\wedge} (y_* \wedge m) = (x_{**} \vee y_{**})_* \wedge m$.

Now we prove the following.

Theorem 3.7: Let A be an ADL with a dual pseudo-complementation $*$. Then the map $f : A \rightarrow A_*$ defined by $f(x) = x_{**} \wedge m$ is an epimorphism.

Proof: Let $x, y \in A$. Then $f(x \wedge y) = [x \wedge y]_{**} \wedge m$
 $= [x_* \vee y_*]_{**} \wedge m$
 $= [x_{**} \vee y_{**}]_* \wedge m$
 $= (x_{**} \wedge m) \bar{\wedge} (y_{**} \wedge m)$
 $= f(x) \bar{\wedge} f(y)$

and $f(x \vee y) = (x \vee y)_{**} \wedge m = (x_{**} \wedge m) \vee (y_{**} \wedge m) = f(x) \vee f(y)$.

Also $f(x_*) = x_{***} \wedge m = x_* \wedge m$. Hence f is an epimorphism.

Theorem 3.8: Let A be an ADL with two dual pseudo-complementations $*$ and \perp . Then the map $f : A_* \rightarrow A_{\perp}$ defined by $f(x_* \wedge m) = x_{\perp} \wedge m$, is an isomorphism of the Boolean algebras A_* and A_{\perp} .

Proof: By Theorem 3.3, the map $f : A_* \rightarrow A_{\perp}$ defined by $f(x_* \wedge m) = x_{\perp} \wedge m$ for all $x_* \wedge m \in A_*$, is a bijection. Let $x, y \in A$. Then

$$\begin{aligned} f\left((x_* \wedge m) \overline{\wedge} (y_* \wedge m)\right) &= f\left((x_{**} \vee y_{**})_* \wedge m\right) \\ &= (x_{**} \vee y_{**})_{\perp} \wedge m \\ &= (x \vee y)_{**\perp} \wedge m \\ &= (x \vee y)_{\perp\perp\perp} \wedge m \\ &= (x_{\perp\perp} \vee y_{\perp\perp})_{\perp} \wedge m \\ &= (x_{\perp} \wedge m) \overline{\wedge} (y_{\perp} \wedge m) \\ &= f(x_* \wedge m) \overline{\wedge} f(y_* \wedge m) \end{aligned}$$

$$\begin{aligned} \text{and } f\left((x_* \wedge m) \vee (y_* \wedge m)\right) &= f\left((x \wedge y)_* \wedge m\right) = (x \wedge y)_{\perp} \wedge m = (x_{\perp} \wedge m) \vee (y_{\perp} \wedge m). \\ &= f(x_* \wedge m) \vee f(y_* \wedge m) \end{aligned}$$

Definition 3.9: Let A be a dual PCADL with a maximal element m . Then the set $\{x \in A \mid x_* \wedge x_{**} = 0\}$ is called the Dual Stonity of A and it is denoted by $DS(A)$.

In view of Theorem 3.3, we get that the set $DS(A)$ is independent of the dual pseudo-complementation on A . Now we prove the following.

Theorem 3.10: Let A be a dual PCADL. Then $DS(A)$ is a sub ADL of A , closed under dual pseudo-complementation.

Proof: If $x \in DS(A)$, then $x_{**} \wedge x_{***} = x_{**} \wedge x_* = 0$ and hence $x_* \in DS(A)$. Let $x, y \in A$. Then

$$\begin{aligned} (x \wedge y)_* \wedge (x \wedge y)_{**} &= (x_* \vee y_*) \wedge (x \wedge y)_{**} \\ &= \left\{ (x_* \wedge (x \wedge y)_{**}) \vee (y_* \wedge (x \wedge y)_{**}) \right\} \\ &\leq (x_* \wedge x_{**}) \vee (y_* \wedge y_{**}) = 0 \end{aligned}$$

$$\begin{aligned} \text{and } (x \vee y)_* \wedge (x \vee y)_{**} &= (x \vee y)_* \wedge (x_{**} \vee y_{**}) \\ &= \left[(x \vee y)_* \wedge x_{**} \right] \vee \left[(x \vee y)_* \wedge y_{**} \right] \\ &\leq (x_* \wedge x_{**}) \vee (y_* \wedge y_{**}) = 0. \end{aligned}$$

Therefore $x \wedge y, x \vee y \in DS(A)$. Also, since $0_* \wedge 0_{**} = 0_* \wedge 0 = 0$, we get $0 \in DS(A)$. Hence $DS(A)$ is a sub ADL of A .

4. DUAL STONE ADL

The concept of Stone lattice was given by T.P. Speed. Later, this concept was extended to the class of ADLs in [7]. In this section, we introduce the concept of a dual Stone ADL and study its properties. We derive necessary and sufficient conditions for an Dual PCADL to become a Dual Stone ADL. We begin with the following definition.

Definition 4.1: An ADL A with a dual pseudo-complementation $*$ is called a dual Stone ADL if $DS(A) = A$, or equivalently, $x_* \wedge x_{**} = 0$ for all $x \in A$. Here afterwards A stands for an ADL $(A, \vee, \wedge, 0)$ with a dual pseudo-complementation $*$ and with a maximal element m .

Theorem 4.2: A is a dual Stone ADL if and only if $(DS(A))_* = A_*$.

Proof: Suppose $(DS(A))_* = A_*$. Let $x \in A$. Then $x_* \in A_* = (DS(A))_*$ and hence $x_{**} \wedge x_{***} = x_* \wedge x_{**} = 0$. Thus $x \in DS(A)$. Therefore A is a dual Stone ADL. Converse is trivial. \square

Theorem 4.3: The following are equivalent:

- (i) A is a dual Stone ADL.
- (ii) $x^+ \vee y^+ = A$ whenever $x, y \in A$ and $x \vee y$ is maximal.
- (iii) $x_* \wedge y_* = 0$ whenever $x, y \in A$ and $x \vee y$ is maximal.

Proof: (i) \Rightarrow (ii): suppose A is a dual Stone ADL and let $x, y \in A$ such that $x \vee y$ is maximal. Then $(x \vee y) \wedge m = m$ and hence $(x_* \vee y) \wedge m = y \wedge m$. Thus $y \wedge x_* = x_*$.

Now $(x_{**} \vee y) \wedge m = [(y \wedge x_*) \vee y] \wedge m = (x_{**} \vee y_* \vee y) \wedge m = m$. So that $x_{**} \vee y$ is maximal and hence $x_{**} \in y^+$. Since $x \vee x_*$ is maximal, we get $x_* \in x^+$. Thus $0 = x_* \wedge x_{**} \in x^+ \vee y^+$.

Hence $x^+ \vee y^+ = A$. (ii) \Rightarrow (iii): Let $x, y \in A$ such that $x \vee y$ is maximal. Then $0 = s \wedge t$ for some $s \in x^+$ and $t \in y^+$. So that $s \vee x$ is maximal and hence $s \wedge x_* = x_*$. Similarly, $t \wedge y_* = y_*$. Hence $x_* \wedge y_* = s \wedge x_* \wedge t \wedge y_* = 0$. (iii) \Rightarrow (i): If $x \in A$, then $x \vee x_*$ is maximal, and hence $x_* \wedge x_{**} = 0$. Thus A is a dual Stone ADL.

Theorem 4.4: Let A be a dual Stone ADL. Then, for any $x, y \in A$, $(x_* \vee y_*) \wedge m = x_{**} \wedge y_{**} \wedge m$.

Proof: Since $x_* \wedge m \leq (x_* \vee y_*) \wedge m$, we get $(x_* \vee y_*) \wedge m \leq x_{**} \wedge m$. Similarly, we get $(x_* \vee y_*) \wedge m \leq y_{**} \wedge m$.

Hence $(x_* \vee y_*) \wedge m \leq x_{**} \wedge y_{**} \wedge m$. On the other hand, we have

$$\begin{aligned} x_{**} \wedge y_{**} \wedge (x_* \vee y_*) &= (x_{**} \wedge y_{**} \wedge x_*) \vee (x_{**} \wedge y_{**} \wedge y_*) = 0 \\ &\Rightarrow (x_{**} \wedge y_{**} \wedge (x_* \vee y_*)) \vee (x_* \vee y_*)_* = 0 \vee (x_* \vee y_*)_* \\ &\Rightarrow [x_{**} \wedge y_{**} \vee (x_* \vee y_*)_*] \wedge [(x_* \vee y_*) \vee (x_* \vee y_*)_*] \wedge m = (x_* \vee y_*)_* \wedge m \\ &\Rightarrow (x_{**} \wedge y_{**} \vee (x_* \vee y_*)_*) \wedge m = (x_* \vee y_*)_* \wedge m. \end{aligned}$$

Hence $x_{**} \wedge y_{**} \wedge m \leq (x_* \vee y_*)_* \wedge m$. Thus we get the result.

Theorem 4.5: A is a dual Stone ADL if and only if $(x \vee y)_* \wedge m = x_* \wedge y_* \wedge m$ for all $x, y \in A$.

Proof: Suppose A is a dual Stone ADL and $x, y \in A$. Then

$$\begin{aligned} (x \vee y)_* \wedge m &= (x \vee y)_{***} \wedge m \\ &= (x_{**} \vee y_{**})_* \wedge m \\ &= x_{***} \wedge y_{***} \wedge m \text{ (by theorem 4.4)} \\ &= x_* \wedge y_* \wedge m. \end{aligned}$$

Conversely, suppose, for any $x, y \in A$, $(x \vee y)_* \wedge m = x_* \wedge y_* \wedge m$. Since $x \vee x_*$ is maximal, we get $(x \vee x_*) \wedge m = m$. Then $m_* \wedge m = ((x \vee x_*) \wedge m)_* \wedge m$

$$\begin{aligned}
 &= ((x \vee x_*)_* \vee m_*) \wedge m \\
 &= (x_* \wedge x_{**}) \wedge m.
 \end{aligned}$$

Therefore $x_* \wedge x_{**} = 0$ and hence A is a dual Stone ADL.

Theorem 4.6: Let A be a dual Stone ADL. Write $A_* = \{x_* \wedge m \mid x \in A\}$. Then $(A_*, \vee, \wedge, *, 0, 0_* \wedge m)$ is a Boolean algebra.

Proof: Clearly, A_* is a bounded distributive lattice. Let $x_* \wedge m \in A_*$. Now $x_* \wedge m \wedge x_{**} \wedge m = x_* \wedge x_{**} \wedge m = 0$ and $(x_* \wedge m) \vee (x_{**} \wedge m) = (x_* \wedge x_{**}) \wedge m = m$. Hence A_* is a Boolean algebra.

Now we prove the following.

Note 4.7: Since the dual of a distributive lattice is again a distributive lattice, we say that a distributive lattice (A, \vee, \wedge) is a dual Stone lattice if its dual (A, \wedge, \vee) is a Stone lattice. Now we prove the following.

Theorem 4.8: Let A be a dual PCADL with a maximal element m . Then the following are equivalent:

- (i) A is a dual Stone ADL.
- (ii) $[a, a \vee m]$ is a dual Stone lattice for all $a \in A$.
- (iii) $[0, m]$ is a dual Stone lattice.

Proof: (i) \Rightarrow (ii): Suppose A is a dual Stone ADL and $a \in A$. For any $x \in A$, define $x_{\perp} = a \vee (x_* \wedge m)$. Then by Theorem 3.11 [3], we get that \perp is a dual pseudo-complementation on $[a, a \vee m]$. Now for any $x \in [a, a \vee m]$, $x_{\perp} \wedge x_{\perp\perp} = [a \vee (x_* \wedge m)] \wedge [a \vee (x_{**} \wedge m)] = a \vee (x_* \wedge x_{**} \wedge m) = a \vee 0 = a$. Hence $[a, a \vee m]$ is a dual Stone lattice. (ii) \Rightarrow (iii) is trivial. (iii) \Rightarrow (i): Suppose $[0, m]$ is a dual Stone lattice under the unary operation \perp . For $x \in A$, define $x_* = (x \wedge m)_{\perp}$. Then by Theorem 3.11 [3], we get A is a dual PCADL. Now, for any $x \in A$, $x_* \wedge x_{**} = (x \wedge m)_{\perp} \wedge ((x_{\perp} \wedge m)_{\perp} \wedge m)_{\perp} = x_{\perp} \wedge (x_{\perp} \wedge m)_{\perp} = x_{\perp} \wedge x_{\perp\perp} = 0$. Hence A is a dual Stone ADL.

Finally, we conclude this paper with the following.

Theorem 4.9: Let A be an ADL. Then A is a dual Stone ADL if and only if $PI(A)$ is a dual Stone lattice.

Proof: Suppose A is a dual Stone ADL. For any $x \in A$, define $(x]_{+} = (x_*]$. Then by Theorem 3.12 [3], we get $PI(A)$ is a dual pseudo-complemented lattice. Now $(x]_{+} \cap (x]_{++} = (x_*] \cap (x_{**}] = (x_* \wedge x_{**}] = (0]$. Hence $PI(A)$ is a dual Stone lattice. Conversely, suppose $(PI(A), \vee, \wedge, +)$ is a dual Stone lattice. For $x \in A$, define $x_* = a \wedge m$ where $(x]_{+} = (a]$. Then by Theorem 3.12 [3], we get that A is a dual PCADL. Let $x \in A$. Suppose $(x]_{+} = (a]$ and $(x]_{++} = (a]_{+} = (b]$. Since $(x]_{+} \cap (x]_{++} = (0]$, we get $a \wedge b = 0$. Since $(a \wedge m] = (a]$, we get that $x_* \wedge x_{**} = a \wedge m \wedge b \wedge m = 0$. Hence A is a dual Stone ADL.

REFERENCES:

[1] Birkhoff, G.: Lattice Theory. Amer. Math. Soc. Colloq. Publ. XXV, Providence (1967), U.S.A.
 [2] G. Epstein and A. Horn : P-algebras, an abstraction from Post algebras, Vol. 4, Number 1, 195-206,1974, Algebra Universalis.

- [3] Rao, G.C. and Naveen Kumar Kakumanu: dual Pseudo-complemented Almost Distributive Lattice, Published in International Journal of Mathematical Archive Vol.-3(2), Feb.-2012.
- [4] Rao, G.C. and Ravi Kumar, S., Normal Almost Distributive Lattices, Southeast Asian Bulletin of Mathematics, Vol.32 (2008), 831-841.
- [5] Swamy, U.M. and Rao, G.C., Almost Distributive Lattices, J. Aust. Math. Soc. (Series A), Vol.31 (1981), 77-91.
- [6] Swamy,U.M., Rao,G.C. and Rao,G.N., Pseudo-complementation on Almost Distributive Lattices, Southeast Asian Bulletin of Mathematics, Vol.24(2000), 95-104.
- [7] Swamy. U.M., Rao. G.C., Nanaji Rao. G.: Stone Almost Distributive Lattices, Southeast Asian Bulletin of mathematics, 27 (2003), 513-526.

[¹The Author Research is Supported by U.G.C. Under XI Plan]
