

**SEMI-GENERALIZED CLOSED SETS
 IN INTUITIONISTIC FUZZY TOPOLOGICAL SPACES**

¹M. Thirumalaiswamy* & ²K. Ramesh

¹*Asso. Prof., Department of Mathematics, NGM College, Pollachi - 642 001, Tamil Nadu, India*

²*Asst. Prof., Department of Science and Humanities, PA College of Engineering and Technology,
 Pollachi - 642002, Tamil Nadu, India*

(Received on: 30-12-12; Revised & Accepted on: 20-02-13)

ABSTRACT

This paper is devoted to the study of intuitionistic fuzzy topological spaces. In this paper an intuitionistic fuzzy semi-generalized closed set introduced. Some of its properties are studied.

Keywords: *Intuitionistic fuzzy topology, intuitionistic fuzzy semi-generalized closed sets and intuitionistic fuzzy points.*

AMS Subject Classification (2000): 54A40, 03F55.

1. INTRODUCTION

Fuzzy set(FS), proposed by Zadeh [13] in 1965, as a framework to encounter uncertainty, vagueness and partial truth, represents a degree of membership for each member of the universe of discourse to a subset of it. Later on, fuzzy topology was introduced by Chang [3] in 1967. By adding the degree of non-membership to FS, Atanassov [1] proposed intuitionistic fuzzy set (IFS) in 1983 which looks more accurately to uncertainty quantification and provides the opportunity to precisely model the problem based on the existing knowledge and observations. In the last few years various concepts in fuzzy were extended to intuitionistic fuzzy sets. In 1997, Coker [4] introduced the concept of intuitionistic fuzzy topological space. In 2005, R. K. Saraf [8] introduced the concept of fuzzy semi-generalized closed sets in fuzzy topological space. In this paper, we introduce intuitionistic fuzzy semi-generalized closed sets in intuitionistic fuzzy topological space. We established their properties and relationships with other classes of early defined forms.

2. PRELIMINARIES

Definition 2.1: [1] Let X be a non-empty fixed set. An intuitionistic fuzzy set (IFS in short) A in X is an object having the form $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle / x \in X \}$ where the functions $\mu_A: X \rightarrow [0,1]$ and $\nu_A: X \rightarrow [0,1]$ denote the degree of membership (namely $\mu_A(x)$) and the degree of non-membership (namely $\nu_A(x)$) of each element $x \in X$ to the set A , respectively, and $0 \leq \mu_A(x) + \nu_A(x) \leq 1$ for each $x \in X$. Denote by $\text{IFS}(X)$, the set of all intuitionistic fuzzy sets in X .

Definition 2.2: [1] Let A and B be IFSs of the form $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle / x \in X \}$ and $B = \{ \langle x, \mu_B(x), \nu_B(x) \rangle / x \in X \}$. Then

- (i) $A \subseteq B$ if and only if $\mu_A(x) \leq \mu_B(x)$ and $\nu_A(x) \geq \nu_B(x)$ for all $x \in X$,
- (ii) $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$,
- (iii) $A^c = \{ \langle x, \nu_A(x), \mu_A(x) \rangle / x \in X \}$,
- (iv) $A \cap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \nu_A(x) \vee \nu_B(x) \rangle / x \in X \}$,
- (v) $A \cup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \nu_A(x) \wedge \nu_B(x) \rangle / x \in X \}$.

For the sake of simplicity, we shall use the notation $A = \langle x, \mu_A, \nu_A \rangle$ instead of $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle / x \in X \}$.

Definition 2.3: [4] An intuitionistic fuzzy topology (IFT in short) on X is a family τ of IFSs in X satisfying the following axioms:

- (i) $0, 1 \in \tau$,
- (ii) $G_1 \cap G_2 \in \tau$, for any $G_1, G_2 \in \tau$,
- (iii) $\cup G_i \in \tau$ for any family $\{G_i / i \in J\} \subseteq \tau$.

Corresponding author: ¹M. Thirumalaiswamy*

¹*Asso. Prof., Department of Mathematics, NGM College, Pollachi - 642 001, Tamil Nadu, India*

In this case the pair (X, τ) is called an *intuitionistic fuzzy topological space* (IFTS in short) and any IFS in τ is known as an intuitionistic fuzzy open set (IFOS in short) in X . The complement A^c of an IFOS A in an IFTS (X, τ) is called an intuitionistic fuzzy closed set (IFCS in short) in X .

Definition 2.4: [4] Let (X, τ) be an IFTS and $A = \langle x, \mu_A, \nu_A \rangle$ be an IFS in X . Then the intuitionistic fuzzy interior and an intuitionistic fuzzy closure are defined by
 $\text{int}(A) = \cup \{ G / G \text{ is an IFOS in } X \text{ and } G \subseteq A \}$,
 $\text{cl}(A) = \cap \{ K / K \text{ is an IFCS in } X \text{ and } A \subseteq K \}$.

Definition 2.5: [5] An IFS A of an IFTS (X, τ) is an
 (i) *intuitionistic fuzzy semiclosed set* (IFSCS in short) if $\text{int}(\text{cl}(A)) \subseteq A$,
 (ii) *intuitionistic fuzzy semiopen set* (IFSOS in short) if $A \subseteq \text{cl}(\text{int}(A))$.

Definition 2.6: [5] An IFS A of an IFTS (X, τ) is an
 (i) *intuitionistic fuzzy preclosed set* (IFPCS in short) if $\text{cl}(\text{int}(A)) \subseteq A$,
 (ii) *intuitionistic fuzzy preopen set* (IFPOS in short) if $A \subseteq \text{int}(\text{cl}(A))$.

Note that every IFOS in (X, τ) is an IFPOS in X .

Definition 2.7: [5] An IFS A of an IFTS (X, τ) is an intuitionistic fuzzy semipre closed set (IFSPCS for short) if there exists an IFPCS B such that $\text{int}(B) \subseteq A \subseteq B$.

Definition 2.8: [5] An IFS A of an IFTS (X, τ) is an
 (i) *intuitionistic fuzzy α -closed set* (IF α CS in short) if $\text{cl}(\text{int}(\text{cl}(A))) \subseteq A$,
 (ii) *intuitionistic fuzzy α -open set* (IF α OS in short) if $A \subseteq \text{int}(\text{cl}(\text{int}(A)))$,
 (iii) *intuitionistic fuzzy regular closed set* (IFRCS in short) if $A = \text{cl}(\text{int}(A))$,
 (iv) *intuitionistic fuzzy regular open set* (IFROS in short) if $A = \text{int}(\text{cl}(A))$,
 (v) *intuitionistic fuzzy β -closed set* (IF β CS in short) if $\text{int}(\text{cl}(\text{int}(A))) \subseteq A$,
 (vi) *intuitionistic fuzzy β -open set* (IF β OS in short) if $A \subseteq \text{cl}(\text{int}(\text{cl}(A)))$.

Definition 2.9: [12] An IFS A of an IFTS (X, τ) is an
 (i) *intuitionistic fuzzy semipre closed set* (IFSPCS for short) if there exists an IFPCS B such that $\text{int}(B) \subseteq A \subseteq B$,
 (ii) *intuitionistic fuzzy semipre open set* (IFSPOS for short) if there exists an IFPOS B such that $B \subseteq A \subseteq \text{cl}(B)$.

Definition 2.10: [11] An IFS A of an IFTS (X, τ) is called an intuitionistic fuzzy w -closed set (IFWCS in short) if $\text{cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is an IFSOS in (X, τ) . An IFS A of an IFTS (X, τ) is called an intuitionistic fuzzy w -open set (IFWOS in short) if A^c is an IFWCS in (X, τ) .

Definition 2.11: [6] An intuitionistic fuzzy set A of an intuitionistic fuzzy topological space (X, τ) is called an
 (i) *intuitionistic fuzzy semi generalized closed set* (IFSGCS in short) if $\text{scl}(A) \subseteq O$ whenever $A \subseteq O$ and O is an IFOS in (X, τ) ,
 (ii) *intuitionistic fuzzy semi generalized open set* (IFSGOS in short) if its complement A^c is an IFSGCS in (X, τ) .

Definition 2.12: [7] An IFS A in an IFTS (X, τ) is said to be an intuitionistic fuzzy generalized semipre closed set (IFGSPCS) if $\text{spcl}(A) \subseteq U$ whenever $A \subseteq U$ and U is an IFSOS in (X, τ) . An IFS A of an IFTS (X, τ) is called an intuitionistic fuzzy generalized semipre open set (IFGSPOS in short) if A^c is an IFGSPCS in (X, τ) .

Definition 2.13: [5] Two IFSs A and B are said to be q -coincident ($A_q B$ in short) if and only if there exists an element $x \in X$ such that $\mu_A(x) > \nu_B(x)$ or $\nu_A(x) < \mu_B(x)$.

Definition 2.14: [5] Two IFSs are said to be not q -coincident ($A_q^c B$ in short) if and only if $A \subseteq B^c$.

Definition 2.15: [9] Let $\alpha, \beta \in [0, 1]$ and $\alpha + \beta \leq 1$. An intuitionistic fuzzy point (IFP for short) $x_{(\alpha, \beta)}$ of X is an IFS of X defined by

$$x_{(\alpha, \beta)}(y) = \begin{cases} (\alpha, \beta) & \text{if } y = x \\ (0, 1) & \text{if } y \neq x \end{cases}$$

Definition 2.16: [7] An IFS A in (X, τ) is an IFQ-set if $\text{int}(\text{cl}(A)) = \text{cl}(\text{int}(A))$.

Result 2.17: [2] For an IFS A in an IFTS (X, τ) , we have $\text{spcl}(A) \supseteq A \cup (\text{int}(\text{cl}(\text{int}(A))))$.

Result 2.18: [10] Every IFOS (IFCS) in an IFTS (X, τ) is an IFSOS (IFSCS).

3. INTUITIONISTIC FUZZY SEMIPRE GENERALIZED CLOSED SET

In this section we have introduced intuitionistic fuzzy semipre generalized closed set and have studied some of its properties.

Definition 3.1: An IFS A in an IFTS (X, τ) is said to be an intuitionistic fuzzy semipre generalized closed set (IFSPGCS for short) if $\text{spcl}(A) \subseteq U$ whenever $A \subseteq U$ and U is an IFSOS in (X, τ) . The family of all IFSPGCSs of an IFTS (X, τ) is denoted by IFSPGC (X) .

For the sake of simplicity, we shall use the notation $A = \langle x, (\mu, \nu), (v, v) \rangle$ instead of $A = \langle x, (a/\mu_a, b/\nu_b), (a/\nu_a, b/\nu_b) \rangle$ in all the examples used in this paper.

Example 3.2: Let $X = \{a, b\}$ and $G = \langle x, (0.5, 0.4), (0.5, 0.6) \rangle$. Then $\tau = \{0_-, G, 1_-\}$ is an IFT on X and the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS in (X, τ) .

Theorem 3.3: Every IFCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let $A \subseteq U$ and U be an IFSOS in (X, τ) . Then $\text{spcl}(A) \subseteq \text{cl}(A) = A \subseteq U$, by hypothesis. Hence A is an IFSPGCS in (X, τ) .

Example 3.4: In Example 3.2., the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS but not an IFCS in (X, τ) .

Theorem 3.5: Every IFSPCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFSPCS and $A \subseteq U$, U be an IFSOS in (X, τ) . Then since $\text{spcl}(A) = A$ and $A \subseteq U$, we have $\text{spcl}(A) \subseteq U$. Hence A is an IFSPGCS in (X, τ) .

Example 3.6: Let $X = \{a, b\}$ and $G_1 = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle$, $G_2 = \langle x, (0.2, 0.1), (0.8, 0.9) \rangle$, $G_3 = \langle x, (0.5, 0.6), (0.5, 0.4) \rangle$ and $G_4 = \langle x, (0.6, 0.7), (0.4, 0.3) \rangle$. Then $\tau = \{0_-, G_1, G_2, G_3, G_4, 1_-\}$ is an IFT on X and the IFS $A = \langle x, (0.9, 0.6), (0.1, 0.4) \rangle$ is an IFSPGCS but not an IFSPCS in (X, τ) in (X, τ) .

Theorem 3.7: Every IF β CS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IF β CS and $A \subseteq U$, U be an IFSOS in (X, τ) . By hypothesis $\beta\text{cl}(A) = A$ and $A \subseteq U$, we have $\text{spcl}(A) \subseteq U$. Hence A is an IFSPGCS in (X, τ) .

Example 3.8: Let $X = \{a, b\}$ and $G_1 = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle$, $G_2 = \langle x, (0.2, 0.1), (0.8, 0.9) \rangle$, $G_3 = \langle x, (0.5, 0.6), (0.5, 0.4) \rangle$ and $G_4 = \langle x, (0.6, 0.7), (0.4, 0.3) \rangle$. Then $\tau = \{0_-, G_1, G_2, G_3, G_4, 1_-\}$ is an IFT on X and the IFS $A = \langle x, (0.9, 0.6), (0.1, 0.4) \rangle$ is an IFSPGCS but not an IF β CS in (X, τ) in (X, τ) .

Theorem 3.9: Every IFSCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFSCS in (X, τ) and let $A \subseteq U$, U be an IFSOS in (X, τ) . Since $\text{spcl}(A) \subseteq \text{scl}(A) = A \subseteq U$. Hence $\text{spcl}(A) \subseteq U$. Therefore A is an IFSPGCS in (X, τ) .

Example 3.10: In Example 3.2., the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS but not an IFSCS in (X, τ) .

Theorem 3.11: Every IFPCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFPCS in (X, τ) and let $A \subseteq U$, U be an IFSOS in (X, τ) . By hypothesis, $\text{pcl}(A) = A$. Since $\text{spcl}(A) \subseteq \text{pcl}(A)$. Therefore $\text{spcl}(A) \subseteq A \subseteq U$. Hence is an IFSPGCS in (X, τ) .

Example 3.12: Let $X = \{a, b\}$ and $G = \langle x, (0.3, 0.2), (0.6, 0.6) \rangle$. Then $\tau = \{0_-, G, 1_-\}$ is an IFT on X and the IFS $A = \langle x, (0.3, 0.2), (0.6, 0.6) \rangle$ is an IFSPGCS in (X, τ) but not an IFPCS in (X, τ) .

Theorem 3.13: Every IFSGCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFSGCS in (X, τ) and $A \subseteq U$, U is an IFSOS in (X, τ) . Since $\text{spcl}(A) \subseteq \text{scl}(A)$ and $\text{scl}(A) \subseteq U$, by hypothesis. Hence A is an IFSPGCS in (X, τ) .

Example 3.14: Let $X = \{a, b\}$ and let $\tau = \{0_-, G_1, G_2, 1_-\}$ where $G_1 = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle$ and $G_2 = \langle x, (0.6, 0.7), (0.4, 0.3) \rangle$. Then the IFS $A = \langle x, (0.6, 0.4), (0.4, 0.3) \rangle$ is an IFSPGCS in (X, τ) but not an IFSGCS in (X, τ) .

Theorem 3.15: Every IFWCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFWCS in (X, τ) and $A \subseteq U$, U is an IFSOS in (X, τ) . By hypothesis $\text{cl}(A) \subseteq U$. Since $\text{spcl}(A) \subseteq \text{cl}(A)$. Therefore $\text{spcl}(A) \subseteq U$. Hence A is an IFSPGCS in (X, τ) .

Example 3.16: In Example 3.2., the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS but not an IFWCS in (X, τ) .

Theorem 3.17: Every IF α CS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IF α CS in (X, τ) and let $A \subseteq U$, U is an IFSOS in (X, τ) . Since every IF α CS is an IFPCS, by Theorem 3.7., A is an IFSPGCS in (X, τ) .

Example 3.18: In Example 3.2., the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS but not an IF α CS in (X, τ) .

Theorem 3.19: Every IFRCS in (X, τ) is an IFSPGCS in (X, τ) but not conversely.

Proof: Let A be an IFRCS in (X, τ) . Since every IFRCS is an IFCS, by Theorem 3.3., A is an IFSPGCS in (X, τ) .


Example 3.20: In Example 3.4., the IFS $A = \langle x, (0.4, 0.2), (0.6, 0.7) \rangle$ is an IFSPGCS but not an IFRCS in (X, τ) .

Theorem 3.21: Every IFSPGCS in (X, τ) is an IFGSPCS in (X, τ) but not conversely.

Proof: Let A be an IFSPGCS in (X, τ) and let $A \subseteq U$, U is an IFOS in (X, τ) . Since every IFOS in (X, τ) is an IFSOS in (X, τ) and by hypothesis $\text{spcl}(A) \subseteq U$. Hence A is an IFGSPCS in (X, τ) .

Example 3.22: Let $X = \{a, b\}$ and $G = \langle x, (0.3, 0.6), (0.7, 0.4) \rangle$. Then $\tau = \{0., G, 1.\}$ is an IFT on X and the IFS $A = \langle x, (0.7, 0.6), (0.3, 0.4) \rangle$ is an IFGSPCS in (X, τ) but not an IFSPGCS in (X, τ) .

In the following diagram, we have provided the relation between various types of intuitionistic fuzzy closedness.



In this diagram by “ $A \rightarrow B$ ” we mean A implies B but not conversely.

Remark 3.23: The union of any two IFSPGCSs in (X, τ) is not an IFSPGCS in (X, τ) in general as seen from the following example.

Example 3.24: Let $X = \{a, b\}$ and let $\tau = \{0., G_1, G_2, 1.\}$ where $G_1 = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle$ and $G_2 = \langle x, (0.6, 0.7), (0.4, 0.3) \rangle$. Then the IFSs $A = \langle x, (0.6, 0.4), (0.4, 0.3) \rangle$ and $B = \langle x, (0.4, 0.8), (0.4, 0.2) \rangle$ are IFSPGCSs in (X, τ) but $A \cup B = \langle x, (0.6, 0.8), (0.4, 0.2) \rangle$ is not an IFSPGCS in (X, τ) . Let $U = \langle x, (0.6, 0.8), (0.4, 0.2) \rangle$ be an IFSOS in (X, τ) .

Since $A \cup B \subseteq U$ but $\text{spcl}(A \cup B) = 1. \notin U$.

Remark 3.25: The intersection of any two IFSPGCSs in (X, τ) is not an IFSPGCS in (X, τ) in general as seen from the following example.

Example 3.26: Let $X = \{a, b\}$ and let $\tau = \{0., G_1, G_2, G_3, G_4, 1.\}$ where $G_1 = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle$, $G_2 = \langle x, (0.2, 0.1), (0.8, 0.9) \rangle$, $G_3 = \langle x, (0.5, 0.6), (0.5, 0.4) \rangle$ and $G_4 = \langle x, (0.6, 0.7), (0.4, 0.3) \rangle$. Then the IFSs $A = \langle x, (0.9, 0.6), (0.1, 0.4) \rangle$ and $B = \langle x, (0.5, 0.9), (0.5, 0) \rangle$ are IFSPGCSs in (X, τ) but $A \cap B = \langle x, (0.5, 0.6), (0.5, 0.4) \rangle$ is not an IFSPGCS in (X, τ) . Let $U = \langle x, (0.5, 0.6), (0.5, 0.4) \rangle$ be an IFSOS in (X, τ) . Since $A \cap B \subseteq U$ but $\text{spcl}(A \cap B) = \langle x, (0.7, 0.8), (0.3, 0.2) \rangle \notin U$.

Theorem 3.27: Let (X, τ) be an IFTS. Then for every $A \in \text{IFSPGC}(X)$ and for every $B \in \text{IFS}(X)$, $A \subseteq B \subseteq \text{spcl}(A)$ implies $B \in \text{IFSPGC}(X)$.

Proof: Let $B \subseteq U$ and U be an IFSOS in (X, τ) . Then since $A \subseteq B$, $A \subseteq U$. Since A is an IFSPGCS, it follows that $\text{spcl}(A) \subseteq U$. Now $B \subseteq \text{spcl}(A)$ implies $\text{spcl}(B) \subseteq \text{spcl}(\text{spcl}(A)) = \text{spcl}(A)$. Thus, $\text{spcl}(B) \subseteq U$. This proves that $B \in \text{IFSPGC}(X)$.

Theorem 3.28: If A is an IFSOS and an IFSPGCS in (X, τ) , then A is an IFSPCS in (X, τ) .

Proof: Since $A \subseteq A$ and A is an IFSOS in (X, τ) , by hypothesis, $\text{spcl}(A) \subseteq A$. But since $A \subseteq \text{spcl}(A)$. Therefore $\text{spcl}(A) = A$. Hence A is an IFSPCS in (X, τ) .

Theorem 3.29: An IFS A of an IFTS (X, τ) is an IFSPGCS in (X, τ) if and only if $A \subseteq_q F \Rightarrow \text{spcl}(A) \subseteq_q F$ for every IFSCS F of X .

Proof: Necessity: Let F be an IFSCS in (X, τ) and $A \subseteq_q F$. Then by Definition 2.14, $A \subseteq F^c$, where F^c is an IFSOS in (X, τ) . Then $\text{spcl}(A) \subseteq F^c$, by hypothesis. Hence again by Definition 2.14, $\text{spcl}(A) \subseteq_q F$.

Sufficiency: Let U be an IFSOS in (X, τ) such that $A \subseteq U$. Then U^c is an IFSCS in (X, τ) and $A \subseteq (U^c)^c$. By hypothesis, $A \subseteq_q U^c \Rightarrow \text{spcl}(A) \subseteq_q U^c$. Hence by Definition 2.14, $\text{spcl}(A) \subseteq (U^c)^c = U$. Therefore $\text{spcl}(A) \subseteq U$. Hence A is an IFSPGCS in (X, τ) .

Theorem 3.30: Let A be an IFSPGCS in (X, τ) and $c(\alpha, \beta)$ be an IFP in X such that $c(\alpha, \beta) \subseteq_q \text{scl}(A)$. Then $\text{scl}(c(\alpha, \beta)) \subseteq_q A$.

Proof: Let A be an IFSPGCS in (X, τ) and $c(\alpha, \beta) \subseteq_q \text{scl}(A)$. If $\text{scl}(c(\alpha, \beta)) \subseteq_q A$, then by Definition 2.14, $A \subseteq (\text{scl}(c(\alpha, \beta)))^c$ where $(\text{scl}(c(\alpha, \beta)))^c$ is an IFSOS in (X, τ) . Then by hypothesis, $\text{scl}(A) \subseteq (\text{scl}(c(\alpha, \beta)))^c \subseteq (c(\alpha, \beta))^c$. Therefore by Definition 2.14, $c(\alpha, \beta) \subseteq_q \text{scl}(A)$, which is a contradiction to the hypothesis. Hence $\text{scl}(c(\alpha, \beta)) \subseteq_q A$.

Theorem 3.31: For an IFS A in (X, τ) , the following conditions are equivalent:

- (i) A is an IFOS and an IFSPGCS in (X, τ) ,
- (ii) A is an IFROS in (X, τ) .

Proof: (i) \Rightarrow (ii) Let A be an IFOS and an IFSPGCS in (X, τ) . Since every IFOS in (X, τ) is an IFSOS in (X, τ) . Then $\text{spcl}(A) \subseteq A$. This implies that $\text{int}(\text{cl}(\text{int}(A))) \subseteq A$. Since A is an IFOS, $\text{int}(A) = A$. Therefore $\text{int}(\text{cl}(A)) \subseteq A$. Since A is an IFOS, it is an IFPOS. Hence $A \subseteq \text{int}(\text{cl}(A))$. Therefore $A = \text{int}(\text{cl}(A))$. Hence A is an IFROS in (X, τ) .

(ii) \Rightarrow (i) Let A be an IFROS in (X, τ) . Therefore $A = \text{int}(\text{cl}(A))$. Since every IFROS is an IFOS, A is an IFOS and $A \subseteq A$. This implies that $\text{int}(\text{cl}(A)) \subseteq A$. Therefore A is an IFSCS. Hence by Theorem 3.9., A is an IFSPGCS in (X, τ) .

Theorem 3.32: Let $F \subseteq A \subseteq X$ where A is an IFSOS and an IFSPGCS in X . Then F is an IFSPGCS in A if and only if F is an IFSPGCS in X .

Proof: Necessity: Let U be an IFSOS in X and $F \subseteq A$. Also let F be an IFSPGCS in A . Then clearly $F \subseteq A \cap U$ and $A \cap U$ is an IFSOS in A . Hence the semipre closure of F in A , $\text{spcl}_A(F) \subseteq A \cap U$. By Theorem 3.28, A is an IFSPCS. Therefore $\text{spcl}(A) = A$ and the semi-pre closure of F in X , $\text{spcl}(F) \subseteq \text{spcl}(F) \cap \text{spcl}(A) = \text{spcl}(F) \cap A = \text{spcl}_A(F) \subseteq A \cap U \subseteq U$. That is $\text{spcl}(F) \subseteq U$ whenever $F \subseteq U$. Hence F is an IFSPGCS in X .

Sufficiency: Let V be an IFSOS in A such that $F \subseteq V$. Since A is an IFSOS in X , V is an IFSOS in X . Therefore $\text{spcl}(F) \subseteq V$, since F is an IFSPGCS in X . Thus $\text{spcl}_A(F) = \text{spcl}(F) \cap A \subseteq V \cap A \subseteq V$. Hence F is an IFSPGCS in A .

Theorem 3.33: For an IFOS A in (X, τ) , the following conditions are equivalent:

- (i) A is an IFCS in (X, τ) ,
- (ii) A is an IFSPGCS and an IFQ-set in (X, τ) .

Proof: (i) \Rightarrow (ii) Since A is an IFCS, it is an IFSPGCS in (X, τ) . Now $\text{int}(\text{cl}(A)) = \text{int}(A) = A = \text{cl}(A) = \text{cl}(\text{int}(A))$, by hypothesis. Hence A is an IFQ-set in (X, τ) .

(ii) \Rightarrow (i) Since A is an IFOS and an IFSPGCS in (X, τ) , by Theorem 3.31, A is an IFROS in (X, τ) . Therefore $A = \text{int}(\text{cl}(A)) = \text{cl}(\text{int}(A)) = \text{cl}(A)$, by hypothesis. Hence A is an IFCS in (X, τ) .

Theorem 3.34: Let (X, τ) be an IFTS. Then for every $A \in \text{IFSPC}(X)$ and for every IFS B in X , $\text{int}(A) \subseteq B \subseteq A \Rightarrow B \in \text{IFSPGC}(X)$.

Proof: Let A be an IFSPCS in X . Then by Definition 2.9, there exists an IFPCS, say C such that $\text{int}(C) \subseteq A \subseteq C$. By hypothesis, $B \subseteq A$. Therefore $B \subseteq C$. Since $\text{int}(C) \subseteq A$, $\text{int}(C) \subseteq \text{int}(A)$ and $\text{int}(C) \subseteq B$. Thus $\text{int}(C) \subseteq B \subseteq C$ and by Definition 2.9, $B \in \text{IFSPC}(X)$. Hence by Theorem 3.5, $B \in \text{IFSPGC}(X)$.

REFERENCES

- [1] K. Atanassov, *Intuitionistic fuzzy sets, Fuzzy Sets and Systems*, 20(1986), 87-96.

- [2] A. Bhattacharjee and R. N. Bhaumik, *Pre-semi closed sets and pre-semi separation axioms in intuitionistic fuzzy topological spaces*, Gen.Math.Notes, Vol.8, No.2, 2012, pp.11-17.
- [3] C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. 24 (1968), 182- 190.
- [4] D. Coker, *An introduction to intuitionistic fuzzy topological space*, *Fuzzy Sets and Systems*, 88(1997), 81-89.
- [5] H. Gurcay, Es. A. Haydar and D. Coker, *On fuzzy continuity in intuitionistic fuzzy topological spaces*, J. Fuzzy Math.5 (2) (1997), 365-378.
- [6] R. Santhi and K. Arun Prakash, *On Intuitionistic Fuzzy Semi-Generalized Closed Sets and its Applications*, Int. J. Contemp. Math. Sciences, Vol. 5, 2010, no. 34, 1677 - 1688.
- [7] R. Santhi and D. Jayanthi, *Intuitionistic fuzzy generalized semi-pre closed sets* (accepted).
- [8] R. K. Saraf, Govindappa Navalagi and Meena Khanna, *On fuzzy semi-pre generalized closed sets*, Bull. Malays. Math. Sci. Soc. (2)28(1)(2005),19-30.
- [9] Seok Jong Lee and Eun Pyo Lee, *The category of intuitionistic fuzzy topological spaces*, Bull. Korean Math. Soc. 37 (2000), No. 1, pp. 63-76.
- [10] Shyamal Debnath, *On semi open sets and semi continuous functions in Intuitionistic Fuzzy Topological Spaces*, IOSR-J. Math. Vol. 3(2012)35-38.
- [11] Thakur, S. S., Bajpai Pandey Jyoti, *Intuitionistic Fuzzy w-closed sets and intuitionistic fuzzy w-continuity*, International Journal of Contemporary Advanced Mathematics, Vol. 1, 2010, No. 1, 1–15.
- [12] Young Bae Jun and Seok- Zun Song, *Intuitionistic fuzzy semi-pre open sets and Intuitionistic fuzzy semi-pre continuous mappings*, Jour. of Appl. Math & computing, 2005, 467-474.
- [13] L. A. Zadeh, *Fuzzy sets*, Information and control, 8(1965)338-353.

Source of support: Nil, Conflict of interest: None Declared