

# ON SUPER EDGE BIMAGIC TOTAL LABELING OF $P_m \times C_n$ AND ITS RELATED GRAPHS

K. Manimekalai\*

*Department of Mathematics, Bharathi women's College (Autonomous), Affiliated by University of Madras, Chennai-600 108, Tamil Nadu, India*

K. Thirusangu

*Department of Mathematics, S.I.V.E.T College, Affiliated by University of Madras, Chennai -600 073, Tamil Nadu, India*

(Received on: 16-06-13; Revised & Accepted on: 13-07-13)

## ABSTRACT

A Graph  $G(p, q)$  is said to have edge bimagic total labeling if there exists a bijection  $f: V(G) \cup E(G) \rightarrow \{1, 2, \dots, p+q\}$  such that for each edge  $e = (u, v) \in E(G)$ ,  $f(u) + f(v) + f(e) = k_1$  or  $k_2$ , where  $k_1$  and  $k_2$  are two constants. Moreover,  $G$  is said to have super edge bimagic total labeling if  $f(V(G)) = \{1, 2, \dots, p\}$ . In this paper we prove that the super edge bimagic total labeling for generalized prism  $P_m \times C_n$ , the Mongolian Ger  $M(n, m)$ , the generalized web  $W(m, n)$ , the generalized web without centre  $W_0(m, n)$  and its related graph.

**Key Words:** edge bimagic total labeling, generalized prism, generalized web, super edge bimagic labeling and total labeling.

**AMS Subject Classification:** 05C78.

## 1. INTRODUCTION

As a standard notation, assume that  $G = G(V, E)$  is a finite, simple and undirected graph with  $p$  vertices and  $q$  edges. By a labeling we mean a one-to-one mapping that carries a set of graph elements onto a set of numbers, called labels (usually the set of integers). Edge bimagic labelings of graphs were introduced and studied by Baskar Babujee J. [1, 2]. Let  $x$  be any real number. Then  $[x]$  denotes the largest integer less than or equal to  $x$ . Terms and terminology as in Harary [4].

**Definition 1.1 [1]:** A graph  $G(p, q)$  is said to have edge bimagic total labeling with two common edge counts  $k_1$  and  $k_2$  if there exists a bijection  $f: V \cup E \rightarrow \{1, 2, \dots, p+q\}$  such that for each  $e = (u, v) \in E$ ,  $f(u) + f(v) + f(e) = k_1$  or  $k_2$ . A total edge bimagic graph is called super edge bimagic if  $f$  maps  $V$  onto  $\{1, 2, \dots, p\}$ .

That is, a bijection  $f: V(G) \cup E(G) \rightarrow \{1, 2, \dots, p+q\}$  is said to be an edge bimagic labeling with two magic constants  $k_1$  and  $k_2$ , if there exists an induced edge map  $f^*: E(G) \rightarrow \{k_1, k_2\}$  such that for every  $e = (u, v) \in E$ ,  $f^*(e) = f(u) + f(v) + f(e) = k_1$  or  $k_2$ .

**Definition 1.2:** The generalized prism  $P_m \times C_n$  is the graph with the vertex set  $V(P_m \times C_n) = \{v_{i,j} : 1 \leq i \leq m, 1 \leq j \leq n\}$  and the edge set  $E(P_m \times C_n) = \{v_{i,j} v_{i,j+1} : 1 \leq i \leq m, 1 \leq j \leq n\} \cup \{v_{i,j} v_{i+1,j} : 1 \leq i \leq m-1, 1 \leq j \leq n\}$ , where  $j$  is taken modulo  $n$  (replacing 0 by  $n$ ).

**Definition 1.3:** The generalized web  $W(m, n)$  is the graph with vertex set  $V(W(m, n)) = \{v_{i,j} : 1 \leq i \leq m+1, 1 \leq j \leq n\} \cup \{v_0\}$  and the edge set  $E(W(m, n)) = \{v_{i,j} v_{i,j+1} : 1 \leq i \leq m, 1 \leq j \leq n\} \cup \{v_{i,j} v_{i+1,j} : 1 \leq i \leq m, 1 \leq j \leq n\} \cup \{v_0 v_{1,j} : 1 \leq j \leq n\}$ , where  $j$  is taken modulo  $n$  (replacing 0 by  $n$ ). The generalized web without centre is denoted by  $W_0(m, n)$ .

**Definition 1.4:** For any integer  $n > 2$  and  $h > 1$ , the Mongolian Ger is the graph  $M(n, h)$  with the vertex set  $V(M(n, h)) = \{v_0, v_{i,j} : 1 \leq i \leq h, 1 \leq j \leq n\}$  and the edge set  $E(M(n, h)) = \{v_0 v_{1,j} : 1 \leq j \leq n\} \cup \{v_{i,j} v_{i,j+1} : 1 \leq i \leq h, 1 \leq j \leq n\} \cup \{v_{i,j} v_{i+1,j} : 1 \leq i \leq h-1, 1 \leq j \leq n\}$ , where  $j$  is taken modulo  $n$  (replacing 0 by  $n$ ).

**Corresponding author: K. Manimekalai\***

*Department of Mathematics, Bharathi women's College (Autonomous), Affiliated by University of Madras, Chennai-600 108, Tamil Nadu, India*

In this paper we study the super edge bimagic total labeling for  $P_m \times C_n$ , and some of its related graphs viz., the Mongolian Ger  $M(n, m)$ , the generalized web  $W(m, n)$ , the generalized web without centre  $W_0(m, n)$ , etc.

## 2. MAIN RESULT

**Theorem 2.1:** For all  $m \geq 2$  and odd  $n \geq 3$ , the generalized prism  $P_m \times C_n$  has super edge bimagic total labeling.

**Proof:** Let  $V$  be the vertex set and  $E$  be the edge set of the graph  $P_m \times C_n$ . Then  $|V| = mn$  and  $|E| = (2m-1)n$ . Denote the vertices of the innermost cycle of  $P_m \times C_n$  as  $v_{1,1}, v_{1,2}, \dots, v_{1,n}$  and the vertices adjacent to  $v_{1,1}, v_{1,2}, \dots, v_{1,n}$  on the second cycle as  $v_{2,1}, v_{2,2}, \dots, v_{2,n}$  respectively. Next denote the vertices adjacent to  $v_{2,1}, v_{2,2}, \dots, v_{2,n}$  on the third cycle as  $v_{3,1}, v_{3,2}, \dots, v_{3,n}$  respectively and so on. Thus the vertices adjacent to  $v_{m-1,1}, v_{m-1,2}, \dots, v_{m-1,n}$  on the  $m$ th cycle as  $v_{m,1}, v_{m,2}, \dots, v_{m,n}$  respectively.

For  $1 \leq i \leq m$ , denote

$$k(i) = \left\lfloor \frac{i-1}{n} \right\rfloor n^2 \quad (1)$$

$$\delta(i) = \begin{cases} a & ; \text{ if } i \equiv a \pmod{n} \text{ \& } a > 0 \\ n & ; \text{ if } i \equiv 0 \pmod{n} \end{cases} \quad (2)$$

We define the bijection  $f: V \cup E \rightarrow \{1, 2, \dots, (3m-1)n\}$  as follows:

Initially we assign the label to the vertices of  $P_m \times C_n$ .

For  $1 \leq i \leq m, 1 \leq j \leq n$

$$f(v_{i,j}) = \begin{cases} k(i) + (\delta(i)-1)(n+1) + j & ; 1 \leq j \leq n - \delta(i) + 1 \\ k(i) + (\delta(i)-1)(n+1) - n + j & ; n - \delta(i) + 2 \leq j \leq n \end{cases} \quad (3)$$

$$\text{Let } e = (v_{i,j}, v_{s,t}) \text{ be any edge in } P_m \times C_n \text{ and let } f'(e) = f(v_{i,j}) + f(v_{s,t}) \quad (4)$$

We denote the edges of  $P_m \times C_n$  as follows:

For  $1 \leq j \leq \frac{n-1}{2}$ , denote the edge by  $e_j^{(1)}$ , if the sum of the labels of its end vertices is equal to  $2j+1$ .

For  $0 \leq j \leq (m-1)n$ , denote the edge by  $e_j^{(2)}$ , if the sum of the labels of its end vertices is equal to  $n+1+j$ .

For  $0 \leq j \leq (m-1)n$ , denote the edge by  $e_j^{(3)}$ , if the sum of the labels of its end vertices is equal to  $mn+2+j$ .

For  $1 \leq j \leq \frac{n-3}{2}$ , denote the edge by  $e_j^{(4)}$ , if the sum of the labels of its end vertices is equal to  $2mn-n+2j+2$ .

Hence from equation (4), we have

$$\begin{aligned} f'(e_j^{(1)}) &= 2j+1 & , & \quad 1 \leq j \leq \frac{n-1}{2} \\ f'(e_j^{(2)}) &= n+1+j & , & \quad 0 \leq j \leq (m-1)n \\ f'(e_j^{(3)}) &= mn+2+j & , & \quad 0 \leq j \leq (m-1)n \\ f'(e_j^{(4)}) &= 2mn-n+2(j+1) & , & \quad 1 \leq j \leq \frac{n-3}{2} . \end{aligned} \quad (5)$$

Now, let us label the edges of  $P_m \times C_n$  as

$$\begin{aligned} f(e_j^{(1)}) &= 2mn+1-2j & , & \quad 1 \leq j \leq \frac{n-1}{2} \\ f(e_j^{(2)}) &= 2mn-n+1-j & , & \quad 0 \leq j \leq (m-1)n \\ f(e_j^{(3)}) &= 3mn-n-j & , & \quad 0 \leq j \leq (m-1)n \\ f(e_j^{(4)}) &= 2(mn-j) & , & \quad 1 \leq j \leq \frac{n-3}{2} . \end{aligned}$$

The induced map  $f^*$  on  $E$  defined by  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)})$  for any edge  $e_j^{(k)} \in E$  satisfies the conditions:

$$\begin{aligned} f^*(e_j^{(1)}) &= f'(e_j^{(1)}) + f(e_j^{(1)}) \\ &= (2j+1) + (2mn+1-2j) \\ &= 2mn+2, & 1 \leq j \leq \frac{n-1}{2} \\ f^*(e_j^{(2)}) &= f'(e_j^{(2)}) + f(e_j^{(2)}) \\ &= (n+1+j) + (2mn-n+1-j) \\ &= 2mn+2, & 0 \leq j \leq (m-1)n \\ f^*(e_j^{(3)}) &= f'(e_j^{(3)}) + f(e_j^{(3)}) \\ &= (mn+2+j) + (3mn-n-j) \\ &= 4mn-n+2, & 0 \leq j \leq (m-1)n \\ f^*(e_j^{(4)}) &= f'(e_j^{(4)}) + f(e_j^{(4)}) \\ &= 2mn-n+2(j+1) + 2(mn-j) \\ &= 4mn-n+2, & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Clearly, it is observed that for each edge  $e_j^{(k)} \in E$ ,  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) = 2mn+2$  or  $4mn-n+2$ . Since there exists two common edge counts  $k_1 = 2mn+2$  and  $k_2 = 4mn-n+2$ , the graph  $P_m \times C_n$  has edge-bimagic total labeling. Moreover  $f(V) = \{1, 2, \dots, mn\}$ ,  $f$  is a super labeling. Hence  $P_m \times C_n$  has super edge bimagic total labeling with magic counts  $2mn+2$  and  $4mn-n+2$ , where  $n$  is odd and  $m \geq 2$ .

**Theorem 2.2:** For all  $m \geq 2$  and odd  $n \geq 3$ , the Mongolian Ger  $M(n, m)$  has super edge bimagic total labeling.

**Proof:** Let  $G_1(V_1, E_1) \cong$  Mongolian Ger  $M(n, m)$ . Then the vertex set  $V_1 = V \cup \{v_0\}$  and the edge set  $E_1 = E \cup E_0$ , where  $V$  and  $E$  are as defined in the proof of the Theorem 2.1 and  $E_0 = \{v_0 v_{1,j} / 1 \leq j \leq n\}$ . Then  $|V_1| = mn+1$  and  $|E_1| = 2mn$ . We define the bijection  $f: V_1 \cup E_1 \rightarrow \{1, 2, \dots, 3mn+1\}$  as follows:

Initially we assign labels to the vertices of  $G_1$ .

$f(v_0) = mn+1$  and  $f$  on the vertices in  $V$  is as in Theorem 2.1. Then  $f(V_1) = \{1, 2, \dots, mn+1\}$ .

Let  $e = (v_{i,j}, v_{s,t})$  be any edge in  $G_1$  and let  $f'(e) = f(v_{i,j}) + f(v_{s,t})$  (6)

We denote the edges of  $G_1$  as follows:

For  $1 \leq j \leq n$ , denote each edge  $(v_0, v_{1,j}) \in E_0$  by  $e_j^{(0)}$  and the edges in  $E$  as denoted in the proof of the Theorem 2.1.

Then from equation (6), we have

$$f'(e_j^{(0)}) = mn+1+j, \quad 1 \leq j \leq n \text{ and}$$

for each edge  $e_j^{(k)} \in E$ ,  $f'(e_j^{(k)})$  is as defined in equation (5) of Theorem 2.1.

Now we assign labels to the edges of  $G_1$ .

$$\begin{aligned} f(e_j^{(0)}) &= mn+n+2-j, & 1 \leq j \leq n \\ f(e_j^{(1)}) &= 2mn+n+2-2j, & 1 \leq j \leq \frac{n-1}{2} \\ f(e_j^{(2)}) &= 2mn+2-j, & 0 \leq j \leq (m-1)n \\ f(e_j^{(3)}) &= 3mn+1-j, & 0 \leq j \leq (m-1)n \\ f(e_j^{(4)}) &= 2mn+n+1-2j, & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Thus  $f(E_1) = \{mn+2, mn+3, \dots, 3mn+1\}$ .

The induced edge map  $f^*$  on  $E_1$  defined by  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)})$  for every edge  $e_j^{(k)} \in E_1$  satisfies the conditions:

$$\begin{aligned} f^*(e_j^{(0)}) &= f'(e_j^{(0)}) + f(e_j^{(0)}) \\ &= 2mn + n + 3, \quad 1 \leq j \leq n \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(1)}) &= f'(e_j^{(1)}) + f(e_j^{(1)}) \\ &= 2mn + n + 3, \quad 1 \leq j \leq \frac{n-1}{2} \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(2)}) &= f'(e_j^{(2)}) + f(e_j^{(2)}) \\ &= 2mn + n + 3, \quad 0 \leq j \leq (m-1)n \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(3)}) &= f'(e_j^{(3)}) + f(e_j^{(3)}) \\ &= 4mn + 3, \quad 0 \leq j \leq (m-1)n \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(4)}) &= f'(e_j^{(4)}) + f(e_j^{(4)}) \\ &= 4mn + 3, \quad 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Clearly it is observed that for each edge  $e_j^{(k)} \in E_1$ ,  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) = 2mn + n + 3$  or  $4mn + 3$ . Since there exists two common edge counts  $k_1 = 2mn + n + 3$  and  $k_2 = 4mn + 3$ , the graph  $G_1$  has super edge bimagic total labeling. Hence for all  $m \geq 2$  and odd  $n \geq 3$ , the Mongolian Ger  $M(n, m)$  has super edge bimagic total labeling.

**Example 2.1:** In Figure 2.1, we give a super edge bimagic total labeling for the Mongolian Ger  $M(5, 3)$ .




Fig. 2.1 : super edge bimagic total labeling of the Mongolian Ger  $M(5, 3)$ .

In the next section, we prove that for all  $m \geq 2$  and odd  $n \geq 3$ , the generalized web without centre,  $W_0(m, n)$  is super edge bimagic. The graph  $W_0(m, n)$  is the same as the graph obtained from  $P_m \times C_n$  by attaching a pendent vertex at each vertex of the outermost cycle of  $P_m \times C_n$ . Denote the pendent vertices adjacent to  $v_{m,1}, v_{m,2}, \dots, v_{m,n}$  of the  $m$ th cycle of  $P_m \times C_n$  as  $v_{m+1,1}, v_{m+1,2}, \dots, v_{m+1,n}$  respectively.

**Theorem 2.3:** For all  $m \geq 2$  and odd  $n \geq 3$ , the graph  $W_0(m, n)$  has super edge-bimagic total labeling.

**Proof:** Let  $V'$  be the vertex set and  $E'$  be the edge set of  $W_0(m, n)$ . Then  $V' = V \cup \{v_{m+1,j} / 1 \leq j \leq n\}$  and  $E' = E \cup \{v_{m,j} v_{m+1,j} / 1 \leq j \leq n\}$ , where  $V$  and  $E$  are as defined in Theorem 2.1. Then  $|V'| = (m+1)n$  and  $|E'| = 2mn$ .

For  $1 \leq i \leq m+1$ , denote  $k(i)$  and  $\delta(i)$  as defined in equations (1) and (2) of Theorem 2.1.

We define the bijection  $f : V' \cup E' \rightarrow \{1, 2, \dots, (3m+1)n\}$  as follows:

Initially we assign labels to the vertices  $W_0(m, n)$ .

For  $1 \leq i \leq m+1$  and  $1 \leq j \leq n$  each  $v_{i,j} \in V'$ , define  $f(v_{i,j})$  as in equation (3) of Theorem 2.1.

Then  $f(V') = \{1, 2, \dots, (m+1)n\}$ .

Let  $e = (v_{i,j}, v_{s,t})$  be any edge in  $W_0(m,n)$  and let  $f'(e) = f(v_{i,j}) + f(v_{s,t})$  (7)

We denote the edges in  $E'$  as follows:

For  $1 \leq j \leq \frac{n-1}{2}$ , denote the edge by  $e_j^{(1)}$ , if the sum of the labels of its end vertices is equal to  $2j+1$ .

For  $0 \leq j \leq mn$ , denote the edge by  $e_j^{(2)}$ , if the sum of the labels of its end vertices is equal to  $n+1+j$ .

For  $0 \leq j \leq (m-1)n$ , denote the edge by  $e_j^{(3)}$ , if the sum of the labels of its end vertices is equal to  $(m+1)n + 2 + j$ .

For  $1 \leq j \leq \frac{n-3}{2}$ , denote the edge by  $e_j^{(4)}$ , if the sum of the labels of its end vertices is equal to  $2(mn + j+1)$ .

Then from equation (7), we have

$$\begin{aligned} f'(e_j^{(1)}) &= 2j + 1, & 1 \leq j \leq \frac{n-1}{2} \\ f'(e_j^{(2)}) &= n + 1 + j, & 0 \leq j \leq mn \\ f'(e_j^{(3)}) &= (m+1)n + 2 + j, & 0 \leq j \leq (m-1)n \\ f'(e_j^{(4)}) &= 2(mn + j+1), & 1 \leq j \leq \frac{n-3}{2}. \end{aligned} \quad (8)$$

Now, let us label the edges of  $W_0(m, n)$  by

$$\begin{aligned} f(e_j^{(1)}) &= 2mn + 2n - 2j + 1, & 1 \leq j \leq \frac{n-1}{2} \\ f(e_j^{(2)}) &= 2mn + n + 1 - j, & 0 \leq j \leq mn \\ f(e_j^{(3)}) &= 3mn + n - j, & 0 \leq j \leq (m-1)n \\ f(e_j^{(4)}) &= 2(mn + n - j), & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Then  $f(E') = \{mn+n+1, mn+n+2, \dots, 3mn+n\}$ .

The induced map  $f^*$  on  $E'$  defined by  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)})$  for every edge  $e_j^{(k)} \in E'$  satisfies the conditions:

$$\begin{aligned} f^*(e_j^{(1)}) &= f'(e_j^{(1)}) + f(e_j^{(1)}) \\ &= 2(mn + n + 1), & 1 \leq j \leq \frac{n-1}{2} \\ f^*(e_j^{(2)}) &= f'(e_j^{(2)}) + f(e_j^{(2)}) \\ &= 2(mn + n + 1), & 0 \leq j \leq mn \\ f^*(e_j^{(3)}) &= f'(e_j^{(3)}) + f(e_j^{(3)}) \\ &= 2(2mn + n + 1) & 0 \leq j \leq (m-1)n \\ f^*(e_j^{(4)}) &= f'(e_j^{(4)}) + f(e_j^{(4)}) \\ &= 2(2mn + n + 1), & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Clearly, it is observed that for each edge  $e_j^{(k)} \in E'$ ,  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) = 2(mn + n + 1)$  or  $2(2mn + n + 1)$ . Since there exists two common edge counts  $k_1 = 2(mn + n + 1)$  and  $k_2 = 2(2mn + n + 1)$ , the graph  $W_0(m, n)$  has super edge-bimagic total labeling, for all  $m \geq 2$  and odd  $n \geq 3$ .

**Theorem 2.4:** Let  $G_2(V_2, E_2)$  be the graph obtained from  $P_m \times C_n$  by attaching  $s$  pendent vertices at each vertex of the outermost cycle. Then the graph  $G_2$  has a super edge bimagic total labeling for all  $m \geq 2$  and odd  $n \geq 3$ .

**Proof:** The graph  $G_2(V_2, E_2)$  is the same as the graph obtained from  $W_0(m, n)$  by appending  $s-1$  pendent edges at each vertex  $v_{mj}$  ( $j = 1, 2, \dots, n$ ) of the outermost cycle of  $W_0(m, n)$ . Denote the newly attached pendent vetices at  $v_{mj}$  as  $v_j^{(1)}, v_j^{(2)}, \dots, v_j^{(s-1)}$  ( $j=1,2,\dots,n$ ). Then  $V_2 = V' \cup V_3$  where  $V_3 = \{v_j^{(l)}; j=1,2,\dots,n; l=1,2,\dots,s-1\}$  and the edge set  $E_2 = E' \cup E_3$ , where  $E_3 = \{v_{mj} v_j^{(l)} / 1 \leq j \leq n; 1 \leq l \leq s-1\}$  and  $V'$  and  $E'$  are as defined in the proof of the Theorem 2.3. Then  $|V_2| = (m+s)n$  and  $|E_2| = (2m+s-1)n$ . We define the bijection  $f: V_2 \cup E_2 \rightarrow \{1, 2, \dots, (3m+2s-1)n\}$  as follows:

Initially we assign labels to the vertices of  $G_2$ .

For  $1 \leq i \leq m+1$  and  $1 \leq j \leq n$  each  $v_{ij} \in V'$ , define  $f(v_{ij})$  as in the equation (3) of Theorem 2.1 and for  $1 \leq j \leq n; 1 \leq l \leq s-1$  each  $v_j^{(l)} \in V_3$ , define  $f(v_j^{(l)}) = f(v_{m+1,j}) + ln$ . Then  $f(V_2) = \{1, 2, \dots, (m+s)n\}$ .

Let  $e = (v_{ij}, v_{st})$  be any edge in  $G_2$  and let  $f'(e) = f(v_{ij}) + f(v_{st})$  (9)

We denote the edges of  $G_2$  as follows.

For  $1 \leq j \leq \frac{n-1}{2}$ , denote the edge by  $e_j^{(1)}$ , if the sum of the labels of its end vertices is equal to  $2j+1$ .

For  $0 \leq j \leq (m+s-1)n$ , denote the edge by  $e_j^{(2)}$ , if the sum of the labels of its end vertices is equal to  $n+1+j$ .

For  $0 \leq j \leq (m-1)n$ , denote the edge by  $e_j^{(3)}$ , if the sum of the labels of its end vertices is equal to  $mn+sn+2+j$ .

For  $1 \leq j \leq \frac{n-3}{2}$ , denote the edge by  $e_j^{(4)}$ , if the sum of the labels of its end vertices is equal to  $(2m+s-1)n+2j+2$ .

Then from equation (9), we have

$$\begin{aligned} f'(e_j^{(1)}) &= 2j+1, & 1 \leq j \leq \frac{n-1}{2} \\ f'(e_j^{(2)}) &= n+1+j, & 0 \leq j \leq (m+s-1)n \\ f'(e_j^{(3)}) &= mn+sn+2+j, & 0 \leq j \leq (m-1)n \\ f'(e_j^{(4)}) &= (2m+s-1)n+2j+2, & 1 \leq j \leq \frac{n-3}{2}. \end{aligned} \quad (10)$$

Now let us label the edges of  $G_2$  by

$$\begin{aligned} f(e_j^{(1)}) &= 2mn+2sn-2j+1, & 1 \leq j \leq \frac{n-1}{2} \\ f(e_j^{(2)}) &= 2mn+2sn-n+1-j, & 0 \leq j \leq (m+s-1)n \\ f(e_j^{(3)}) &= 3mn+2sn-n-j, & 0 \leq j \leq (m-1)n \\ f(e_j^{(4)}) &= 2(mn+sn-j), & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Then  $f(E_2) = \{mn+sn+1, mn+sn+2, \dots, (3m+2s-1)n\}$ .

The induced map  $f^*$  on  $E_2$  defined by  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)})$  for every edge  $e_j^{(k)} \in E_2$  satisfies the conditions:

$$\begin{aligned} f^*(e_j^{(1)}) &= f'(e_j^{(1)}) + f(e_j^{(1)}) \\ &= 2(mn+sn+1) = k_1, & 1 \leq j \leq \frac{n-1}{2} \\ f^*(e_j^{(2)}) &= f'(e_j^{(2)}) + f(e_j^{(2)}) \\ &= 2(mn+sn+1) = k_1, & 0 \leq j \leq (m+s-1)n \\ f^*(e_j^{(3)}) &= f'(e_j^{(3)}) + f(e_j^{(3)}) \\ &= (4m+3s-1)n+2 = k_2, & 0 \leq j \leq (m-1)n \\ f^*(e_j^{(4)}) &= f'(e_j^{(4)}) + f(e_j^{(4)}) \\ &= (4m+3s-1)n+2 = k_2, & 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$

Clearly, it is observed that for every edge  $e_j^{(k)} \in E_2$ ,  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) = 2(mn + sn + 1)$  or  $(4m + 3s - 1)n + 2$ . Since there exists two common edge counts  $k_1 = 2(mn + sn + 1)$  and  $k_2 = (4m + 3s - 1)n + 2$ , the graph  $G_2$  has super edge bimagic total labeling, for  $n$  odd and  $m \geq 2$ .

In Figure 2.3, we give a super edge bimagic total labeling for the graph  $G_2$  in Theorem 2.4 with  $m = 4$ ,  $n = 5$  and  $s = 3$ .




Figure 2.3

**Theorem 2.5:** For  $m \geq 2$  and odd  $n \geq 3$ , the generalized web  $W(m, n)$  has super edge bimagic total labeling.

**Proof:** The generalized web  $W(m, n)$  is the same as the one obtained from  $W_0(m, n)$  by joining each vertex of the innermost cycle to a new vertex  $v_0$  by an edge. Let  $V''$  be the vertex set and  $E''$  be the edge set of  $W(m, n)$ . Then  $V'' = V' \cup \{v_0\}$  and  $E'' = E' \cup E_0$ , where  $V'$  and  $E'$  are as defined in Theorem 2.3 and  $E_0 = \{v_0 v_{1,j} : 1 \leq j \leq n\}$ . Then  $|V''| = (m+1)n + 1$  and  $|E''| = (2m+1)n$ .

For  $1 \leq i \leq m+1$ , denote  $k(i)$  and  $\delta(i)$  as defined in equations (1) and (2) of Theorem 2.1.

We define the bijection  $f: V'' \cup E'' \rightarrow \{1, 2, \dots, (3mn+2)n+1\}$  as follows:

Initially we assign labels to the vertices of  $W(m, n)$ .

For  $1 \leq i \leq m+1$  and  $1 \leq j \leq n$  each vertex  $v_{i,j} \in V'$ , define  $f(v_{i,j})$  as in the equation (3) of Theorem 2.1 and  $f(v_0) = mn + n + 1$ . Then  $f(V'') = \{1, 2, \dots, mn + n + 1\}$ .

Let  $e = (v_{i,j}, v_{s,t})$  be any edge in  $W(m, n)$  and let  $f'(e) = f(v_{i,j}) + f(v_{s,t})$  (11)

We denote the edges of  $W(m, n)$  as follows:

For  $0 \leq j \leq n$ , denote each edge  $v_0 v_{1,j} \in E_0$  by  $e_j^{(0)}$  and denote the edges in  $E'$  of  $W(m, n)$  as denoted in Theorem 2.3.

Then from the equation (11), we have

$$f'(e_j^{(0)}) = mn + n + 1 + j, \quad 0 \leq j \leq n$$

and for each  $e_j^{(k)} \in E'$ ,  $f'(e_j^{(k)})$  is as defined in equation (8) of Theorem 2.3.

Now, let us label the edges of  $W(m, n)$  by

$$\begin{aligned} f(e_j^{(0)}) &= mn + 2n + 2 - j, \quad 0 \leq j \leq n \\ f(e_j^{(1)}) &= 2mn + 3n - 2j + 2, \quad 1 \leq j \leq \frac{n-1}{2} \\ f(e_j^{(2)}) &= 2mn + 2n + 2 - j, \quad 0 \leq j \leq mn \end{aligned}$$

$$f(e_j^{(3)}) = 3mn + 2n + 1 - j, 0 \leq j \leq (m-1)n$$

$$f(e_j^{(4)}) = 2mn + 3n + 1 - 2j, 1 \leq j \leq \frac{n-3}{2}.$$

Then  $f(E'') = \{mn+n+2, mn+n+3, \dots, 3mn+2n+1\}$ .

Define the induced map  $f^*$  on  $E''$  by

$$f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) \text{ for every edge } e_j^{(k)} \in E''.$$

It satisfies the conditions:

$$\begin{aligned} f^*(e_j^{(0)}) &= f'(e_j^{(0)}) + f(e_j^{(0)}) \\ &= (mn+n+1+j) + (mn+2n+2-j) \\ &= 2mn+3n+3 = k_1, \quad 0 \leq j \leq n. \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(1)}) &= f'(e_j^{(1)}) + f(e_j^{(1)}) \\ &= (2j+1) + (2mn+3n-2j+2) \\ &= 2mn+3n+3 = k_1, \quad 1 \leq j \leq \frac{n-1}{2} \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(2)}) &= f'(e_j^{(2)}) + f(e_j^{(2)}) \\ &= (n+1+j) + (2mn+2n+2-j) \\ &= 2mn+3n+3 = k_1, \quad 0 \leq j \leq mn \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(3)}) &= f'(e_j^{(3)}) + f(e_j^{(3)}) \\ &= (mn+n+2+j) + (3mn+2n+1-j) \\ &= 4mn+3n+3 = k_2, \quad 0 \leq j \leq (m-1)n \end{aligned}$$

$$\begin{aligned} f^*(e_j^{(4)}) &= f'(e_j^{(4)}) + f(e_j^{(4)}) \\ &= 2(mn+j+1) + (2mn+3n+1-2j) \\ &= 4mn+3n+3 = k_2, \quad 1 \leq j \leq \frac{n-3}{2}. \end{aligned}$$




Fig.2.4: Generalized web  $W(4,7)$



Clearly, it is observed that for each edge  $e_j^{(k)} \in E'$ ,  $f^*(e_j^{(k)}) = f'(e_j^{(k)}) + f(e_j^{(k)}) = 2mn + 3n + 3$  or  $4mn + 3n + 3$ . Since there exists two common edge counts  $k_1 = 2(mn + n + 1)$  and  $k_2 = 2(2mn + n + 1)$ , the generalized web  $W(m, n)$  has super edge bimagic total labeling for all  $m \geq 2$  and odd  $n \geq 3$ .

In Figure 2.4, we give a super edge bimagic total labeling for the generalized web  $W(4, 7)$ .

## CONCLUSION

In this paper, we have proved the super edge bimagic total labeling for generalized prism  $P_m \times C_n$ , the Mongolian Ger  $M(n, m)$ , the generalized web  $W(m, n)$ , the generalized web without centre  $W_0(m, n)$  and its related graph. Further we extend this work by examining the existence of certain other labelings for these graphs.

## REFERENCE

- [1] Baskar Babujee J., Bimagic labeling in path graphs, The Mathematics Education, Volume 38, No. 1. 2004, p. 12-16.
- [2] Baskar Babujee J., On Edge Bimagic Labeling, Journal of Combinatorics Information & System Sciences, Vol. 28, No. 1-4, 239-244 (2004).
- [3] Gallian J. A., A Dynamic Survey of Graph Labeling, Electronic Journal of Combinatorics, 17 (2010), # DS6.
- [4] Harary F. "Graph Theory", Narosa Publishing House, New Delhi, (1998).

**Source of support: Nil, Conflict of interest: None Declared**