

## A NOTE ON HARMONIC MEAN GRAPHS

S. S. Sandhya\*<sup>1</sup> and S. Somasundaram<sup>2</sup>

<sup>1</sup>Department of Mathematics, Sree Ayyappa College for Women, Chunkankadai: 629003, India.

<sup>2</sup>Department of Mathematics, M. S. University, Tirunelveli-627012, India.

(Received On: 17-01-15; Revised & Accepted On: 10-02-15)

### ABSTRACT

A Graph  $G = (V, E)$  with  $p$  vertices and  $q$  edges is said to be a Harmonic mean graph if it is possible to label the vertices  $x \in V$  with distinct labels  $f(x)$  from  $1, 2, \dots, q+1$  in such a way that when each edge  $e=uv$  is labeled with  $f(e=uv) = \left\lfloor \frac{2f(u)f(v)}{f(u)+f(v)} \right\rfloor$  (or)  $\left\lceil \frac{2f(u)f(v)}{f(u)+f(v)} \right\rceil$ , then the resulting edge labels are distinct. In this case  $f$  is called Harmonic mean labeling of  $G$ . In this paper we prove that Triple Triangular, Alternate Triple Triangular, Triple Quadrilateral and Alternate Triple Quadrilateral snakes are Harmonic mean graphs.

**Keywords:** Graph, Harmonic mean graph, Triple Triangular snake, Alternate Triple Triangular snake, Triple Quadrilateral snake, Alternate Triple Quadrilateral snake graphs.

### 1. INTRODUCTION

Throughout this paper we consider only finite, undirected and simple graphs. Let  $G = (V, E)$  be a graph with  $p$  vertices and  $q$  edges. For all terminologies and notations we follow Harary [1]. There are several types of labeling and detailed survey can be found in [2]. The concept of mean labeling has been introduced in [3] and the Harmonic mean labeling was introduced in [4]. The concept of Double Triangular snake and Double Quadrilateral snake has been proved in [5] and [6]. In this paper we prove that the Harmonic mean labeling behaviour of Triple Triangular snake, Triple Quadrilateral snake graphs. Also we wish to investigate Alternate Triple Triangular snake and Alternate Triple Quadrilateral snake graphs are Harmonic mean graphs. The following definitions are necessary for our present investigation.

**Definition 1.1:** A Graph  $G = (V, E)$  with  $p$  vertices and  $q$  edges is said to be a Harmonic mean graph if it is possible to label the vertices  $x \in V$  with distinct labels  $f(x)$  from  $1, 2, \dots, q+1$  in such a way that when each edge  $e=uv$  is labeled with  $f(e=uv) = \left\lfloor \frac{2f(u)f(v)}{f(u)+f(v)} \right\rfloor$  (or)  $\left\lceil \frac{2f(u)f(v)}{f(u)+f(v)} \right\rceil$ , then the resulting edge labels are distinct. In this case  $f$  is called Harmonic mean labeling of  $G$ .

**Definition 1.2:** A Triangular snake  $T_n$  is obtained from a path  $u_1 u_2 \dots u_n$  by joining  $u_i$  and  $u_{i+1}$  to a new vertex  $v_i$  for  $1 \leq i \leq n-1$ . That is every edge of a path is replaced by a triangles  $C_3$ .

**Definition 1.3:** An Alternate Triangular snake  $A(T_n)$  is obtained from a path  $u_1, u_2, \dots, u_n$  by joining  $u_{i+1}$  (Alternatively) to new vertex  $v_i$ . That is, every alternate edge of a path is replaced by  $C_3$ .

**Definition 1.4:** A Double Triangular snake  $D(T_n)$  consists of two Triangular snakes that have a common path.

**Definition 1.5:** An Alternate Double Triangular snake  $A[D(T_n)]$  consists of two Alternate Triangular snake that have a common path.

**Definition 1.6:** A Triple Triangular snake  $T(T_n)$  consists of three Triangular snakes that have a common path.

**Definition 1.7:** An Alternate Triple Triangular snake  $A(T(T_n))$  consists of three Alternate Triangular snakes that have a common path.

**Definition 1.8:** A Quadrilateral snake  $Q_n$  is obtained from a path  $u_1 u_2 \dots u_n$  by joining  $u_i$  and  $u_{i+1}$  to new vertices  $v_i, w_i$  respectively and then joining  $v_i$  and  $w_i$ ,  $1 \leq i \leq n-1$ . That is, every edge of a path is replaced by cycle  $C_4$ .

Corresponding Author: S. S. Sandhya\*<sup>1</sup>

**Definition 1.9:** An Alternate Quadrilateral snake  $A(Q_n)$  is obtained from a path  $u_1u_2\dots u_n$  by joining  $u_i$  and  $u_{i+1}$  (Alternatively) to new vertices  $v_i$  and  $w_i$  respectively and then joining  $v_i$  and  $w_i$ . That is, every alternate edge of path is replaced by a cycle  $C_4$ .

**Definition 1.10:** A Double Quadrilateral snake  $D(Q_n)$  consists of two Quadrilateral snakes that have a common path.

**Definition 1.11:** An Alternate Double Quadrilateral snake  $A[D(Q_n)]$  consists of two Alternate Quadrilateral snakes that have a common path.

**Definition 1.12:** An Triple Quadrilateral snakes  $T(Q_n)$  consists of three Quadrilateral snakes that have a common path.

**Definition 1.13:** An Alternate Triple Quadrilateral snake  $A[T(Q_n)]$  consists of three Alternate Quadrilateral snakes that have a common path.

## 2. MAIN RESULTS

**Theorem 2.1:** Triple Triangular snakes are Harmonic mean graphs.

**Proof:** Let  $G$  be the graph obtained from a path  $u_1u_2\dots u_n$  by joining  $u_i$  and  $u_{i+1}$  to three new vertices  $v_i$ ,  $w_i$  and  $t_i$ ,  $1 \leq i \leq n-1$ .

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_1) = 3$$

$$f(u_i) = 7i - 6, \quad 1 \leq i \leq n$$

$$f(v_1) = 1$$

$$f(v_i) = 7i - 5, \quad 1 \leq i \leq n-1$$

$$f(w_i) = 7i - 3, \quad 1 \leq i \leq n-1$$

$$f(t_i) = 7i, \quad 1 \leq i \leq n-1.$$

From the above labeling pattern, we get distinct edge labels.

Thus  $f$  provides a Harmonic mean labeling for  $G$ .

**Example 2.2:** The labeling pattern of Triple Triangular snake obtained from six vertices is given below




Figure: 1

**Theorem 2.3:** Alternate Triple Triangular snakes are Harmonic mean graphs.

**Proof:** Let  $G = A[T(T_n)]$  be the Alternate Triple Triangular snake graph and its vertices be  $v_i$ ,  $w_i$  and  $t_i$ ,  $1 \leq i \leq n-1$ . Here we consider two cases.

**Case (i):** If the triangle starts from  $u_1$  then

We need to considered two subcases.

**Subcase (i) (a):** If  $n$  is odd then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by  
 $f(u_i) = 3$

$$f(u_i) = \begin{cases} 4i-3, & \forall i = 3, 5, 7, \dots, n \\ 4i, & \forall i = 2, 4, 6, \dots, n-1. \end{cases}$$

$$f(v_1) = 1$$

$$f(v_i) = 8i-6, \quad \forall i = 2, 3, 4, \dots, \frac{n-1}{2}$$

$$f(w_i) = 8i-4, \quad \forall i = 1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(t_i) = 8i-2, \quad \forall i = 1, 2, 3, \dots, \frac{n-1}{2}$$

From the above labeling pattern, we get the edge labels are all distinct.

Thus  $f$  provide a Harmonic mean labeling for  $G$ . Harmonic mean labeling of  $G$  obtained from seven vertices is given below.




Figure: 2

**Subcase (i) (b):** If  $n$  is even then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by  
 $f(u_i) = 3$

$$f(u_i) = \begin{cases} 4i, & \forall i = 2, 4, 6, \dots, n \\ 4i-3 & \forall i = 3, 5, 7, \dots, n-1 \end{cases}$$

$$f(v_i) = 8i-6, \quad \forall i = 2, 3, 4, \dots, \frac{n}{2}$$

$$f(w_i) = 8i-4, \quad \forall i = 1, 2, 3, \dots, \frac{n}{2}$$

$$f(t_i) = 8i-2, \quad \forall i = 1, 2, 3, \dots, \frac{n}{2}$$

From the above labeling pattern, we get the edges labels are all distinct.

Thus  $f$  provide a Harmonic mean labeling for  $G$  the labeling pattern is shown in the following figure.




Figure: 3

**Case (ii):** If the triangular starts from  $u_2$  then also we consider two subcases

**Subcase (ii) (a):** If  $n$  is odd then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_1) = 1$$

$$f(u_2) = 4$$

$$f(u_i) = \begin{cases} 4i-3, & \forall i=3, 5, \dots, n. \\ 4i-6, & \forall i=4, 6, \dots, n-1. \end{cases}$$

$$f(v_1) = 2$$


$$f(v_i) = 8i-5, \quad \forall i=2, 3, \dots, \frac{n-1}{2}$$

$$f(w_i) = 8i-3, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}.$$

$$f(t_i) = 8i, \quad \forall i=2, 4, 6, \dots, \frac{n-1}{2}.$$

From above labeling pattern, we get distinct edge labels.

Thus  $f$  is Harmonic mean labeling of  $G$  and its labeling pattern of seven vertices is shown below.



**Figure: 4**

**Sub case (ii) (b):** If  $n$  is even then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_1) = 1$$

$$f(u_2) = 4$$

$$f(u_i) = \begin{cases} 4i-3, & \forall i=3, 5, \dots, n. \\ 4i-6, & \forall i=4, 6, \dots, n-1. \end{cases}$$

$$f(v_1) = 2$$

$$f(v_i) = 8i-5, \quad \forall i=2, 3, \dots, \frac{n-2}{2}$$

$$f(w_i) = 8i-3, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}.$$

$$f(t_i) = 8i, \quad \forall i=2, 4, 6, \dots, \frac{n-2}{2}.$$

Then we get distinct edge labels.

Thus  $f$  provide a Harmonic mean labeling of  $G$  and its labeling pattern of  $G$  is obtained eight vertices is shown below.




Figure: 5

From above all the cases, we conclude that Alternate Triple Triangular snakes are Harmonic mean graphs.

**Theorem 2.4:** Triple Quadrilateral snakes are Harmonic mean graphs.

**Proof:** Let  $G$  be the graph obtained from a path  $u_1 u_2 \dots u_n$  by joining  $u_i$  and  $u_{i+1}$  (alternatively) to new vertices  $v_i, w_i, x_i, y_i, z_i$  and  $t_i$  respectively and then joining  $v_i w_i, x_i y_i$  and  $z_i, t_i$   $1 \leq i \leq n-1$  and its labeling pattern is shown below




Figure: 6

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_1) = 4$$

$$f(u_i) = 10i-9, \quad 2 \leq i \leq n.$$

$$f(v_1) = 1$$

$$f(v_i) = 10i-8, \quad 1 \leq i \leq n-1$$

$$f(w_i) = 10i-3, \quad 1 \leq i \leq n-1.$$

$$f(x_i) = 10i-7, \quad 1 \leq i \leq n-1.$$

$$f(y_i) = 10i-1, \quad 1 \leq i \leq n-1.$$

$$f(z_i) = 10i-5, \quad 1 \leq i \leq n-1.$$

$$f(t_i) = 10i, \quad 1 \leq i \leq n-1.$$

From above labeling pattern, we get the edge labels are all distinct.

Thus  $f$  provide a Harmonic mean labeling for  $G$ .

**Example 2.5:** Harmonic mean labeling of  $G$  is obtained from five vertices is given below.




Figure: 7

**Theorem 2.6:** Alternate Triple Quadrilateral snakes are Harmonic mean graphs

**Proof:** Let  $G$  be an Alternate Triple Quadrilateral snake and its vertices be  $v_i, w_i, x_i, y_i, z_i$  and  $t_i$  ( $1 \leq i \leq n-1$ ).

**Case (i):** If the Quadrilateral starts from  $u_1$  then

We have two cases.

**Sub case (i)(a):** If  $n$  is odd then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by  
 $f(u_1) = 3$

$$f(u_i) = \begin{cases} \frac{11i}{2}, & \forall i = 2, 4, 6, \dots, n. \\ \frac{11(i-1)}{2} + 1, & \forall i = 3, 5, 7, \dots, n-1. \end{cases}$$

$$f(v_1) = 1$$

$$f(v_i) = 11i-9, \quad \forall i = 2, 3, 4, \dots, \frac{n-1}{2}$$

$$f(w_i) = 11i-5, \quad \forall i = 1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(x_i) = 11i-7, \quad \forall i = 1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(y_1) = 10$$

$$f(y_i) = 11i-2, \quad \forall i = 2, 3, 4, \dots, \frac{n-1}{2}$$

$$f(z_1) = 6$$

$$f(z_i) = 11i-5, \quad \forall i = 1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(t_1) = 8$$

$$f(t_i) = 11i-2, \quad \forall i = 2, 3, 4, \dots, \frac{n-1}{2}$$

From the above labeling pattern we get the edge labels are all distinct.  $G$  is a Harmonic mean graph.

Labeling pattern of  $G$  is obtained from seven vertices is shown in the following figure.




Figure: 8

**Subcase (i) (b):** If  $n$  is even then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by  
 $f(u_i) = 3$

$$f(u_i) = \begin{cases} \frac{11i}{2}, & \forall i = 2, 4, 6, \dots, n. \\ \left\lfloor \frac{11(i-1)}{2} \right\rfloor + 1, & \forall i = 3, 5, 7, \dots, n-1. \end{cases}$$

$$f(v_1) = 1$$

$$f(v_i) = 11i-9, \quad \forall i = 2, 3, 4, \dots, \frac{n}{2}$$

$$f(w_i) = 11i-5, \quad \forall i = 1, 2, 3, \dots, \frac{n}{2}$$

$$f(x_i) = 11i-7, \quad \forall i = 1, 2, 3, \dots, \frac{n}{2}$$

$$f(y_1) = 10$$

$$f(y_i) = 11i-2, \quad \forall i = 2, 3, 4, \dots, \frac{n}{2}$$

$$f(z_1) = 6$$

$$f(z_i) = 11i-5, \quad \forall i = 1, 2, 3, \dots, \frac{n}{2}$$

$$f(t_1) = 8$$

$$f(t_i) = 11i-2, \quad \forall i = 2, 3, 4, \dots, \frac{n}{2}$$

From the above labeling pattern we get the edge labels are all distinct.  $G$  is a Harmonic mean graph.

Labeling pattern of  $G$  is obtained from seven vertices is shown in the following figure.




Figure: 9

**Case (ii):** If the Quadrilateral starts from  $u_2$ .

Here we consider two subcases

**Subcase(ii)(a):** If n is odd then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_2) = 4$$

$$f(u_i) = \frac{11(i-1)}{2} + 1, \quad \forall i=1, 3, 5, \dots, n.$$

$$f(u_i) = \frac{11i}{2} - 9 \quad \forall i=4, 6, \dots, n-1.$$

$$f(v_1) = 2$$

$$f(v_i) = 11i - 8, \quad \forall i=2, 3, 4, \dots, \frac{n-1}{2}$$

$$f(w_i) = 11i - 3, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(x_i) = 11i - 6, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(y_i) = 11i - 1, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(z_i) = 11i - 4, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}$$

$$f(t_i) = 11i, \quad \forall i=1, 2, 3, \dots, \frac{n-1}{2}$$

From the above labeling pattern we get distinct edge labels.

Thus f provides a Harmonic mean labeling for G and it's labeling pattern obtained from seven vertices is given below.




Figure: 10

**Subcase (ii) (b):** If n is even then

Define a function  $f: V(G) \rightarrow \{1, 2, \dots, q+1\}$  by

$$f(u_2) = 4$$

$$f(u_i) = \frac{11(i-1)}{2} + 1, \quad \forall i=1, 3, 5, \dots, n.$$

$$f(u_i) = \frac{11i}{2} - 9 \quad \forall i=4, 6, \dots, n-1.$$

$$f(v_1) = 2$$

$$f(v_i) = 11i - 8, \quad \forall i=2, 3, 4, \dots, \frac{n-2}{2}$$

$$f(w_i) = 11i - 3, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}$$

$$f(x_i) = 11i - 6, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}$$

$$f(y_i) = 11i - 1, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}$$

$$f(z_i) = 11i - 4, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}$$

$$f(t_i) = 11i, \quad \forall i=1, 2, 3, \dots, \frac{n-2}{2}$$

From the above labeling pattern we get distinct edge labels.

Hence G is a Harmonic mean graph.

The labeling pattern of G is obtained from eight vertices is shown in the following figure




Figure: 11

From all the above cases it is clear that Alternate Triple Quadrilateral snake graphs are Harmonic mean graph.

## REFERENCES

1. J.A.Gallian, A dynamic survey of graph labeling. The Electronic Journal of combinatorics 17#DS6.
2. F.Harary, Graph theory, Narosa publishing House New Delhi.
3. S. Somasundram and R.Ponraj, Mean labeling of graphs, National Academy of Science letters vol.26, p210-2013.
4. S. Somasundaram and S.S.Sandhya Harmonic Mean Labeling of graphs, Communicated to Journal of Combinatorial Mathematics and Combinatorial Computing.
5. C. David Raj, C. Jayasekaran and S.S.Sandhya, 'Harmonic Mean Labeling On Double Triangular Snakes' Global Journal of Theoretical and Applied Mathematics Sciences Volume 3, Number 2 (2013) pp.67-72.
6. C. Jayasekaran, S.S.Sandhya and C. David Raj, 'Harmonic Mean Labeling On Double Quadrilateral Snakes' International Journal of Mathematics Research, Volume 5, Number 2 (2013), pp.251-256.

Source of support: Nil, Conflict of interest: None Declared

*[Copy right © 2015. This is an Open Access article distributed under the terms of the International Journal of Mathematical Archive (IJMA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]*